

UNIVERSITY OF CALIFORNIA
SANTA CRUZ

**AN INCLUSIVE SEARCH FOR THE DECAY OF A BOOSTED
HIGGS BOSON IN THE $H \rightarrow b\bar{b}$ CHANNEL WITH THE ATLAS
DETECTOR**

A dissertation submitted in partial satisfaction of the
requirements for the degree of

DOCTOR OF PHILOSOPHY

in

PARTICLE PHYSICS

by

Jacob Martin Pasner

October 2019

The Dissertation of Jacob Martin Pasner
is approved:

Professor Jason Nielsen, Chair

Professor Abraham Seiden

Professor Michael Hance

Dean Lori Kletzer
Vice Provost and Dean of Graduate Studies

Copyright © by
Jacob Martin Pasner
2019

Table of Contents

List of Figures	vi
List of Tables	vii
Abstract	viii
Dedication	ix
Acknowledgments	x
1 Introduction	1
I Theoretical Motivations and the Standard Model	2
2 The Standard Model and Beyond	3
2.1 The Standard Model	3
2.2 Quantum Chromodynamics	3
2.3 Quantum Electrodynamics	3
2.4 Spontaneous Symmetry Breaking	4
2.5 The Higgs Mechanism	4
2.6 Parton Distribution Function	4
II Experimental Apparatus and Associated Facilities	5
3 The Large Hadron Collider	6
3.1 Accelerator complex layout and design	7
3.2 Particle Injection Chain	7
3.3 Performance	7

4	The ATLAS Detector	8
4.1	Tracking with the Inner Detector	8
4.2	Calorimetry	8
4.3	Muon Spectrometer	8
5	Boosted Higgs at the LHC	9
5.1	Physics beyond the Standard Model	10
5.2	Higgs Production Mechanisms	10
5.3	Branching Ratios	10
5.4	Discovery	10
5.5	Fermion Decay Modes	10
5.6	Boosted Higgs	10
III	The HbbISR Analysis	11
6	Data and Simulation Preparation	12
6.1	Data Used	12
6.2	Monte Carlo Samples	12
7	Physics Object Selection	13
7.1	Calorimeter Jets	14
7.2	Track Jets	14
7.3	Fat Jets	14
7.4	B-tagged Jets	14
7.5	Muons	14
7.6	Overlap Removal	14
8	Event Selection	15
8.1	Selected Triggers	15
8.2	Pre-selection Studies	15
8.3	Signal Selection	15
8.4	Optimisation	15
9	Background Estimation	16
9.1	Multi-jet QCD estimation	16
9.2	$t\bar{t}$ control region	16
9.3	Single top estimation	16
9.4	Hadronic vector boson channel	16
10	Systematic Uncertainties	17
10.1	Theoretical Uncertainties	17
10.2	Experimental Uncertainties	17

11 Statistical Fit	18
11.1 Profile Likelihood Function	18
11.2 Fit Configuration	18
11.3 Statistical Tests	18
12 Results	19
12.1 Expectations	19
12.2 Statistical Analysis Results	19
12.3 Measurements and Limits	19
IV Conclusion	20
13 Conclusion	21
Bibliography	22
A Hadronic Vqq Sherpa Studies	23

List of Figures

List of Tables

Abstract

An Inclusive Search for the decay of a Boosted Higgs boson in the $H \rightarrow b\bar{b}$
channel with the ATLAS detector

by

Jacob Martin Pasner

This is an abstract placeholder

To my family

You never doubted me

And always made me laugh

Acknowledgments

I would like to thank my committee for their dutiful efforts to make this document one I can be proud of for the rest of my life. Furthermore, I would like to thank the SCIPP collaboration and UCSC Physics Department for their support in both academic and personal arenas.

Chapter 1

Introduction

Every dissertation should have an introduction. You might not realize it, but the introduction should introduce the concepts, background, and goals of the dissertation.

Part I

Theoretical Motivations and the Standard Model

Chapter 2

The Standard Model and Beyond

2.1 The Standard Model

The pinnacle of humanities ability to represent the fundamental fields and particles that build the universe, the Standard Model is the guiding theoretical basis of particle physics.

2.2 Quantum Chromodynamics

Quantum chromodynamics is super wack

2.3 Quantum Electrodynamics

Quantum Electrodynamics is the first model created in the QFT image.

2.4 Spontaneous Symmetry Breaking

Spontaneous symmetry breaking occurs when a system loses an inherent symmetry in order to attain a lower energy configuration.

2.5 The Higgs Mechanism

The Higgs Mechanism is the system by which particles attain mass through the spontaneous breaking of the Higgs potential, thus causing all particles it interacts with to have mass.

2.6 Parton Distribution Function

Before QFT the proton was thought to be a hard ball containing no smaller constituents. However, we know now that the strong field inside the proton allows for any strong object to exist with some probability which changes based off of the total energy of the proton. This behavior is represented then by a Probability Distribution Function.

Part II

Experimental Apparatus and Associated Facilities

Chapter 3

The Large Hadron Collider

Located 100 meters under the Swiss / French boarder lies the 26.7 kilometer Large Hadron Collider (LHC). The culmination of a huge international collaboration, this apparatus is used to produce proton and heavy ion collisions for observation by the four major experiments at the LHC: ATLAS, CMS, LHCb, and ALICE. The system was designed for a maximum center-of-mass energy of $\sqrt{s} = 14$ TeV and a peak instantaneous luminosity of $L = 10^{34}\text{cm}^{-2}\text{s}^{-1}$.

The first LHC workshop was held in 1984 in Lausanne at the European Organization for Nuclear Reserach (CERN). The nearly 30 year old case for a machine that would push towards the discovery of the elusive Higgs Boson was presented using the existing CERN accerlerator facilities and the Large Electron Positron (LEP) collider tunnel. The proposal became reality on September 10, 2008 when the first proton beams were circulated, only to have calamity strike 9 days later in the form of a catastrophic electrical fault. The repairs and improvements lasted until November 2009 when the

LHC restarted. Since then this modern marvel has worked wonderfully and, as hoped, lead to the discovery of the Higgs Boson by the CMS and ATLAS collaborations July 4, 2013.

The following chapter provides a brief introduction to the worlds most powerful accelerator starting with the little red bottle of hydrogen in building XXX, and ending with the interaction point where protons collide at the highest energies ever produced.

3.1 Accelerator complex layout and design

We begin with the most common element in the Universe, hydrogen, as our source of protons. A bottle of hydrogen gas provides raw H_2 which is then injected into a Duoplasmatron which uses a strong electric field to break the molecule into bare H^+ aka a proton! The proton is then accelerated by a 90 kV supply and leaves the Duoplasmatron with 1.4% speed of light (4000 km/s).

3.2 Particle Incjecton Chain

3.3 Performance

Chapter 4

The ATLAS Detector

4.1 Tracking with the Inner Detector

4.2 Calorimetry

4.3 Muon Spectrometer

Chapter 5

Boosted Higgs at the LHC

Its July 4th, 2012 and the walls of building 500 are reverberating as Particle Physicists around the world rejoice the discovery of the particle that gives all things mass, the Higgs Boson.

5.1 Physics beyond the Standard Model

5.2 Higgs Production Mechanisms

5.3 Branching Ratios

5.4 Discovery

5.5 Fermion Decay Modes

5.6 Boosted Higgs

Part III

The HbbISR Analysis

Chapter 6

Data and Simulation Preparation

In order to compare data to theory ATLAS has developed an analysis chain which runs both real data and simulated samples through the same processing, assuring a final result which is as comparable as possible.

6.1 Data Used

6.2 Monte Carlo Samples

Chapter 7

Physics Object Selection

After the ATHENA Digitization step both data and monte carlo have the same format, representing the three dimensional energy deposits. In order to analyze these deposits they are cleaned, clustered and checked for overlap resulting in physics objects useful for our specific analysis.

7.1 Calorimeter Jets

7.2 Track Jets

7.3 Fat Jets

7.4 B-tagged Jets

7.5 Muons

7.6 Overlap Removal

Chapter 8

Event Selection

Having created our physics objects we begin to make selections of what types of events we want to consider given the goal of our analysis. In our boosted topology this means considering things like momentum, jet collection efficiencies and background rejection.

8.1 Selected Triggers

8.2 Pre-selection Studies

8.3 Signal Selection

8.4 Optimisation

Chapter 9

Background Estimation

The dominant background was QCD. I worked on the $t\bar{t}$ control region.

The Vqq and single top backgrounds were estimated from monte carlo.

9.1 Multi-jet QCD estimation

9.2 $t\bar{t}$ control region

9.3 Single top estimation

9.4 Hadronic vector boson channel

Chapter 10

Systematic Uncertainties

10.1 Theoretical Uncertainties

10.2 Experimental Uncertainties

Chapter 11

Statistical Fit

The statistical fit in our analysis was accomplished using a framework developed for Higgs searches.

11.1 Profile Likelihood Function

11.2 Fit Configuration

11.3 Statistical Tests

Chapter 12

Results

12.1 Expectations

12.2 Statistical Analysis Results

12.3 Measurements and Limits

Part IV

Conclusion

Chapter 13

Conclusion

I conclude that this section is the conclusion

Bibliography

Appendix A

Hadronic Vqq Sherpa Studies

Ancillary material should be put in appendices, which appear after the bibliography.