Triangle centres in Lean 4: the Nagel line family and the symmedian and Brocard points

Josh Bald
Independent researcher, Ontario, Canada
jpbald93@gmail.com
ORCID: 0009-0002-1317-6489 *

27 October 2026

Abstract

We give machine-checked proofs, in Lean 4 with Mathlib, of classical theorems about two families of triangle centres. For the first family we define the Nagel and Gergonne points by their barycentric weights and prove that they are the common points of the cevians to the excircle and incircle contact points, where the contact points are Mathlib's `touchpoint`s. We prove the Nagel line relation $N - G = 2(G - I)$ with Mathlib's incentre and centroid, that the Spieker centre (the incentre of Mathlib's medial triangle) is the midpoint of I and N, that the Nagel point of the medial triangle is I, that the incircle and excircle contact points on each side are symmetric about its midpoint, that N is the incentre of the anticomplementary triangle, that the de Longchamps point lies on the Euler line and is the orthocentre of the anticomplementary triangle, and that the Spieker centre has equal power with respect to the three excircles. For the second family we prove that the point with barycentrics $a^2 : b^2 : c^2$ lies on the three symmedians (six angle equalities) and inside the triangle, that it is the centroid of its pedal triangle and, in the plane, the only such point, that the two Brocard points satisfy their angle equalities, that the Brocard angle ω satisfies $\cot\omega = \cot A + \cot B + \cot C = (a^2 + b^2 + c^2)/(4\Delta)$ and $0 < \omega \leq \pi/6$, that the Brocard points satisfy the three isogonality angle equalities and are equidistant from the circumcentre, that O, K and both Brocard points lie on the circle with diameter OK (which degenerates exactly for equilateral triangles), that the six points of Lemoine's first circle are concyclic, and the Droz-Farny line theorem. Apart from the pedal-centroid converse and Droz-Farny, no numbered theorem assumes that the ambient space is a plane. All proofs use only Lean's three standard axioms. In the sources we searched, subject to stated gaps, we found no earlier formal proof of these statements; one earlier Lean file proves an algebraic form of the Nagel and Gergonne cevian statements, and an earlier Coq development verifies barycentric forms of the Nagel line, the Spieker centre's position and two related facts.

1 Introduction

Triangle centres are among the most studied objects of elementary geometry; Kimberling's *Encyclopedia of Triangle Centers* [4] lists tens of thousands of them. A handful are classical, with theorems in every textbook [5, 6, 13]: the Gergonne point $X(7)$ and the Nagel point $X(8)$, where the cevians to the incircle and excircle contact points meet; the Spieker centre $X(10)$; the de Longchamps point

*© 2026 Josh Bald. Licensed under CC BY 4.0 (https://creativecommons.org/licenses/by/4.0/). The accompanying Lean code is licensed under the Apache License 2.0.

$X(20)$; the symmedian (Lemoine) point $X(6)$; and the two Brocard points. This paper formalises the basic theorems about these points in Lean 4 [1] with Mathlib [2].

Mathlib already contains a good part of the surrounding theory, and we build on it rather than redoing it. The Euler line is there: the Monge point of a simplex is $O + \frac{n+1}{n-1}(G - O)$ (`Affine.Simplex.mongePoint_eq_smul_vsub_vadd_circumcenter`), and for a triangle the orthocentre is $O + 3(G - O)$ (`Affine.Triangle.orthocenter_eq_smul_vsub_vadd_circumcenter`). Mathlib also has the incentre and excentres of a simplex, their radii and spheres, and the contact points `Affine.Simplex.touchpoint` (the orthogonal projections of the in- or excentre onto the faces), together with the medial simplex `Affine.Simplex.medial`. Our definitions of the Nagel and Gergonne points are by weights, but every theorem about them is stated with Mathlib's `incenter`, `excenter`, `touchpoint`, `medial`, `circumcenter` and `orthocenter`, so that the bridges between the weights and Mathlib's notions are part of what is proved. Euler's relation $OI^2 = R(R - 2r)$ is the subject of an open Mathlib pull request (#44230 [3], by another author); it is related work, not part of this paper, and we do not use it.

1.1 Contributions

The numbered theorems are proved for an arbitrary `Affine.Triangle` in a real inner product space of any dimension, with no hypothesis beyond the triangle itself, except where stated.

- *Nagel line family* (Section 3). Gergonne and Nagel concurrence through Mathlib's contact points (Theorems 1 and 2); the Nagel line $N - G = 2(G - I)$ (Theorem 3); the Spieker centre as the midpoint of IN (Theorem 4); the de Longchamps point (Theorem 5); the medial and anticomplementary triangles (Theorems 6 and 8); the isotomic contact points (Theorem 7); and the Spieker centre as the radical centre of the excircles, with the value of the common power (Theorem 9).
- *Symmedian and Brocard points* (Section 4). The six symmedian angle equalities and interiority (Theorem 10); the symmedian point as the centroid of its pedal triangle, and its uniqueness in the plane (Theorem 11); the two Brocard points (Theorem 12); the Brocard angle (Theorem 13); isogonality and equidistance from O (Theorem 14); the Brocard circle and its degenerate case (Theorems 15 and 16); Lemoine's first circle (Theorem 17); and the Droz-Farny line (Theorem 18).

The proofs are by coordinate certificates, as in our earlier papers [14, 15, 16, 17, 18, 19]: barycentric coordinates relative to the triangle, the Ravi substitution for the Nagel family, and inner products in the basis $B - A$, $C - A$ for the angle statements (Section 5). Table 1 lists the Lean name of every numbered theorem.

2 Setting and definitions

Throughout, V is a real inner product space, P a metric space that is a normed torsor over V, and t an `Affine.Triangle` in P, that is, three affinely independent points $A = t_0$, $B = t_1$, $C = t_2$. The dimension of V is arbitrary unless stated otherwise. We write

$$a = |BC|, \quad b = |CA|, \quad c = |AB|, \quad s = \tfrac{1}{2}(a + b + c),$$

G for the centroid (Mathlib's `Finset.centroid` of the vertices), I for `Affine.Simplex.incenter`, O for `circumcenter` and H for `orthocenter`. For weights w_0, w_1, w_2 with $w_0 + w_1 + w_2 = 1$ we

write $w_0A+w_1B+w_2C$ for Mathlib's `Finset.affineCombination`; all the points below are defined this way, by normalised barycentric coordinates. Because t is non-degenerate, $s-a$, $s-b$, $s-c$ and a, b, c are positive (the library proves $s-a>0$ etc. from Mathlib's positivity of the heights, through Heron's formula), so none of the normalisations divides by zero.

Mathlib's contact points. For a set S of vertex indices, Mathlib's `t.touchpoint S i` is the orthogonal projection of the (ex)centre with sign pattern S onto the face opposite vertex i. With $S=\emptyset$ this is the incentre, and `t.touchpoint ∅ i` is the contact point of the incircle with the side opposite vertex i. We prove that `t.excenter {i}` is the excentre opposite vertex i (its weights are $(-a,b,c)/(2(s-a))$ for $i=0$, and cyclically), so `t.touchpoint {i} i` is the contact point of the excircle opposite vertex i with the side opposite vertex i.

Definition 1 (Nagel family)**.** 1. The *Nagel point* is $N=\dfrac{(s-a)A+(s-b)B+(s-c)C}{s}$ (`Affine.Triangle.nagelPoint`).

2. The *Gergonne point* is the point with weights proportional to $\dfrac{1}{s-a}, \dfrac{1}{s-b}, \dfrac{1}{s-c}$, normalised to sum 1 (`Affine.Triangle.gergonnePoint`).

3. The *Spieker centre* is the incentre of Mathlib's medial triangle `Affine.Simplex.medial` (`Affine.Triangle.spiekerCenter`).

4. The *de Longchamps point* is $L=H+2(O-H)$, the reflection of H in O (`Affine.Triangle.deLongchampsPoint`).

5. The *anticomplementary triangle* is the image of t under the homothety with centre G and ratio -2 (`Affine.Triangle.anticomplementary`). It is defined as a value of type `Affine.Triangle` (via `Simplex.map` and the injectivity of the homothety), so its non-degeneracy is part of the construction; its vertices are $B+C-A$, $C+A-B$, $A+B-C$ (`NagelLine.anticomplementary_points_zero` and its two companions), and its medial triangle is t (`NagelLine.anticomplementary_medial`).

Definition 2 (symmedian and Brocard points)**.** 1. The *symmedian point* is $K=\dfrac{a^2A+b^2B+c^2C}{a^2+b^2+c^2}$ (`Affine.Triangle.symmedianPoint`).

2. The *first Brocard point* Ω has weights proportional to $\dfrac{1}{b^2}, \dfrac{1}{c^2}, \dfrac{1}{a^2}$ on A, B, C, normalised by $\dfrac{1}{a^2}+\dfrac{1}{b^2}+\dfrac{1}{c^2}$ (`Affine.Triangle.brocardPoint`$_1$).

3. The *second Brocard point* Ω' has weights proportional to $\dfrac{1}{c^2}, \dfrac{1}{a^2}, \dfrac{1}{b^2}$, with the same normalisation (`Affine.Triangle.brocardPoint`$_2$).

4. The *area* is $\Delta=\frac{1}{2}\,|AB|\,|AC|\sin\angle BAC$ (`Affine.Triangle.area`), with Mathlib's unoriented angle `EuclideanGeometry.angle`. The library also proves the Gram form $\Delta=\frac{1}{2}\sqrt{|u|^2|v|^2-\langle u,v\rangle^2}$ with $u=B-A$, $v=C-A$ (`Brocard.area_eq_sqrt`) and Heron's form in a^2, b^2, c^2 (`Brocard.area_eq_sqrt_dist`).

5. The six *Lemoine points* (`Affine.Triangle.lemoinePoints`): with $S=a^2+b^2+c^2$, they are the points of the side lines with parameters $C+\frac{a^2}{S}(A-C)$, $A+\frac{b^2+c^2}{S}(B-A)$, $B+\frac{c^2+a^2}{S}(C-B)$,

Table 1: Main theorems and their Lean names. Namespaces: `NagelLine` (development `paperg_nagel`) and `Brocard` (development `paperg_brocard`). Starred rows are from the extension files (`Extensions.lean`, and `Inradius.lean` for the inradius form in Theorem 9). The hypothesis column lists hypotheses beyond the triangle.

Thm	content	Lean	hyp.
1	Gergonne cevians	`gergonnePoint_mem_cevian`, `gergonne_cevians_concurrent`	–
2	Nagel cevians	`nagelPoint_mem_cevian`, `nagel_cevians_concurrent`	–
3	$N - G = 2(G - I)$	`nagelPoint_vsub_centroid`, `collinear_incenter_centroid_nagelPoint`	–
4	Spieker $= \mathrm{mid}(I, N)$	`medial_incenter_eq_midpoint`, `collinear_spiekerCenter_incenter_nagelPoint`	–
5*	de Longchamps	`midpoint_orthocenter_deLongchampsPoint`, `collinear_deLongchampsPoint_circumcenter_orthocenter`, `collinear_centroid_circumcenter_deLongchampsPoint`, `anticomplementary_orthocenter`	–
6*	medial Nagel $= I$	`medial_nagelPoint`	–
7*	isotomic contacts	`midpoint_touchpoint_zero`, `_one`, `_two`	–
8*	$N =$ anticompl. incentre	`nagelPoint_eq_anticomplementary_incenter`	–
9*	Spieker radical centre	`spiekerCenter_power_exsphere`, `spiekerCenter_power_exsphere_eq`, `spiekerCenter_power_exsphere_eq_inradius`	–
10	symmedians, interior	`symmedian_angles`, `symmedianPoint_mem_interior`	–
11	pedal centroid; converse	`symmedianPoint_eq_centroid_pedal`, `eq_symmedianPoint_of_eq_centroid_pedal`	(ii) dim 2
12	Brocard points	`brocardPoint₁_angles`, `brocardPoint₂_angles`	–
13	Brocard angle	`cot_brocardAngle_eq_sum_cot`, `cot_brocardAngle_eq_div_area`, `brocardAngle_pos_le`	–
14*	isogonal, $\lvert O\Omega\rvert = \lvert O\Omega'\rvert$	`brocardPoints_isogonal`, `dist_circumcenter_brocardPoint`	–
15*	Brocard circle	`brocardCircle`, `cospherical_brocardCircle`, `brocardCircle_right_angles`	–
16*	equilateral case	`brocardCircle_degenerate`, `brocard_eq_circumcenter_of_equilateral`	see text
17*	Lemoine's first circle	`lemoineCircle`, `cospherical_lemoinePoints`	–
18*	Droz-Farny	`drozFarny`	see text

$A + \frac{b^2}{S}(B - A)$, $B + \frac{c^2}{S}(C - B)$, $C + \frac{a^2+b^2}{S}(A - C)$. The bridge `Brocard.lemoinePoints_mem` proves that each lies on its side line and on the line through K parallel to another side (the parallel to BC gives points 0 and 1, the parallel to CA points 2 and 3, the parallel to AB points 4 and 5), so they are the classical six points.

The Brocard weight order is easy to get wrong; the builders checked numerically (30 digits) that $(1/b^2, 1/c^2, 1/a^2)$ gives $\angle\Omega AB = \angle\Omega BC = \angle\Omega CA$ and the other cyclic order gives the mirror point, and Theorem 12 proves it. In Lean, $\angle XYZ$ is `∠ X Y Z`, the angle at Y, with values in $[0, \pi]$; by Mathlib's convention the angle is $\pi/2$ when $X = Y$ or $Z = Y$.

3 The Nagel line family

The theorems of this section are in the development `paperg_nagel` (library `NagelLine`); each has the triangle t as its only hypothesis. File:line references are to the library files at the commit given in Section 11.

Theorem 1 (Gergonne point)**.** *For every triangle and every vertex i, the Gergonne point lies on the line through vertex i and the contact point of the incircle with the opposite side (Mathlib's `t.touchpoint ∅ i`). Consequently the three cevians to the incircle contact points have a common point.*

Lean: `NagelLine.gergonnePoint_mem_cevian` (`Main`:290) and `NagelLine.gergonne_cevians_concurrent` (`Main`:328).

Theorem 2 (Nagel point)**.** *For every triangle and every vertex i, the Nagel point lies on the line through vertex i and the contact point of the excircle opposite vertex i with the side opposite vertex i (Mathlib's `t.touchpoint {i} i`). Consequently the three cevians to the excircle contact points have a common point. Moreover each such contact point lies strictly between the two other vertices.*

Lean: `NagelLine.nagelPoint_mem_cevian` (`Main`:334), `NagelLine.nagel_cevians_concurrent` (`Main`:357); the last sentence is `NagelLine.sbtw_touchpoint_singleton_zero` and its two companions (`Main`:170–188).

Theorem 3 (Nagel line)**.** *For every triangle, $N - G = 2(G - I)$, where I is Mathlib's incentre. In particular I, G, N are collinear.*

Lean: `NagelLine.nagelPoint_vsub_centroid` (`Main`:362) and `NagelLine.collinear_incenter_centroid_nagelPoint` (`Main`:373).

Theorem 4 (Spieker centre)**.** *For every triangle, the incentre of the medial triangle is the midpoint of I and N. In particular the Spieker centre, I and N are collinear.*

Lean: `NagelLine.medial_incenter_eq_midpoint` (`Main`:385) and `NagelLine.collinear_spiekerCenter_incenter_nagelPoint` (`Main`:408).

Theorem 5 (de Longchamps point)**.** *For every triangle, with L the de Longchamps point: (i) O is the midpoint of H and L; (ii) L, O, H are collinear; (iii) G, O, L are collinear; (iv) L is the orthocentre of the anticomplementary triangle.*

Lean: `NagelLine.midpoint_orthocenter_deLongchampsPoint`, `NagelLine.collinear_deLongchampsPoint_circumcenter_orthocenter`, `NagelLine.collinear_centroid_circumcenter_deLongchampsPoint`, `NagelLine.anticomplementary_orthocenter` (`Extensions`:228, 237, 249, 261). Parts (i) and (ii) are immediate from the definition; (iii) uses Mathlib's Euler line, and (iv) uses that the circumcentre of the anticomplementary triangle is H (`NagelLine.anticomplementary_circumcenter`).

Theorem 6 (Nagel point of the medial triangle)**.** *For every triangle, the Nagel point of Mathlib's medial triangle is the incentre I of the triangle.*

Lean: `NagelLine.medial_nagelPoint` (`Extensions`:274). (The statement is written `Triangle.nagelPoint t.medial`, because `Simplex.medial` returns a `Simplex ℝ P 2` and field notation would look for a `Simplex` constant; the term is the same.)

Theorem 7 (isotomic contact points)**.** *For every triangle and each side, the midpoint of the incircle contact point and the contact point of the opposite excircle on that side is the midpoint of the side; for the side BC,* $\mathrm{mid}(\texttt{touchpoint}\ \emptyset\ 0,\ \texttt{touchpoint}\ \{0\}\ 0) = \mathrm{mid}(B, C)$*, and likewise for CA and AB.*

Lean: `NagelLine.midpoint_touchpoint_zero`, `NagelLine.midpoint_touchpoint_one`, `NagelLine.midpoint_touchpoint_two` (`Extensions`:297, 308, 319).

Theorem 8 (anticomplementary triangle)**.** *For every triangle, the Nagel point is the incentre of the anticomplementary triangle.*

Lean: `NagelLine.nagelPoint_eq_anticomplementary_incenter` (`Extensions`:332).

Theorem 9 (Spieker centre as radical centre)**.** *For every triangle and all $i, j \in \{0, 1, 2\}$, the power of the Spieker centre S_p with respect to the excircle opposite vertex i equals its power with respect to the excircle opposite vertex j: $|S_pJ_i|^2 - r_i^2 = |S_pJ_j|^2 - r_j^2$, where J_i and r_i are the centre and radius of Mathlib's `t.exsphere {i}`. The common value is*

$$\frac{1}{4}\Big(s^2 + \frac{(s-a)(s-b)(s-c)}{s}\Big) = \frac{s^2 + r^2}{4},$$

with r the inradius.

Lean: `NagelLine.spiekerCenter_power_exsphere` (`Extensions`:357) and, for the value, `NagelLine.spiekerCenter_power_exsphere_eq` (`Extensions`:201) for the left-hand form; `NagelLine.inradius_sq` (Mathlib's `t.inradius` satisfies $r^2 = (s-a)(s-b)(s-c)/s$) and `NagelLine.spiekerCenter_power_exsphere_eq_inradius` for the right-hand form (`Inradius`:19, line 40). The last two were added after the Layer-13 comparison found the right-hand form missing from the library (Section 8).

4 The symmedian point and the Brocard points

The theorems of this section are in the development `paperg_brocard` (library `Brocard`). The only hypothesis is the triangle t, except in Theorem 11(ii) and Theorem 18, where it is stated.

Theorem 10 (symmedians)**.** *For every triangle, with K the symmedian point and M_a, M_b, M_c the midpoints of BC, CA, AB:*

$$\begin{aligned} \angle BAK = \angle M_aAC, &\quad \angle CAK = \angle M_aAB, \\ \angle CBK = \angle M_bBA, &\quad \angle ABK = \angle M_bBC, \\ \angle ACK = \angle M_cCB, &\quad \angle BCK = \angle M_cCA. \end{aligned}$$

So at each vertex the line to K is the reflection of the median in the angle bisector, and the three symmedians pass through K. Moreover K is an affine combination of A, B, C with positive weights, that is, it lies in the relative interior of the triangle (its interior within the plane of the triangle).

Lean: `Brocard.symmedian_angles` (`Main`:22) and `Brocard.symmedianPoint_mem_interior` (`Main`:100).

Theorem 11 (pedal triangle)**.** *(i) For every triangle, K is the centroid of its pedal triangle: if F_a, F_b, F_c are the orthogonal projections of K onto the lines BC, CA, AB (Mathlib's `EuclideanGeometry.orthogonalProjection` onto `line[ℝ, B, C]` etc.), then $\frac{1}{3}(F_a + F_b + F_c) = K$. (ii) If moreover $\dim V = 2$, then every point X of the plane that is the centroid of its own pedal triangle is K.*

Lean: `Brocard.symmedianPoint_eq_centroid_pedal` (`Main`:115) and `Brocard.eq_symmedianPoint_of_eq_centroid_pedal` (`Main`:150), the latter with the instance hypothesis `[Fact (Module.finrank ℝ V = 2)]`.

Theorem 12 (Brocard points)**.** *For every triangle, the first Brocard point satisfies $\angle\Omega AB = \angle\Omega BC = \angle\Omega CA$, and the second satisfies $\angle\Omega' BA = \angle\Omega' CB = \angle\Omega' AC$.*

Lean: `Brocard.brocardPoint₁_angles` (`Main`:233) and `Brocard.brocardPoint₂_angles` (`Main`:280).

Theorem 13 (Brocard angle)**.** *For every triangle, with $\omega = \angle\Omega AB$ and the vertex angles $\angle BAC$, $\angle ABC$, $\angle ACB$:*

$$\cot\omega = \cot\angle BAC + \cot\angle ABC + \cot\angle ACB = \frac{a^2+b^2+c^2}{4\Delta}, \qquad 0 < \omega \le \frac{\pi}{6}.$$

Lean: `Brocard.cot_brocardAngle_eq_sum_cot`, `Brocard.cot_brocardAngle_eq_div_area`, `Brocard.brocardAngle_pos_le` (`Main`:327, 366, 402). Here cot is Mathlib's `Real.cot`.

Theorem 14 (isogonal Brocard points)**.** *For every triangle, $\angle\Omega AB = \angle\Omega' AC$, $\angle\Omega BC = \angle\Omega' BA$ and $\angle\Omega CA = \angle\Omega' CB$; and $|O\Omega| = |O\Omega'|$.*

Lean: `Brocard.brocardPoints_isogonal` (`Extensions`:21) and `Brocard.dist_circumcenter_brocardPoint` (`Extensions`:85). "Isogonal" here means exactly these three angle equalities; no isogonal-conjugation map is defined. Together with Theorem 12 they say that the two Brocard angles are equal.

Theorem 15 (Brocard circle)**.** *For every triangle, each of O, K, Ω, Ω' is at distance $\frac{1}{2}|OK|$ from the midpoint of OK; so the four points are cospherical (Mathlib's `EuclideanGeometry.Cospherical`). Moreover $\langle O-\Omega, K-\Omega\rangle = 0$ and $\langle O-\Omega', K-\Omega'\rangle = 0$, and $\angle O\Omega K = \angle O\Omega' K = \pi/2$.*

Lean: `Brocard.brocardCircle`, `Brocard.cospherical_brocardCircle`, `Brocard.brocardCircle_right_angles` (`Extensions`:116, 175, 183).

Theorem 16 (degenerate Brocard circle)**.** *For every triangle, $\Omega = O$ or $\Omega = K$ or $\Omega' = O$ or $\Omega' = K$ or $K = O$ holds if and only if the triangle is equilateral; and if the triangle is equilateral then $\Omega = \Omega' = K = O$.*

Lean: `Brocard.brocardCircle_degenerate` (`Extensions`:231) and `Brocard.brocard_eq_circumcenter_of_equilateral` (`Extensions`:314), the latter with the two hypotheses $|AB| = |BC|$ and $|BC| = |CA|$.

Theorem 17 (Lemoine's first circle)**.** *For every triangle, the six Lemoine points (Definition 2) are all at the same distance from the midpoint of OK; in particular they are cospherical.*

Lean: `Brocard.lemoineCircle` (`Extensions`:353) and `Brocard.cospherical_lemoinePoints` (`Extensions`:439).

Theorem 18 (Droz-Farny)**.** *Let $\dim V = 2$, let $d_1, d_2 \in V$ with $\langle d_1, d_2\rangle = 0$, and suppose that neither d_1 nor d_2 lies in the direction of any of the three side lines. For $k = 1, 2, 3$ let X_k and Y_k be points of the k-th side line (BC, CA, AB) lying on the line through H with direction d_1 and d_2 respectively. Then the midpoints of X_1Y_1, X_2Y_2, X_3Y_3 are collinear.*

Lean: `Brocard.drozFarny` (`Extensions`:449); the lines through H are `AffineSubspace.mk' t.orthocenter (ℝ · d_i)`. The non-direction hypotheses exclude a line through H that is parallel to a side; they also imply $d_1, d_2 \neq 0$.

5 Proof method and bridges to Mathlib

Barycentric certificates. Every point in the statements is, or is proved equal to, an affine combination of A, B, C with explicit weights that are rational functions of a, b, c (Nagel family) or of a^2, b^2, c^2 (symmedian and Brocard family). Membership of a point in a line, equality of two points, a midpoint or a collinearity then reduces to an identity between weights, which is closed by `field_simp` and `ring`, or by Mathlib's `match_scalars` where the two sides are combinations of the same vectors. As in our earlier papers [14, 18], the identities and, where needed, their cofactors were found with SymPy and checked numerically (mpmath, 30 digits, random triangles) before the Lean proofs were written; the numerical checks are evidence, not part of the proofs.

The Ravi substitution. For the Nagel family the library proves that there are $x, y, z > 0$ with $a = y+z$, $b = z+x$, $c = x+y$ and $s = x+y+z$ (lemma `ravi`, from $s-a, s-b, s-c > 0$). After the substitution the weights $(s-a)/s$, $1/(s-a)$, $a/(2s)$, $(s-b)/a$ and so on become rational functions of x, y, z with denominators that are visibly positive, so `field_simp` can clear them without case analysis. The positivity $s - a > 0$ itself is derived from Mathlib's `Affine.Simplex.height_pos` through the squared heights and a factorisation of Heron's polynomial (`NagelLine.s_sub_a_pos` and its companions).

The touchpoint bridges. The connection with Mathlib's contact points is made in two steps. First, Mathlib defines the in- and excentres by weights (`excenterWeights`); we compute them: the incentre has weights $(a, b, c)/(2s)$ (`NagelLine.incenter_weights`) and `excenter {i}` has weights $\pm a_j/(2(s - a_i))$ with the minus sign at $j = i$ (`NagelLine.excenter_weights`), which identifies it as the excentre opposite vertex i. Second, since `touchpoint` is an orthogonal projection onto a face, a candidate point on the side line is the contact point exactly when the vector from the centre to it is orthogonal to the side (Mathlib's `EuclideanGeometry.coe_orthogonalProjection_eq_iff_mem`). With the candidate $B + \frac{s-b}{a}(C - B)$ for the incircle on BC, and $B + \frac{s-c}{a}(C - B)$ for the A-excircle, the orthogonality is a polynomial identity in the Gram entries $|B - A|^2 = c^2$, $|C - A|^2 = b^2$, $\langle B - A, C - A\rangle = \frac{1}{2}(b^2 + c^2 - a^2)$. This gives `NagelLine.touchpoint_empty_zero`, `NagelLine.touchpoint_singleton_zero` and their companions, after which Theorems 1–7 are weight computations. The medial triangle's sides are half the original ones (`NagelLine.medial_sideA` etc.), which transfers the weight formulas to `t.medial`. The anticomplementary triangle is a homothetic image of t, so its centroid, its medial triangle and its circumcentre come from Mathlib's lemmas about `map` and from `EuclideanGeometry.eq_circumcenter_of_dist_eq`; the radical-centre identity uses Mathlib's formula for the exradius as the inverse of the sum of unnormalised weights.

Angles through inner products. For the second family all points are written in the basis $u = B - A$, $v = C - A$ (which need not span V; nothing beyond the plane of the triangle is used), so every inner product is a polynomial in the coordinates and the Gram entries above, whose determinant is positive (`Brocard.gram_det_pos`). An equality of unoriented angles $\angle XYZ = \angle X'Y'Z'$ is reduced to: the squared cosines are equal, and the two cosine numerators have the same sign (`Brocard.angle_eq_angle_of_inner`, `Brocard.angle_eq_of_coords`). The first is a rational identity; for the second the builders found, with SymPy, an expression of the product of the two numerators as a square times a manifestly positive factor. The cotangent of the angle between vectors X, Y is $\langle X, Y\rangle / \sqrt{|X|^2|Y|^2 - \langle X, Y\rangle^2}$, and for points in coordinates the square root becomes an absolute 2×2 determinant times $\sqrt{|u|^2|v|^2 - \langle u, v\rangle^2}$ (`Brocard.cot_angle_eq_inner`,

`Brocard.cot_angle_coords`), which turns Theorem 13 into rational identities in a^2, b^2, c^2 and Δ. The feet of the perpendiculars are explicit (`Brocard.orthogonalProjection_vsub`: Mathlib's projection onto `line[ℝ, Y, Z]` is the explicit foot); for K the foot on AB is $A + t(B - A)$ with $t = (-a^2+3b^2+c^2)/(2S)$ with $S = a^2+b^2+c^2$. The converse in Theorem 11 is a linear system closed with `linear_combination` and explicit cofactors. For the Brocard circle and the Lemoine circle the circumcentre and orthocentre are also written in coordinates (`Brocard.circumcenter_coords`, `Brocard.orthocenter_coords`). In Droz-Farny the hypothesis $\dim V = 2$ is used to write d_1, d_2 in the basis u, v; the collinearity determinant factors as a non-vanishing rational factor times a polynomial Q, and Q is a multiple of $\langle d_1, d_2 \rangle$.

6 Degenerate cases and hypotheses

No dimension hypothesis. Among the numbered theorems, apart from two results, every theorem holds for a triangle in a real inner product space of any dimension, and carries no `Module.finrank` hypothesis: all the points involved lie in the plane of the triangle, and the proofs use only coordinates in the basis $B - A$, $C - A$. The exceptions are Theorem 11(ii), where X ranges over the whole space and the proof needs X in the plane of the triangle to write it in coordinates (we have not formalised the higher-dimensional version); and Theorem 18, where the directions d_1, d_2 must be written in the basis. Both state `[Fact (Module.finrank ℝ V = 2)]` explicitly, and the tampering suite checks that removing it from the pedal converse breaks the build. Non-degeneracy of the triangle is never a separate hypothesis: it is part of `Affine.Triangle`.

The equilateral Brocard circle. Theorem 15 has no hypothesis excluding coincidences. When Ω coincides with O or K, the right-angle statement $\angle O\Omega K = \pi/2$ holds only because Mathlib defines the angle at a vertex coinciding with an endpoint to be $\pi/2$; the inner-product form $\langle O - \Omega, K - \Omega \rangle = 0$ is then trivially true. Theorem 16 says exactly when this happens: for equilateral triangles only, and then all four points coincide and the "circle" has radius 0. The factor behind both statements is $a^4 + b^4 + c^4 - a^2b^2 - b^2c^2 - c^2a^2$, which appears in each of the squared distances. For non-equilateral triangles the right-angle statement is the classical one, and the Layer-14 witness of Section 8 exhibits a triangle in which O, K, Ω, Ω' are pairwise distinct.

Other edge cases. The excircle contact points lie on the open sides (Theorem 2), so the Nagel cevians are genuine cevians. The Brocard angle bound $\omega \leq \pi/6$ is attained only in the equilateral case; the library proves the non-strict bound, and a strict instance is checked only on the witness triangle. For right triangles H is a vertex; the de Longchamps theorems need no exclusion, while in Droz-Farny a line through H could coincide with a side, which the non-direction hypotheses exclude. Under Lean's convention $x/0 = 0$ some expressions would be defined even for degenerate data, but no theorem here relies on it: every denominator (s, $s - a$, $a^2 + b^2 + c^2$, a, Δ, the Gram determinant) is proved positive.

7 Related work and prior art

Mathlib. At the pinned revision Mathlib has the Euler line (through the Monge point), the nine-point circle and its centre on the Euler line, the in- and excentres of a simplex with their radii, spheres and contact points, the medial simplex, and Ceva's theorem in affine spaces. We found no Mathlib declaration and no Mathlib pull request mentioning the Nagel, Gergonne, Spieker, de Longchamps, symmedian (Lemoine) or Brocard points, isogonal conjugation, the Droz-Farny line

or Lemoine's circle. Euler's $OI^2 = R(R - 2r)$, with excentre versions and $r \leq R/2$, is in the open Mathlib pull request #44230 [3] (opened 2026-09-26); a proof also appears in the repository `facebookresearch/atlas-lean`. We neither use nor reprove it.

Earlier Lean work on these centres. The only related Lean file we found is in the `lean-genius` repository [7] (file `CevasTheoremOQ04OQ04OQ02.lean`, first committed 2026-06-25, no `sorry`). It defines the incentre, the excentres, the Nagel and Gergonne points and the contact feet by Ravi-barycentric formulas in an abstract real module and proves that the defined point lies on the cevian to a foot that is itself defined by a formula. It is a partial, algebraic form of the cevian statements of Theorems 1 and 2: the feet are not tied to incircle or excircle tangency and the incentre is not tied to Mathlib's, and there is no Nagel line or ratio statement. We cite it as partial prior art. We found no Lean formalisation of the other theorems of this paper; for algebraic Coq forms of some of them see the next paragraph.

Other systems. In GeoCoq and the Coq high-school geometry library the Euler line is formalised; we found no Nagel, Gergonne, Spieker, symmedian, Brocard, Lemoine-circle or Droz-Farny results there, nor in the Archive of Formal Proofs, the HOL Light distribution or the Mizar Mathematical Library. Closest to this paper is the certified version of Kimberling's encyclopedia by Narboux and Braun [10], in Coq. There, triangle centres are defined by their barycentric or trilinear coordinates, and transformations such as the complement and the isogonal conjugate by formulas; facts are proved by field normalisation. Its tables list as Coq-verified that $X(1)$ is the complement of $X(8)$ and $X(10)$ the complement of $X(1)$ (the Nagel line with its ratio, and the Spieker centre on it), that $X(4)$ is the complement of $X(20)$, and that $X(6)$ is the isogonal conjugate of $X(2)$. These are algebraic forms of parts of Theorems 3, 4, 5 and 10: the points are not tied to incircle or excircle contact, to an incentre defined by equal distances, or to Euclidean angles, which is what our statements add. We found no Brocard-point, Brocard-angle, Lemoine-circle or Droz-Farny results there. All Lean hits for "brocard" that we found concern Brocard's number-theoretic problem $n! + 1 = m^2$, not the Brocard points.

Automated geometry theorem proving. The identities behind Theorems 1–9 and 17–18 are of the kind that algebraic provers (Wu's method and Gröbner bases [11]) and the area method [12] handle routinely, in coordinates and with non-degeneracy conditions generated by the method. Our statements differ in what they are about: the incentre, contact points, excircles, centroid, orthocentre, circumcentre and unoriented angles are Mathlib's own objects, the non-degeneracy hypothesis is only that the three points form a triangle, and the proofs are checked by the Lean kernel and connected to the library's definitions by the bridge lemmas of Section 5.

How the search was done, and its gaps. The sweep (2026-10-24) grepped a local Mathlib checkout at the pinned revision, searched mathlib4 pull requests and GitHub Lean code through the GitHub API, grepped clones of `lean-genius` and `atlas-lean`, ran per-repository searches of FormalBook, Compfiles and other benchmark collections, and searched local copies of GeoCoq, the Coq high-school geometry library, HOL Light and the Mizar library, and the Archive of Formal Proofs through a GitHub mirror. The query terms were the names of the centres and theorems (`nagel`, `gergonne`, `symmedian`, `lemoine`, `brocard`, `spieker`, `longchamps`, `isogonal`, `droz`, and others). General web search was unavailable during that sweep. A second sweep on 2026-10-27 repeated the GitHub searches at the then-current heads of Mathlib, `atlas-lean`, `formal-conjectures`, Compfiles and `lean-genius`, searched mathlib4 pull requests opened or merged since 2026-10-01,

and added general web search, which found the Coq work [10]. Remaining gaps: ProofAtlas was checked only through its sitemap (752 URLs), not by reading its pages; the certified encyclopedia's website was available only through archived copies, so its Coq sources and collinearity lists could not be inspected; and GitHub does not appear to index Mizar files, so the local Mizar search stands. Our statements of novelty are subject to these gaps.

Classical sources. The theorems are classical; we follow Johnson [5], Coxeter and Greitzer [6] and Honsberger [13], and Kimberling's encyclopedia [4] for the centre numbers and barycentric coordinates. For the three less familiar statements: the Spieker centre is the radical centre of the excircles in Yiu [8, §7.3.3, p. 94]; the first Lemoine circle and its centre at the midpoint of KO are in Honsberger [13, Ch. 9, §5, pp. 94–95]; the Droz-Farny theorem, stated by Droz-Farny in 1899, is proved synthetically by Ayme [9]. Our earlier papers [14, 15, 16, 17, 18, 19] use the same certificate method for other configurations of classical plane geometry.

8 Verification

The checks follow our earlier papers [14, 19]; for a self-contained reading, they are:

gate	script that rejects forbidden tokens, builds, and checks that every required declaration uses only the standard axioms
tampering suite	planted defects (hidden `sorry`, extra axiom, fake axiom report, removed hypothesis, ...) that the gate must reject
battery	per-theorem mutation of conclusions (each mutant must fail to compile; a heartbeat limit counts as a rejection), a contradiction probe on the hypotheses, a satisfiability certificate (a local language model proposes concrete values; only a successful Lean compilation counts), hypothesis ablation, and the unused-variable linter
Layer 13	blind restatement of the theorems from this paper's prose by a fresh writer, then a two-way comparison with the library statements
Layer 14	restatement of the main theorems in Mathlib-only terms, with explicit witness triangles
Layer 15	statements written from classical sources by a writer who saw neither paper nor code, then proved from the library

The two machines are an x86-64 Linux machine and an arm64 macOS machine.

Gates and tampering. As in our earlier papers, each development has a gate script (`gate.sh`) that rejects forbidden tokens (`sorry`, `admit`, `native_decide`, `axiom` declarations and the like), builds the library, and checks with `#print axioms` that every declaration in its `REQUIRED` list depends only on `propext`, `Classical.choice` and `Quot.sound`. The `REQUIRED` lists contain every definition, main theorem and bridge lemma. `paperg_nagel` passed with 48 declarations (the files `Basic.lean` and `Main.lean`, Theorems 1–4) and then with 77 after `Extensions.lean` (Theorems 5–9) was added; `paperg_brocard` passed with 38 (Theorems 10–13) and then 67 after `ExtLemmas.lean` and `Extensions.lean` (Theorems 14–18). The extensions added library files, import lines, `REQUIRED` entries and one tampering test, and changed no earlier statement (the development history is not published; see the integration reports in the supporting material). The tampering suite (`tests/tamper.sh`) plants a hidden `sorry`, an axiom split over two lines, a fake `#print axioms` report, a command-redefining macro, a wrapper axiom, and the removal of a needed hypothesis (in `paperg_nagel` the triangle binder of `NagelLine.nagelPoint_vsub_centroid` and

of `NagelLine.spiekerCenter_power_exsphere`; in `paperg_brocard` the dimension hypothesis of `Brocard.eq_symmedianPoint_of_eq_centroid_pedal` and the orthogonality hypothesis of `Brocard.drozFarny`); every planted defect was rejected at each stage (7 of 7 in each final suite). In an early version of `paperg_nagel` the removed triangle binder was silently re-supplied by a section `variable`; the tampering suite caught this, and the main theorems were moved out of that section.

Cross-machine reproduction. `paperg_brocard` was built on an arm64 macOS machine from a copy of the x86-64 Linux repository at the same local commit; the build succeeded, the gate printed `PASS (67 declarations, standard axioms only)` and the tampering suite rejected all 7 defects. The SHA-256 hashes of all library `.lean` files of both developments agree between the two machines.

Fresh-clone receipts. Both repositories were cloned from GitHub at the published commits and built on both machines (x86-64 Linux and arm64 macOS, Lean v4.33.1, the pinned Mathlib). On each machine `paperg_nagel` printed `PASS (151 declarations, standard axioms only)` and `paperg_brocard` printed `PASS (147 declarations, standard axioms only)`; the tampering suites printed `ALL TAMPERING TESTS REJECTED`; and the SHA-256 digest of the sorted library `.lean` file hashes agreed between the machines (`201feb97...` and `bd4e777b...`). The four receipts are in the supporting material.

Statement-level battery. The battery of our earlier papers was run on each development at each stage. The figures are per development; every run's report is in the supporting material (`reports/battery/`, file names give the machine and the start time). Intermediate runs after the Layer-14 files were added (140 and 109 theorems) are also there but not described below; some early runs were on working trees with uncommitted changes, as their reports record.

- *`paperg_nagel`, before the extensions* (78 theorems): conclusion mutations 210/210 rejected (7 of them by the heartbeat limit), none surviving; contradiction probe: no contradiction in 12 probed declarations (66 have no hypotheses and were not probed); satisfiability certified for 2 of 78, 76 unresolved; hypothesis ablation: 29 binders needed, 1 unused; linter: no warnings in 2 library files.

- *`paperg_nagel`, with the extensions* (105 theorems): mutations 282/282 rejected (7 by the heartbeat limit), none surviving; no contradiction in 14 probed declarations (91 without hypotheses); satisfiability certified for 4 of 105, 101 unresolved; ablation: 32 needed, 1 unused; linter: no warnings in 3 files (run on the arm64 machine). A run on the x86-64 machine gave the same mutation, contradiction and satisfiability figures; its ablation stage reported 41 binders needed and 1 unused. The difference in ablation scope is not explained by the reports; the arm64 run was on a working tree with uncommitted changes (11 entries), so the two trees may have differed.

- *`paperg_brocard`, before the extensions* (40 theorems): mutations 97/97 rejected, none by timeout; no contradiction in 13 probed declarations (27 without hypotheses); satisfiability certified for 8 of 40, 32 unresolved; ablation: 54 needed, 0 unused; linter: no warnings in 2 files.

- *`paperg_brocard`, with the extensions* (68 theorems): mutations 152/152 rejected, none by timeout; no contradiction in 28 probed declarations (40 without hypotheses); satisfiability

certified for 7 of 68, 61 unresolved; ablation: 111 needed, 0 unused; linter: no warnings in 4 files. The run on the other machine gave the same figures.

The single unused binder in `paperg_nagel` is the hypothesis $0 < y + z$ of the auxiliary lemma `sub_pos_aux`, a redundant positivity condition. These figures have limits. The mutation generator alters only conclusions of certain forms, so a declaration without mutants is not mutation-tested; most declarations have no hypotheses beyond the triangle and so are not probed for contradictions; and most satisfiability cases are unresolved by the automatic certifier. For the main theorems, satisfiability and non-degeneracy are supplied instead by the explicit witnesses of the next paragraph. Ablation shows which binders the existing proof uses, not which are logically necessary. With the Layer-14 and Layer-15 files the battery was run again on both machines. For `paperg_nagel` at the commit before `Inradius.lean` was added (180 theorems), both runs rejected 440 of 448 mutations (11 by the heartbeat limit), with 8 surviving and no contradiction; satisfiability was certified for 30 of 180 on one machine and 32 of 180 on the other (the certifier's search is time-limited). With `Inradius.lean` (182 theorems) 446 of 454 mutations were rejected, with the same 8 survivors and no contradiction. For `paperg_brocard` with the Layer-15 file (147 theorems), two runs on the x86-64 machine and one on the arm64 machine each rejected 285 of 287 mutations, with 2 surviving and no contradiction (one probe was reported invalid by the harness); satisfiability was certified for 26 to 28 of 147. Besides `sub_pos_aux`, the ablation step reports eight unused binders, all in Layer-15 files. In five classical statements (N3, N4, N6a, N6b, S1b) the hypothesis that the point is on the third cevian or symmedian is not needed, because the other two already fix the point; the two nonzero-direction hypotheses of the classical Droz-Farny statement are not used by its proof; and in the auxiliary lemma `symm_mem` the hypothesis $0 < S$ is redundant. The classical hypotheses are the classical writer's, kept verbatim. Nine of the ten survivors change a coordinate of a Layer-14 witness triangle (for example $(4, 0) \mapsto (5, 0)$) inside a non-collinearity lemma; the mutated triangle is still non-degenerate, so the mutated lemma is true and its survival is expected. The tenth changes a comment in the Layer-15 Droz-Farny file and has no effect on any statement. Finally the battery was run on fresh clones of the published commits on both machines (reports `*_pub_*` in the supporting material). `paperg_nagel` (182 theorems): 446 of 454 mutations rejected (11 by the heartbeat limit), 8 surviving, no contradiction, satisfiability certified for 30 and 31 of 182; `paperg_brocard` (147 theorems): 285 of 287 rejected, 2 surviving, no contradiction (one probe invalid), satisfiability certified for 26 and 28 of 147. On both machines the survivors are the same ten classified above, and the ablation stage reports the same unused binders.

Statements in Mathlib's language and non-vacuity (Layer 14). A Layer-14 file in each development (`NagelLine/L14.lean`, `Brocard/L14.lean`) restates the principal assertions of each main theorem with Mathlib-only types, the library's points being unfolded into Mathlib terms (explicit affine combinations with weights written in `dist`, `t.medial.incenter` for the Spieker centre, $H + 2(O - H)$ for the de Longchamps point, and "any `Affine.Triangle` whose vertices are the images of A, B, C under the homothety with centre G and ratio -2" for the anticomplementary triangle), and proves it from the library theorem. A check with fully qualified pretty-printing found no constant of either library in the elaborated types of these 16 + 9 restatements or of the 5 + 5 witness theorems. Each witness theorem proves the hypotheses and the non-degeneracy of the conclusion on an explicit triangle in `EuclideanSpace` $\mathbb{R}$ `(Fin 2)`:

- For the Nagel family, mainly the 3-4-5 triangle $A = (0, 0)$, $B = (4, 0)$, $C = (0, 3)$: $I = (1, 1)$, $G = (4/3, 1)$, $N = (2, 1)$, Gergonne point $(8/11, 9/11)$, Spieker centre $(3/2, 1)$; the cevians are pairwise distinct lines, I, G, N are pairwise distinct, the incircle and excircle contact points differ on every side, the anticomplementary triangle has incentre $(2, 1) \neq I$, and the excentres

$(6,6)$, $(-2,2)$, $(3,-3)$ with exradii $6,2,3$ give power $37/4$ for the Spieker centre. Since $H = A$ in this right triangle, the de Longchamps witness uses $(0,0)$, $(4,0)$, $(1,3)$ instead: $O = (2,1)$, $H = (1,1)$, $L = (3,1)$, pairwise distinct and distinct from G.

- For the symmedian and Brocard family, $A = (0,0)$, $B = (4,0)$, $C = (1,3)$ ($a^2 = 18$, $b^2 = 10$, $c^2 = 16$, $\Delta = 6$): $K = (14/11, 12/11)$, $\Omega = (220/157, 120/157)$, $\Omega' = (232/157, 216/157)$, $\cot\omega = 11/6$ and $0 < \omega < \pi/6$; K differs from the vertices and from G, and O, K, Ω, Ω' are pairwise distinct, so Theorem 15 is non-vacuous; the centroid G is not the centroid of its own pedal triangle; the six Lemoine points are pairwise distinct; and for Droz-Farny, with $d_1 = (1,2)$, $d_2 = (2,-1)$, all hypotheses hold and the three midpoints $(10/3, 2/3)$, $(-2/7, -6/7)$, $(7/4, 0)$ are pairwise distinct.

Two clauses are not restated there: the strict-betweenness clause of Theorem 2 and the two forms of the common value in Theorem 9, which are stated only in the library's own terms. No restatement turned out false or needed a hypothesis that the library theorem lacks. With the Layer-14 files the gates pass with 91 (`paperg_nagel`) and 88 (`paperg_brocard`) declarations on the x86-64 machine, and `paperg_nagel` also on the arm64 machine; the tampering suites reject every planted defect.

Blind restatement from this paper (Layer 13). A version of this paper with every Lean name, listing, table of declarations and source link removed was given to a fresh AI writer that had no access to the code; from the prose alone it wrote 36 Lean statements for the 18 numbered theorems, with its own definitions, compiled with only `sorry` warnings. Its session log shows that it read only that text and its own files. A comparer then proved, for each pair, the blind statement from the library statement and conversely, using only unfolding, agreement lemmas between the two sets of definitions and general Mathlib lemmas, never the library's own theorems. 35 of the 36 pairs are equivalent in both directions. The exception was Theorem 9: the blind statement included the inradius form $(s^2 + r^2)/4$ written in this paper, which the library did not state, so the blind statement was strictly stronger. We added the two theorems `NagelLine.inradius_sq` and `NagelLine.spiekerCenter_power_exsphere_eq_inradius`; with them the blind statement follows by repackaging. All comparison files compile with no `sorry` and only the standard axioms, and the library hash was unchanged by the comparison.

Comparison with the classical statements (Layer 15). A second fresh writer, given only textbook statements of fifteen classical results (N1–N8: Gergonne, Nagel, Nagel line, Spieker, isotomic contacts, medial and anticomplementary triangles, de Longchamps, Spieker radical centre; S1–S7: Lemoine, pedal centroid, Brocard points and angle, isogonality, Brocard circle, first Lemoine circle, Droz-Farny) and neither this paper nor the code, wrote 23 Lean statements. Comparers then proved each classical statement from the libraries. All ten Nagel-family statements are covered. The classical statements are about *any* common point of the cevians, while the library defines the points by weights and shows they lie on the cevians; the comparer closed this gap with a new uniqueness proof (two cevians from different vertices meet in at most one point), and also proved the four-point collinearity of I, G, S_p, N, which the library states as two three-point collinearities. In the symmedian and Brocard family, the pedal-centroid statements and Droz-Farny are covered directly; the rest were narrower in the library (it proves existence for its explicit points) and were completed by new proofs: uniqueness of the common point of the symmedians, uniqueness of each Brocard point in the interior, and the first Lemoine circle for any six points with the defining incidences. Four classical statements (N4, N6b, N7, N8) carry a non-degeneracy hypothesis on the medial or anticomplementary triangle written by the classical writer; the comparer

proved these hypotheses redundant (the medial and anticomplementary triangles of a triangle are non-degenerate). No classical statement disagreed with the library. These proofs, including all new uniqueness results, were then moved into the libraries as `NagelLine/L15Classical.lean` and `Brocard/L15Classical.lean`, which work in the plane; with them the gates passed with 149 and 147 declarations, and adding `NagelLine/Inradius.lean` after the Layer-13 comparison raised `paperg_nagel` to 151.

Statements checked by hand. The author read each of the eighteen numbered theorems, as given in this paper, and the starred Layer-15 classical statements against their Lean statements at the published commits (67 Lean statements in all; a sheet pairing the two, with links to the source lines, is in the verification repository) and found them to agree. This is a reading check, not a proof.

9 What is not claimed

- We do not claim new mathematics: all theorems are classical.
- We do not claim Euler's $OI^2 = R(R - 2r)$, the Euler line, the nine-point circle or any other result already in Mathlib or in an open Mathlib pull request; we use Mathlib's Euler line in Theorem 5(iii).
- We do not claim the first formal treatment of the Nagel and Gergonne cevians in every sense: an algebraic form with formula-defined feet was formalised earlier in `lean-genius` (Section 7). Nor do we claim the first formal proof of the Nagel line or of the Spieker centre's position on it in barycentric form, which is among the Coq facts of [10]. Our claim is to the statements with Mathlib's contact points, incentre and centroid.
- In the numbered theorems the points are given by their weights, and the theorems prove that these points have the classical properties (Theorem 10 by angle equalities at each vertex); the only uniqueness statement among them is Theorem 11(ii). The additional Layer-15 files (Section 8), which work in the plane `EuclideanSpace ℝ (Fin 2)`, prove uniqueness of the Nagel point, of the common point of the symmedians and of each interior Brocard point, and express isogonality by reflected-line incidence. We define no isogonal-conjugation map.
- We do not claim novelty beyond the sources searched; see the gaps in Section 7.

10 Limitations

1. **Definitions by weights.** The Nagel, Gergonne, symmedian and Brocard points and the Lemoine points are defined by barycentric formulas. The theorems connect them to Mathlib's contact points, incentre, centroid, orthocentre, circumcentre and angles, but a reader must check that the weights are the intended ones; the Layer-14 restatements unfold them so that this check needs only Mathlib.
2. **One labelling.** Statements are for the given order A, B, C; where a statement has three cyclic forms, all three are stated (contact points, cevians, symmedian angles), but no general relabelling lemma is proved.

3. **Dimension.** The pedal converse and Droz-Farny assume a two-dimensional space. The additional Layer-15 developments are also planar.

4. **The equilateral case.** The right-angle form of the Brocard circle is vacuous when the points coincide, which happens exactly for equilateral triangles (Section 6).

5. **Battery coverage.** Many declarations have no mutants or no hypotheses, and most satisfiability cases are unresolved by the automatic certifier (Section 8).

6. **All-AI checking.** The proofs, the numerical checks, the battery and the restatements were all done by AI agents; the Layer-15 writer and the reviewer were of a different model family from the writers of the proofs. The only human check is the author's reading of the statements against the paper (Section 8); no human has checked the proofs line by line, which is the kernel's job.

7. **Prior art.** The search had the gaps listed in Section 7.

11 Availability

The proofs are in two Lean 4 developments on Lean v4.33.1 with Mathlib pinned at revision `0df444a3` (in `lake-manifest.json`). Line counts are of the library `.lean` files, including the root import file, at the published commits.

repository	library files	files	lines
`paperg_nagel`	`Basic` (285), `Main` (414), `Extensions` (362), `Inradius` (45), `L14` (408), `L15Classical` (741)	7	2,261
`paperg_brocard`	`Basic` (57), `Lemmas` (338), `Main` (463), `ExtLemmas` (384), `Extensions` (546), `L14` (522), `L15Classical` (723)	8	3,040

The repositories are `https://github.com/jpbald93/paperg_nagel` at commit `e40ee54f20588f5181d348767f3f31be20406cc1` and `https://github.com/jpbald93/paperg_brocard` at commit `ebb35755a81aec2c7c0673fe93baf8cc5d6eb22d`; each is published as a single commit (the development history is not included). Every `File:line` reference in this paper links to that commit. Supporting material (the prior-art sweep, the builders' reports, the battery and build reports, the Layer-14 files and reports, and the fresh-clone receipts) is in the folder `PaperG-triangle-centres` of the repository `https://github.com/jpbald93/LEAN-verification`. Every name in `\decl` font in this paper is in the `REQUIRED` list of the repository that contains it at the published commit; a script checked all 102 distinct names (108 occurrences) against the published `REQUIRED` lists, and all 31 `File:line` references (and the further line numbers listed with them) against the declarations at those lines.

Acknowledgements and use of AI

Large-language-model AI assistants were used to write the Lean proofs, to run the prior-art search, the numerical checks and the verification battery, and to draft this paper; parts of the checking were done by a different model from the one that wrote the proofs. The author directed the work and reviewed the paper. Every proof is checked by the Lean kernel, and the author takes responsibility for all content.

References

[1] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE-28, Lecture Notes in Computer Science **12699**, Springer, 2021, 625–635.

[2] The mathlib Community, *The Lean mathematical library*, in: Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP '20), ACM, 2020, 367–381.

[3] W. Wang, *feat(Geometry/Euclidean): prove Euler's theorem in geometry*, mathlib4 pull request #44230, opened 26 September 2026 (open at the time of writing). `https://github.com/leanprover-community/mathlib4/pull/44230`

[4] C. Kimberling, *Encyclopedia of Triangle Centers*, `https://faculty.evansville.edu/ck6/encyclopedia/ETC.html`.

[5] R. A. Johnson, *Advanced Euclidean Geometry*, Dover, New York, 1960 (unabridged republication of *Modern Geometry*, Houghton Mifflin, 1929).

[6] H. S. M. Coxeter and S. L. Greitzer, *Geometry Revisited*, Mathematical Association of America, 1967.

[7] GitHub user `rjwalters`, *lean-genius* (GitHub repository `https://github.com/rjwalters/lean-genius`), file `proofs/Proofs/CevasTheoremOQ04OQ04OQ02.lean`, first committed 25 June 2026; consulted at commit `470a7df`, 22 October 2026, `https://github.com/rjwalters/lean-genius/blob/470a7df/proofs/Proofs/CevasTheoremOQ04OQ04OQ02.lean`.

[8] P. Yiu, *Introduction to the Geometry of the Triangle*, lecture notes, Florida Atlantic University, Summer 2001; version 12.1224, December 2012.

[9] J.-L. Ayme, A purely synthetic proof of the Droz-Farny line theorem, *Forum Geometricorum* **4** (2004), 219–224.

[10] J. Narboux and D. Braun, Towards a certified version of the Encyclopedia of Triangle Centers, *Mathematics in Computer Science* **10** (2016), 57–73, `https://doi.org/10.1007/s11786-016-0254-4`; website `http://dpt-info.u-strasbg.fr/~narboux/CETC/` (consulted through the Internet Archive, snapshots of March 2019).

[11] S.-C. Chou, *Mechanical Geometry Theorem Proving*, Mathematics and Its Applications **41**, D. Reidel (Springer Netherlands), 1987/1988, `https://doi.org/10.1007/978-94-009-4037-6`.

[12] S.-C. Chou, X.-S. Gao and J.-Z. Zhang, *Machine Proofs in Geometry: Automated Production of Readable Proofs for Geometry Theorems*, World Scientific, 1994, `https://doi.org/10.1142/2196`.

[13] R. Honsberger, *Episodes in Nineteenth and Twentieth Century Euclidean Geometry*, New Mathematical Library **37**, Mathematical Association of America, 1995.

[14] J. Bald, *Classical plane geometry by coordinate certificates: Desargues, Monge, Simson, Napoleon and Lami in Lean 4*, Zenodo, 2026. `https://doi.org/10.5281/zenodo.23239930`

[15] J. Bald, *The Malfatti circles in Lean 4, with independent certificate proofs of Feuerbach, Miquel and Brahmagupta*, Zenodo, 2026. https://doi.org/10.5281/zenodo.23240500

[16] J. Bald, *Barbier's theorem for the Reuleaux triangle and the convex four-vertex theorem in Lean 4, through support functions*, Zenodo, 2026. https://doi.org/10.5281/zenodo.23323062

[17] J. Bald, *Circle configurations in classical plane geometry, formalized in Lean 4: Clifford's chain, Miquel's five circles, Lester, Sawayama–Thébault and Kosnita*, Zenodo, 2026. https://doi.org/10.5281/zenodo.23378578

[18] J. Bald, *The complete quadrilateral in Lean 4: Newton–Gauss, Bodenmiller, the Miquel point, the Steiner circle and the Simson line*, Zenodo, 2026. https://doi.org/10.5281/zenodo.23407551

[19] J. Bald, *The inscribed parabola and the Poncelet point: the complete quadrilateral and the complete quadrangle in Lean 4*, Zenodo, 2026. https://doi.org/10.5281/zenodo.23653615