

Error & One-Sample Tests

PSY 3307

Jonathan A. Pedroza, PhD

Cal Poly Pomona

2022-02-22

Inflated Error Rates

- **familywise error rate** or experimentwise error rate is the change in probability of making a type I error

$$\text{familywise error} = 1 - .95^n$$

$$0.40 = 1 - 0.95^{10}$$

- 40% chance of making a type I error (false positive)

Inflated Error Rates

- **Bonferroni Correction** is a correction to make the alpha accurate for the number of tests you are making
 - alpha is your pre-determined alpha/probability, k is the number of tests

$$P_{crit} = \frac{\alpha}{k}$$

$$P_{crit} = \frac{.05}{4}$$

$$P_{crit} = .0125$$

Statistical Power

- opposite issue from Type II error is known as statistical **power**
 - chance of not finding a true significant finding (false negative)
- false negative is beta
 - .2 of a probability is the norm for beta or power of .8
 - 80% chance that you detect an effect that is truly there

$$1 - \beta$$

Power

- we can think about increasing power with the following information
 - alpha, sample size, size of the effect desired
- calculate the power of the test (beta)
 - we have alpha and sample size
- calculate sample size to achieve a given level of power
 - we have alpha and beta


Power

- Power can be determined a priori (before analyses) or post hoc (after analyses)
- Ways to improve power
 - increase sample size
 - potentially use a one-tailed test over a two-tailed test
 - use parametric tests over nonparametric tests

Confidence Intervals & Statistical Significance

- if your confidence intervals barely touch on the ends then you'll have a significance level approximately at .01
- if there is a gap between your confidence intervals then you have a significance level less than .01
- slight overlap between your confidence intervals and you'll have a significant difference less than .05
- if confidence intervals overlap a lot then you have no difference between groups

Example



Sample Size & Statistical Significance

- even with means and sd at the same level for groups/conditions
 - with a large enough sample size, you'll see a difference
- large samples, small differences can be significant (evidence of significant effect)
 - small samples, large differences can be non-significant (no evidence of significant effect)

Reporting Significance Tests

- 95% confidence intervals are in brackets []
 - The age of participants ($M = 23.5$, $SD = 1.34$, 95% CI [21.1, 24.6])
- The age of students in PSY 3307-01 and PSY 3307-03 are significant different, $p < .05$ (using tables)
 - The age of students in PSY 3307-01 and PSY 3307-03 are significant different, $p = .034$

Effect Sizes

- there are many problems with NHST, but many of the findings from NHST don't tell us how strong of an effect there is
 - most statistics tell us we passed the critical value/threshold
- a standardized measure of an effect/strength of a relationship is known as an **effect size**
 - standardized = compare to other effects from other analyses or other studies
- many types of effect sizes, but let's focus on the most useful for this class
 - Cohen's d

Cohen's d/Hedges' g

- when samples are small ($N < 20$) --> Hedge's g
 - when greater than 20, both are roughly the same

$$g = \frac{\bar{X}_1 - \bar{X}_2}{SD_{pooled}} * \frac{N - 3}{N - 2.25} * \sqrt{\frac{N - 2}{N}}$$

$$\hat{d} = \frac{\bar{X}_1 - \bar{X}_2}{S}$$

- helpful primarily for use in independent samples t-tests
 - where we are comparing two groups
- when groups have equal SD
 - use the SD

Cohen's d/Hedges' g

- when groups have unequal SD
 - use the SD for the control group/baseline OR
 - pool the SD of the two groups (if they are independent of one another)

$$S_p = \sqrt{\frac{(N_1 - 1)S_1^2 + (N_2 - 1)S_2^2}{N_1 + N_2 - 2}}$$

- we'll cover this a lot more when conducting statistics that are not standardized

Logic of Hypothesis Testing

- At least for the current statistical testing we will be conducting (z-test)
- We are interested whether our DV score in our sample is representative of the population (H_0) or if our sample's DV score is significantly different from the population DV score (H_1)
- This means we will be using:
 - probability
 - region of rejection
 - criterion
 - critical value

Example

- We are interested in knowing if CPP students eat more or less fast food than the average CSU student H_0 : H_1 :
- The average number of a times a week that a CSU student eats fast food is 3.7, sd is 1.2
- Sampling from CPP, we find out the average CPP student eats fast food 2.4 times a week
- We would then conduct a z-test to see if our sample is statistically significantly different from CSU students
- Before doing any analyses, we would create our criterion for what counts as a significant finding

The z-Test

1. Randomly selected a sample
2. DV is somewhat normally distributed in population
 - and is ratio or interval scale
3. Know population mean of raw scores under another condition of the IV
4. Know population standard deviation

Sampling Distribution of Two-tailed Test

1. Create sampling distribution of means from population raw scores
 - This will be what our H_0 states
2. Identify what the population mean is for H_0
3. Select an alpha
 - **alpha** greek letter for criterion probability (e.g., .05)
4. Identify regions of rejection
 - One-tailed or two-tailed test
5. Determine critical value
 - two-tailed is $z = 1.96$

Compute the z-score

$$z_{obt} = \frac{\bar{X} - \mu}{\sigma_{\bar{X}}}$$

- Need the standard error of the mean before getting the z-score
- z is now obtained from the data, that is why it is z obt in the formula above
- Choosing some random values
 - mu is 3.7
 - sigma is 1.2
 - N is 100
- We would calculate xbar from our sample, let's just call it 2.4

```
# standard error calculation  
1.2/sqrt(100)
```

```
## [1] 0.12
```

```
# se is .12  
  
(2.4 - 3.7)/.12
```

```
## [1] -10.83333
```

```
# z-score is -10.83
```

Comparing Obtained z-value to Critical Value

- We know that the critical value is ± 1.96
- Since our value is outside of the critical value in the region of rejection
 - we can reject that CPP students eat fast food as much as the rest of the CSU system students
- We have rejected the null hypothesis

Interpretation of STATISTICALLY Significant & Nonsignificant Results

- If you reject the null, you have statistically significant findings
 - H_1 is supported, there is a relationship/there are differences
- If you retain the null, you don't have statistically significant findings
 - H_0 supported, there is no relationship/there are no differences

What Does a Statistically Significant z-test Finding Represent

- Significant MEANS NOTHING
- Statistically significant indicates you reject the null and your sample is different from the population (for z-tests)
- We can't prove that H_0 is false
- We don't know the exact population mean represented by our sample
 - potential **sampling error**, your sample may not be representative of the population

What Does a Nonsignificant z-test Finding Represent

- our sample is representative of the population
 - CPP students eat fast food as much as CSU students
- Don't say insignificant
 - best way of stating this is: "there was no evidence supporting that CPP students are statistically different from CSU students"
- We didn't find evidence of a statistically significant difference/relationship
- We simply have failed to reject the null hypothesis

z-Test Summary

- Create hypothesis/es
- Set up sampling distribution, select alpha, location region of rejection, determine critical value
- compute \bar{x} , standard error and z-score obtained from population mean and standard deviation
- compare obtained z-score to critical value
 - statistically significant = reject H_0
 - nonsignificant = retain H_0
 - don't draw conclusions

The One-Tailed Test

- predict scores in a direction
- interested in whether DV scores are higher or lower, not both
- the alpha still needs to be determined prior to analyses

Example

- H_0 :
- H_1 : CSUDH students will eat more fast food than CSU students
- CSUDH students eat 4.1 times a week

```
# standard error calculation  
1.2/sqrt(100)
```

```
## [1] 0.12
```

```
# se is .12  
  
(4.1 - 3.7)/.12
```

```
## [1] 3.333333
```

```
# z-score is 3.33
```