

# Hydra: a mixture modeling toolkit for subtyping pediatric cancer cohorts using multimodal gene expression signatures

Jacob Pfeil<sup>1,2\*</sup>, Lauren M. Sanders<sup>1,2,3</sup>, Ioannis Anastopoulos<sup>1,2</sup>, A. Geoffrey Lyle<sup>2,3</sup>, Alana S. Weinstein<sup>1,2</sup>, Yuanqing Xue<sup>1,2</sup>, Andrew Blair<sup>1,2</sup>, Holly C. Beale<sup>2,3</sup>, Alex Lee<sup>4</sup>, Stanley G. Leung<sup>4</sup>, Phuong T. Dinh<sup>4</sup>, Avanthi Tayi Shah<sup>4</sup>, Marcus R. Breese<sup>4</sup>, W. Patrick Devine<sup>5</sup>, Isabel Bjork<sup>2</sup>, Sofie R. Salama<sup>1,2,6</sup>, E. Alejandro Sweet-Cordero<sup>4</sup>, David Haussler<sup>1,2,6</sup>, Olena Morozova Vaske<sup>2,3</sup>

**1** Department of Biomolecular Engineering, University of California, Santa Cruz, California, United States of America

**2** Genomics Institute, University of California, Santa Cruz, California, United States of America

**3** Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, California, United States of America

**4** Department of Pediatrics, Division of Hematology and Oncology, University of California, San Francisco, California, United States of America

**5** Department of Anatomic Pathology, University of California, San Francisco, California, United States of America

**6** Howard Hughes Medical Institute, University of California, Santa Cruz, California, United States of America

\* Corresponding author: [jpfeil@ucsc.edu](mailto:jpfeil@ucsc.edu)

## Abstract

Precision oncology has primarily relied on coding mutation status as a readout to define potential therapeutic benefit. Incorporation of transcriptome analysis into precision oncology workflows has proven to be challenging, as relative rather than absolute gene expression level needs to be considered, requiring differential expression analysis across samples. However, cell-of-origin and tumor microenvironment effects limit the effectiveness of these approaches. To address these challenges, we developed an unsupervised clustering approach for discovering differential pathway expression within cancer cohorts using gene expression measurements. Hydra is an unsupervised gene clustering approach that models differential expression as a multi-modal distribution. Multivariate clustering of multimodally expressed genes reveals differential pathway expression and tumor subtype signatures. We demonstrate that the hydra approach is more sensitive than widely-used gene set enrichment approaches for detecting multimodal expression signatures. We applied the hydra pipeline to high-risk neuroblastoma and osteosarcoma samples and discovered expression signatures associated with changes in the tumor microenvironment. These expression signatures were consistent with pathology review of the H&E slide images. Furthermore, we identified an association between ATRX deletions and elevated immune marker expression in high-risk neuroblastoma samples. Hydra is available as a Docker container for easy deployment (<https://hub.docker.com/r/jpfeil/hydra>). The source code is available on GitHub ([www.github.com/jpfeil/hydra](http://www.github.com/jpfeil/hydra)).

## Author summary

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Curabitur eget porta erat. Morbi consectetur est vel gravida pretium. Suspendisse ut dui eu ante cursus gravida non sed sem. Nullam sapien tellus, commodo id velit id, eleifend volutpat quam. Phasellus mauris velit, dapibus finibus elementum vel, pulvinar non tellus. Nunc pellentesque pretium diam, quis maximus dolor faucibus id. Nunc convallis sodales ante, ut ullamcorper est egestas vitae. Nam sit amet enim ultrices, ultrices elit pulvinar, volutpat risus.

## Introduction

Lorem ipsum dolor sit [1] amet, consectetur adipiscing elit. Curabitur eget porta erat. Morbi consectetur est vel gravida pretium. Suspendisse ut dui eu ante cursus gravida non sed sem. Nullam Eq (1) sapien tellus, commodo id velit id, eleifend volutpat quam. Phasellus mauris velit, dapibus finibus elementum vel, pulvinar non tellus. Nunc pellentesque pretium diam, quis maximus dolor faucibus id. [2] Nunc convallis sodales ante, ut ullamcorper est egestas vitae. Nam sit amet enim ultrices, ultrices elit pulvinar, volutpat risus.

$$P_Y = \underbrace{H(Y_n) - H(Y_n|\mathbf{V}_n^Y)}_{S_Y} + \underbrace{H(Y_n|\mathbf{V}_n^Y) - H(Y_n|\mathbf{V}_n^{X,Y})}_{T_{X \rightarrow Y}}, \quad (1)$$

Large cancer sequencing projects, including The Cancer Genome Atlas (TCGA) and Therapeutically Applicable Research to Generate Effective Treatments (TARGET), have facilitated the development of cancer gene expression compendia (Vivian et al. 2017; Pugh et al. 2013; Goldman et al. 2018; Weinstein et al. 2013; Newton et al. 2017) but these compendia often lack corresponding normal tissue expression data. Without the the normal comparator, Hoadley et al. (2018) found that cell-of-origin signals drive integrative clustering of TCGA data, with the exception of tumors that have high immune and stromal expression clustering irrespective of the cell-of-origin. Strong cell-of-origin and tumor microenvironment (TME) signals may complicate the interpretation of gene expression results for precision oncology applications, so careful modeling of the data is necessary to subtract out unwanted signals. The TME includes tumor cells, stromal fibroblasts, infiltrating immune cells, and vasculature (Joyce and Fearon 2015). Similarities in TME composition across tumor samples have led to the identification of TME states (i.e. inflamed, immune-excluded, immune-desert). These states are dynamic and may change as the tumor progresses, but may shed light on the immunogenicity of tumor cells and correlate with response to cancer immunotherapies (Chen and Mellman 2017). The TME cellular composition can be inferred from gene expression data since host cell RNA is sequenced along with the cancer cell RNA. Not accounting for the immune signal when performing gene expression analysis may confound the interpretation of clustering results (Aran et al. 2017; Rhee et al. 2018). Tumor progression and response to therapies is associated with features of the TME, so targeting the TME therapeutically may improve the treatment of some cancers. Immunotherapies that activate the host immune system to eradicate tumors have improved the treatment of several cancer types (Mellman et al. 2011; Page et al. 2014). Pediatric cancers are thought to be less immunogenic because pediatric cancers have lower mutation burdens in general. Gene overexpression is another source of tumor-specific antigens which may play an important role in pediatric tumor immunogenicity. In addition to infiltrating immune cells, cancer-associated fibroblasts

assist in extracellular matrix remodeling and activation of growth factor signaling that facilitate tumor growth, metastasis, and resistance to therapies. Methods are needed to both infer and characterize gene expression subtypes that correlate with tumor microenvironment states to accelerate the development of novel therapies for pediatric cancers. Tumor/normal differential expression analysis in which a cohort of tumor tissues is compared to normal counterparts is an effective approach for identifying gene expression biomarkers (Anders and Huber 2010; Soneson and Delorenzi 2013; Anders et al. 2013), but it is often not possible to conduct this analysis in a clinical setting. Sufficient biological and technical replicates are limited by tumor tissue availability, and healthy neighboring tissue often cannot be isolated. In addition, for many pediatric cancers, the cell-of-origin and thus the appropriate reference normal tissue is not known. Besides differential expression analysis, single-sample pathway analysis can be used to identify upregulation of biological gene sets in tumor subtypes. The most widely used pathway analysis approach is gene set enrichment analysis (GSEA). GSEA identifies coordinated expression of pathway genes using gene ranks and a Kolmogorov-Smirnov-like test statistic (Subramanian et al. 2005; Mootha et al. 2003). GSEA is usually performed on differentially expressed genes, but single-sample versions exist for identifying cancer subtypes. GSEA uses curated pathway gene sets like those in the Molecular Signatures Database (MSigDB) (Liberzon et al. 2011). Cancer gene expression subtypes are traditionally identified using unsupervised clustering analysis such as consensus clustering analysis (Oyelade et al. 2016; John et al. 2018; Wilkerson and Hayes 2010). These methods are generally underpowered because the number of genes greatly exceeds the number of samples. Dimensionality reduction approaches such as Principal Component Analysis (PCA) have been found to underestimate the dimensionality of gene expression data. Lenz et al. (2016) found two cases in which PCA fails to identify a biological signal: when the size of the cluster is small and when the effect size is small. Lenz et al. (2016) suggest investigating multimodally expressed genes to improve identification of subtypes. Cancer subtypes naturally lead to multimodal expression patterns where each subtype expresses genes with distinct levels and correlation patterns. Furthermore, enriching for multimodally expressed genes before clustering has been shown to improve correlations with clinical features of interest (Yi Li et al. 2005). Gaussian mixture models are a powerful class of unsupervised clustering algorithms for detecting multimodally expressed genes (Ghosh 2004). A Gaussian mixture model is appropriate when the expression data can be modeled by two or more Gaussian distributions (Gelman et al. 2014). One limitation of Gaussian mixture models in this context is that the number of clusters in the data is often not known beforehand, so a parameter search is used to identify the best-performing model. However, this is a computationally expensive approach. This problem can be overcome by placing a Dirichlet process prior on the number of expression clusters. The number of clusters is then inferred while fitting the mixture model using Markov chain Monte Carlo sampling (Gelman et al. 2014). This approach has not been widely used in clinical cancer research because these algorithms are also computationally intensive, but recent advances in approximate sampling methods make this approach scalable for precision oncology applications. Here, we describe a Dirichlet process mixture model pipeline called hydra for identifying clinically relevant expression subtypes and classifying N-of-1 tumor samples. The hydra pipeline uses the nonparametric Bayesian Python library `bnpy` (Hughes and Sudderth 2015) and incorporates state-of-the-art cluster profiling software and biological pathway gene sets. We provide an overview of the hydra pipeline, assess performance for detecting differential pathway expression, and apply the approach to better understand expression patterns in high-risk neuroblastoma and osteosarcoma. We propose a novel framework for N-of-1 tumor gene expression analysis and show that this framework can identify distinct immune and stromal

Materials and methods

90

Etiam eget sapien nibh

91

Nulla mi mi, Fig 1 venenatis sed ipsum varius, volutpat euismod diam. Proin rutrum  
vel massa non gravida. Quisque tempor sem et dignissim rutrum. Lorem ipsum dolor sit  
amet, consectetur adipiscing elit. Morbi at justo vitae nulla elementum commodo eu id  
massa. In vitae diam ac augue semper tincidunt eu ut eros. Fusce fringilla erat porttitor  
lectus cursus, S1 Video vel sagittis arcu lobortis. Aliquam in enim semper, aliquam  
massa id, cursus neque. Praesent faucibus semper libero.

92  
93  
94  
95  
96  
97

**Fig 1. Bold the figure title.** Figure caption text here, please use this space for the  
figure panel descriptions instead of using subfigure commands. A: Lorem ipsum dolor  
sit amet. B: Consectetur adipiscing elit.

Synthetic Data Generation and Validation We tested hydra’s ability to detect  
differential pathway expression using synthetic cancer data. We compared hydra to two  
widely used gene set enrichment tools: single-sample gene set enrichment analysis  
(ssGSEA) and gene set variation analysis (GSVA) (Barbie et al. 2009; Hänzelmann et al.  
2013; Tarca et al. 2013). Both methods are implemented in the GSVA R package  
(Hänzelmann et al. 2013). We modeled cancer gene expression as a multivariate  
Gaussian distribution, and then estimated the mean vector and covariance matrix using  
the TARGET MYCN wild-type neuroblastoma cohort (n=70). This approach allowed  
us to model pediatric cancer gene expression data while also controlling for  
subtype-related expression variation. We downloaded the RSEM-quantified TPM  
normalized gene expression measurements from the UCSC Xena Browser (Goldman et  
al. 2018). To reduce heteroscedasticity and the effect of outlier expression levels, we  
then transformed the expression data to  $\log_2(\text{TPM} + 1)$  (Zwiener et al. 2014). We  
defined an expression subtype as a subset of samples with a distinct expression  
signature. We modelled expression subtypes across the top 20 most highly expressed  
MSigDB Hallmark gene sets with at least 100 genes. We tested a range of synthetic  
data parameters related to the number of differentially expressed genes within a gene  
set and the effect size for these genes. We randomly generated the covariance matrix for  
the cancer subtype expression data. We tested the effect of having 10

98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116

TARGET Gene Expression Analysis The TARGET neuroblastoma and  
osteosarcoma RNA-Seq gene expression data were downloaded from the UCSC Xena  
Browser (Goldman et al. 2018). The TARGET clinical data were downloaded from the  
TARGET Data Matrix (<https://ocg.cancer.gov/programs/target/data-matrix>). We  
trained the hydra pipeline on 70 International Neuroblastoma Staging System (INSS)  
stage 4, MYCN non-amplified neuroblastoma and 74 osteosarcoma expression profiles.  
The resulting models were tested using an independent dataset of pediatric RNA-Seq  
data shared through the UCSC Treehouse compendium. The Treehouse compendium  
uses the same gene expression pipeline as the TARGET samples from the UCSC Xena  
Browser (Vivian et al. 2017). The neuroblastoma unsupervised enrichment analysis  
included all genes with a mean expression greater than  $1.0 \log_2(\text{TPM} + 1)$ , a minor  
expression component probability of at least 20

117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128

Statistical Analysis A Kruskal-Wallis test was used to identify statistically significant  
differences across two or more groups, and a Mann-Whitney U test was used for  
pairwise tests using a Holm-Sidak correction for multiple hypothesis testing (Pedregosa  
et al. 2011; Jones et al. 2001). We used the scipy stats implementation of the

129  
130  
131  
132

Kruskal-Wallis test and the scikit-learn post hoc processing implementation of pairwise Mann-Whitney U tests. Spearman rank and Pearson correlation values were calculated using the scipy library (Jones et al. 2001). Correlations between clinical features and clusters were identified using the Fisher Exact Test implemented in the R stats package (R Core Team 2013). Survival analysis was done using the survminer package (Kassambara et al. 2017).

H&E Slide Preparation and Pathologist Review Pediatric tumor samples were flash frozen, embedded in OCT, and 5um cryosections were collected. Slides were hematoxylin and eosin (H&E) stained and imaged on a Leica DMI8, equipped with a HC PL APO 40x/0.85 NA objective and DFC7000T camera. H&E slides were reviewed by a licensed pathologist for evidence of inflammation and graded as having either minimal (≤ 10

## Results

Nulla mi mi, venenatis sed ipsum varius, Table 1 volutpat euismod diam. Proin rutrum vel massa non gravida. Quisque tempor sem et dignissim rutrum. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Morbi at justo vitae nulla elementum commodo eu id massa. In vitae diam ac augue semper tincidunt eu ut eros. Fusce fringilla erat porttitor lectus cursus, vel sagittis arcu lobortis. Aliquam in enim semper, aliquam massa id, cursus neque. Praesent faucibus semper libero.

Table 1. Table caption Nulla mi mi, venenatis sed ipsum varius, volutpat euismod diam.

Heading1				Heading2			
cell1row1	cell2 row 1	cell3 row 1	cell4 row 1	cell5 row 1	cell6 row 1	cell7 row 1	cell8 row 1
cell1row2	cell2 row 2	cell3 row 2	cell4 row 2	cell5 row 2	cell6 row 2	cell7 row 2	cell8 row 2
cell1row3	cell2 row 3	cell3 row 3	cell4 row 3	cell5 row 3	cell6 row 3	cell7 row 3	cell8 row 3

Table notes Phasellus venenatis, tortor nec vestibulum mattis, massa tortor interdum felis, nec pellentesque metus tortor nec nisl. Ut ornare mauris tellus, vel dapibus arcu suscipit sed.

## LOREM and IPSUM nunc blandit a tortor

### 3rd level heading

Maecenas convallis mauris sit amet sem ultrices gravida. Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec euismod ligula. Curabitur fringilla pulvinar lectus consectetur pellentesque. Quisque augue sem, tincidunt sit amet feugiat eget, ullamcorper sed velit. Sed non aliquet felis. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Mauris commodo justo ac dui pretium imperdiet. Sed suscipit iaculis mi at feugiat.

1. react
2. diffuse free particles
3. increment time by dt and go to 1

## Sed ac quam id nisi malesuada congue

Nulla mi mi, venenatis sed ipsum varius, volutpat euismod diam. Proin rutrum vel massa non gravida. Quisque tempor sem et dignissim rutrum. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Morbi at justo vitae nulla elementum commodo eu id massa. In vitae diam ac augue semper tincidunt eu ut eros. Fusce fringilla erat porttitor

lectus cursus, vel sagittis arcu lobortis. Aliquam in enim semper, aliquam massa id, cursus neque. Praesent faucibus semper libero.

- First bulleted item.
- Second bulleted item.
- Third bulleted item.

Hydra Method We developed a Bayesian non-parametric clustering pipeline for identifying biological and technical variation in large cancer gene expression datasets without the need for a reference normal dataset (Figure 1). Bayesian non-parametric models have been proposed for analyzing gene expression, but to our knowledge, this is the first reproducible and widely deployable implementation of a non-parametric mixture model pipeline designed for precision oncology gene expression analysis. The hydra pipeline is an open source software tool hosted on GitHub ([www.github.com/jpfeil/hydra](http://www.github.com/jpfeil/hydra)). The pipeline is written in Python 2.7 (Millman and Aivazis 2011) and R (R Core Team 2013) programming languages. A Docker container is available for deployment across environments ([www.dockerhub.com/jpfeil/hydra](http://www.dockerhub.com/jpfeil/hydra)). The hydra pipeline uses the bnpy Python library to fit Dirichlet process Gaussian mixture models (DP-GMM) to cancer gene expression data. While most clustering algorithms require setting the number of clusters before fitting the data, the DP-GMM learns the number of clusters while clustering the samples. The Dirichlet process is a distribution for modeling distributions. This feature makes the Dirichlet process an effective prior for the number of clusters in a mixture model. The DP-GMM learns the cluster mean vectors and covariance matrices and assigns each sample to a cluster (Hughes and Sudderth 2013, 2015). This process automates the identification of multimodally expressed genes and multivariate expression signatures. Coordinated expression of multimodally expressed genes, which is encoded in the correlation structure across genes, defines subtype expression signatures. The DP-GMM is also effective when clusters overlap, which is a beneficial property in gene expression data analysis. The parameters learned from fitting cancer gene expression data can be used to classify new samples for precision medicine applications. The hydra pipeline starts with gene expression preprocessing by mean centering and subsetting to genes of interest. The pipeline reads gene set annotations and standardizes gene symbols using the NCBI gene alias annotations (Benson et al. 2013). Next, hydra performs multimodal feature selection across all genes of interest. Multimodal feature selection has been shown to improve clustering performance, and the resulting clustering correlates better with clinical features (Yi Li et al. 2005). The multimodal filter removes unimodal expression distributions, selecting for multimodally expressed genes that harbor statistically significant differences in expression in multiple groups of samples within the cohort. The multimodally expressed genes are then used in downstream multivariate clustering. The hydra pipeline can be run in two modes. Mode 1 uses multimodally expressed genes to identify differential pathway expression within user-specified gene sets. Mode 2 looks for the enrichment of biological gene sets before multivariate clustering using the clusterProfiler tools (Yu et al. 2012). We use the gene set analysis (mode 1) for identifying known gene expression signatures and the gene set enrichment analysis (mode 2) for discovering unknown sources of variation. The pipeline is equipped with commonly used gene sets, including the Molecular Signatures Database (MSigDB) (Liberzon et al. 2011), the Gene Ontology Terms (Ashburner et al. 2000; The Gene Ontology Consortium 2019), and the EnrichmentMap gene sets (Merico et al. 2010). The gene set database is configurable to the user's research goals and additional gene sets can be added at runtime. The pipeline includes routines for cluster profiling and N-of-1 tumor analysis. Cluster profiling analysis includes GSEA to identify

pathway expression that characterizes each cluster. GSEA uses all available genes since these methods require non-differentially expressed genes to assess the significance of an enrichment score. A t-statistic is calculated for each gene comparing gene expression values of samples within and outside of a cluster. Cluster profiling GSEA uses the ranked gene-level t-statistics to determine gene set enrichment. The N-of-1 tumor analysis routine classifies a new gene expression profile into one of the inferred clusters, calculates a gene-level z-score for that sample relative to the normalized expression distribution, and performs GSEA. This procedure can identify subtle gene expression signatures that may not be detectable using the entire expression cohort.

**Performance Assessment using Synthetic Gene Expression Data** To assess how well hydra detects differentially expressed pathways as compared to common GSEA approaches, we applied these methods to synthetically-generated cancer gene expression data. We generated synthetic cancer gene expression data based on the TARGET high-risk neuroblastoma cohort and the Hallmark MSigDB gene sets (see Synthetic Data Generation and Validation section). We tested a range of effect sizes and percent differentially expressed genes (%DEG) within the MSigDB gene sets. We generated receiver operator curves (ROC) and calculated the Area Under the receiver operator Curve (AUC) for each analysis. Overall, the hydra pipeline outperformed the single-sample GSEA approaches with a mean AUC of 0.97 (95% CI: 0.96 - 0.98). ssGSEA had a mean AUC of 0.73 (95% CI: 0.72 - 0.75) and GSVA had a mean AUC of 0.69 (95% CI: 0.67 - 0.70) (Figure 2A). We further investigated the performance of these methods by plotting the AUC against the effect size at %DEG of 10 and 25% (Figure 2B). The hydra method performed better across all effect sizes, achieving near perfect performance above an effect size of 1.0 and 0.75 at %DEG of 10% and 25%, respectively. ssGSEA and GSVA performed similarly at low effect sizes, but ssGSEA performed better at %DEG of 10% and an effect size greater than 1.0. The performance difference between ssGSEA and GSVA was less pronounced at a higher %DEG of 25%. The ssGSEA and GSVA methods began to perform similarly to hydra at an effect size of 3.0 and %DEG of 25%. We next examined how the mean expression for a gene set, as a measure of the baseline expression, correlated with performance. We correlated the mean expression of the gene set with AUC and found that both the ssGSEA and GSVA AUC scores were negatively correlated with the mean expression (Pearson correlation: -0.21 and -0.18). The hydra method was positively correlated (Pearson correlation: 0.16) with the mean expression at low effect sizes (Figure 2C).

**Hydra Analysis of High-Risk Neuroblastoma Identifies Distinct Tumor Microenvironment States** After completing the performance assessment with synthetic gene expression data, we applied the hydra unsupervised enrichment analysis to the TARGET high-risk neuroblastoma cohort. High-risk neuroblastoma is an aggressive disease and is resistant to intensive therapy. Further subtyping of high-risk neuroblastoma may identify novel therapeutic targets and improve risk stratification. We hypothesized that unsupervised clustering of Gene Ontology terms would identify expression subtypes of high-risk neuroblastoma tumors. TumorMap analysis showed that the MYCN-non-amplified (MYCN-NA) neuroblastoma samples clustered separately from MYCN-amplified (MYCN-A) and stage 4S neuroblastomas (Supplementary Figure 1) samples. We focused on the MYCN-NA neuroblastoma tumor samples because this is the largest set of samples (N=70), and variation within MYCN-NA tumors is not well understood (Morgenstern et al. 2019). We applied the hydra unsupervised enrichment analysis to the MYCN-NA cohort. The multimodal expression filter identified 428 genes with a minor component probability greater than 20% (Supplementary Table 2). Gene Ontology analysis found enrichment for the following GO terms (FDR  $q \leq 0.01$ ): adaptive immune response (24 genes), mesenchyme development (12 genes), steroid hormone secretion (4 genes), and response to

corticosterone (4 genes). Multivariate Dirichlet process mixture model analysis of the 44 enriched GO term genes identified three clusters of neuroblastoma samples (Figure 3A). The posterior probability for belonging to each cluster was 42%, 34%, and 17%, respectively. The posterior probability for a sample belonging to a new cluster was about 6% in our analysis. We next applied the cluster profiling gene set enrichment analysis (see Hydra Method section) to each cluster using all genes from the pre-filtered expression matrix. Cluster 1 was enriched for adaptive immune response gene sets, cluster 2 was enriched for proliferative signaling gene sets, and cluster 3 was enriched for cancer-associated fibroblast gene sets (Figure 3B). Two of the three clusters were enriched for tumor microenvironment-associated expression. To further validate this signal, we correlated the hydra clusters with enrichment scores from the tumor microenvironment profiling tools xCell (Aran et al. 2017) and ESTIMATE (Yoshihara et al. 2013). Cluster 1 had higher average xCell enrichment scores associated with adaptive immune cell types including B-cells, CD4+ naive T-cells, and CD8+ naive T-cells (Kruskal-Wallis:  $p < 0.001$ ; Supplementary Tables 3-5). Cluster 2 was characterized by the absence of immune and stromal expression and higher tumor purity. The average ESTIMATE tumor purity for each cluster was 88%, 96% and 82%, respectively. Cluster 3 was enriched for fibroblast-associated expression by xCell analysis (Kruskal-Wallis:  $p < 0.001$ ). Clusters 1 and 3 had higher ESTIMATE immune-associated expression levels than cluster 2 (average ImmuneScore per cluster: 58, -612, 56), but cluster 3 had the highest stromal expression signature (average StromalScore per cluster: -1027, -1310, -135). We found no difference in patient survival outcomes across clusters (log-rank test,  $p > 0.05$ ). We investigated associations with clinical covariates, including mutation burden, age, and tumor content as assessed by a clinical pathologist, but found no statistically significant differences (Kruskal-Wallis:  $p > 0.05$ ; Supplementary Figure 2). We then investigated associations between the hydra clusters and neuroblastoma-associated molecular aberrations and clinical features (Supplementary Table 6). ATRX gene deletions were enriched in cluster 1 (Fisher's Exact Test:  $p < 0.05$ ). MKI low tumors were enriched in cluster 2 and 3 (Fisher's Exact Test:  $p < 0.01$ ). Chromosome 17 wild type tumors were enriched in clusters 2 and 3 (Fisher's Exact Test:  $p < 0.01$ ). Consensus clustering is a widely used approach for identifying tumor subtypes using gene expression data. We applied the M3C consensus clustering method, which is a more sophisticated version of consensus clustering that uses a null distribution to assess the statistical significance of the clustering (John et al. 2018; Wilkerson and Hayes 2010). M3C clustering of the MYCN-NA expression data using the 5000 genes with the largest median absolute deviation (MAD) resulted in the identification of two statistically significant clusters. We found that the M3C clusters correlated with the hydra clusters with lower ESTIMATE TumorPurity, but were not able to separate the adaptive immune cell and fibroblast infiltrated clusters. We also applied kmeans clustering using the gap statistic approach for estimating the number of clusters, but this approach did not identify any clusters (Tibshirani et al. 2001; Maechler et al. 2019). These results suggest that the hydra approach is more sensitive at detecting distinct tumor microenvironment states.

**N-of-1 Tumor Analysis for Pediatric Neuroblastoma** We investigated the predictive performance of the hydra approach for identifying clinically relevant signals in the N-of-1 tumor analysis setting. We obtained tumor gene expression data from five stage 4, MYCN-NA neuroblastoma samples (Figure 4). The age at diagnosis ranged from 2 to 6 years. Four out of five samples had a deletion in the ATRX gene. Samples 1D and 1R are diagnosis and resection samples from the same patient (Figure 4). Three of the ATRX-deleted samples clustered with the high immune expression cluster (cluster 1) and one clustered in the low immune, high proliferative signaling cluster (cluster 2). Hydra analysis assigned sample 1D to cluster 1 and sample 1R to cluster 2. The



resection sample 1R was extracted from lymph node tissue, which has a significantly different immune background than the training data. Another possible explanation for this change is that the tumor microenvironment is dynamic and the tumor may evade immune recognition as the disease progresses. We performed GSEA comparing samples 1D and 1R to investigate potential mechanisms leading to immune evasion in sample 1R. GSEA analysis found downregulation of the MHC Class I Antigen Processing & Presentation GO term in sample 1R (adjusted p-value  $\leq 0.002$ ). Loss of antigen processing functions is a common mechanism of immune evasion across cancer types (Reeves and James 2017). We obtained H&E slide images for these samples; the images were reviewed by a licensed pathologist and scored for evidence of inflammation. The N-of-1 predictive function of the hydra pipeline was used to classify samples into the subtypes discovered by the TARGET neuroblastoma analysis. Most of the samples (4 out of 5) clustered in cluster 1, which is characterized by higher immune marker expression. The hydra analysis agreed with the pathologist review in 4 out of 5 samples (Figure 4). Sample 4 was the only discordant sample. Sample 4 is also from lymph node tissue, which may have higher immune expression because of the tissue type. Notably, concordant ESTIMATE values were present in 3 out of 5 samples scored by the pathologist: samples 1D, 1R, and 2.

To further investigate expression patterns within the hydra-identified tumor microenvironment subtypes, we performed GSEA by z-score normalizing each tumor's gene expression data to its tumor microenvironment cluster. This approach revealed additional signal not present at the cohort level (Supplementary Figure 4). For example, enrichment of immune expression signatures within cluster 2 predicted differences in overall survival such that patients with higher immune expression had a better survival rate. Similarly, an elevated cell cycle signal within cluster 3 predicted worse survival compared to other cluster 3 samples with lower cell cycle expression. This approach provides a more appropriate background distribution for determining the significance of gene expression patterns and survival statistics.

**Hydra Analysis Discovers Complex Tissue Signatures** While the MYCN-NA neuroblastoma analysis focused on immune and fibroblast expression signatures, the hydra enrichment pipeline is unsupervised and is not restricted to immune or fibroblast signatures. For example, we applied the hydra enrichment analysis to the TARGET osteosarcoma cohort (N=74) and discovered enrichment of the GO striated muscle contraction term (FDR  $\leq 0.01$ ) (Supplementary Figure 5). Multivariate clustering for the GO striated muscle contraction gene set identified two clusters. xCell analysis of the osteosarcoma cohort found significant enrichment of skeletal muscle expression in the second cluster (Mann-Whitney U test,  $p \leq 0.001$ ). Surprisingly, the M3C clustering approach was not able to detect the strong muscle signature using the 5000 genes with the largest MAD ( $p \leq 0.05$ ). We identified a similar expression signal in an independent cohort of osteosarcoma tumor samples and subsequently confirmed with a licensed pathologist that the tumor sample did in fact contain muscle tissue. Explaining these sources of variation is necessary to derive clinically relevant conclusions from gene expression data. In addition to the muscle signature, we were able to identify innate immune and stromal subtypes in the TARGET osteosarcoma cohort that correlated with differences in overall survival rates (data not shown). Therefore, the hydra approach reveals important gene expression signatures reflecting cell content within the tumor sample and is widely applicable for revealing important gene expression signatures in complex tumor samples.

## Discussion

Nulla mi mi, venenatis sed ipsum varius, Table 1 volutpat euismod diam. Proin rutrum vel massa non gravida. Quisque tempor sem et dignissim rutrum. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Morbi at justo vitae nulla elementum commodo eu id massa. In vitae diam ac augue semper tincidunt eu ut eros. Fusce fringilla erat porttitor lectus cursus, vel sagittis arcu lobortis. Aliquam in enim semper, aliquam massa id, cursus neque. Praesent faucibus semper libero [3].

The hydra pipeline uses model-based clustering to identify recurrent expression patterns within cancer gene expression cohorts. We leveraged recent improvements in model-based clustering algorithms to identify differentially expressed genes without a matched normal distribution. We modeled differential expression as a multimodal Gaussian distribution using nonparametric Bayesian statistics. We then enriched for biologically-annotated Gene Ontology terms and performed multivariate clustering to reveal cancer subtyping expression signatures. The hydra framework can be used for identifying expression subtypes and classifying N-of-1 tumors. The hydra pipeline outperformed standard gene set enrichment tools for identifying overexpression of the MSigDB Hallmark cancer gene sets in synthetic cancer gene expression data and identified tumor microenvironment gene expression signatures in the TARGET pediatric neuroblastoma and osteosarcoma datasets that were not detected by consensus clustering analysis. Multivariate gene expression analysis is typically underpowered because the number of genes greatly exceeds the number of samples. To address this limitation, we propose selecting for multimodally expressed genes before performing multivariate analysis. The hydra multimodal filter reduces the number of genes and enriches for genes that participate in known biological processes, including those curated in the Gene Ontology database. As Yi Li et al. (2005) found in their study on unsupervised clustering of gene expression data, we also found that reducing expression data to multimodally expressed genes improves clustering of clinical subtypes. For example, multimodally expressed genes separate neuroblastoma subtypes by TumorMap analysis better than the standard approach of using all expressed genes. Furthermore, we showed that the hydra approach is more sensitive at resolving tumor microenvironment subtypes than the M3C consensus clustering approach. Significant progress has been made in subtyping neuroblastomas and adapting therapy for aggressive subtypes, but unexplained heterogeneity remains. Not accounting for this heterogeneity decreases the power of standard methods to detect important expression patterns. Identifying biomarkers using genome-wide technology may lead to improved risk stratification and the discovery of novel drug targets (Morgenstern et al. 2019). Hydra analysis of the TARGET MYCN-NA neuroblastoma cohort (N=70) found differential expression of tumor microenvironment markers, including markers of the adaptive immune response. Pediatric cancers are generally thought to be non-immunogenic because they have lower mutation burden than adult cancers, but the immunogenicity of pediatric cancer has not been sufficiently investigated (Majzner et al. 2017). Our analysis found significant variation in immune marker expression and identified ATRX deletions as a potential biomarker of immune infiltrated tumors. Further investigation into gene expression signatures and molecular aberrations that predict response to immunotherapy in pediatric cancers is needed. Hydra analysis may facilitate the development of novel therapies by grouping patients with similar tumor microenvironment properties. We found significant differences in immune and stromal expression that may inform precision medicine applications. The tumor microenvironment has become an important therapeutic consideration, but few methods account for the tumor microenvironment directly. Tumor purity has been identified as a confounding factor in cancer gene expression subtyping efforts (Rhee et al. 2018). For example, tumor purity and tumor microenvironment expression have been shown to

correlate with pancreatic cancer subtypes (Raphael et al. 2017). Furthermore, Aran et al. (2018) found that tumor purity was correlated with the mesenchymal glioblastoma subtype and recommended a differential expression approach to computationally remove the tumor purity signal. However, standard approaches for subtracting the tumor purity effect may not be the best approach because several mechanisms may influence tumor purity, and each mechanism may result in a different expression pattern. For instance, our analysis of MYCN-NA neuroblastoma identified two gene expression signatures that correlated with lower predicted tumor purity. Cluster 1 had an adaptive immune expression signature and cluster 3 had a cancer-associated fibroblast signature. Therefore, the estimated tumor purity signal should not be subtracted without first accounting for the different mechanisms influencing tumor purity. Precision oncology aims to differentiate tumors of the same diagnosis in order to match patients with the best treatment. We have developed the hydra method to discover subtle but recurrent expression patterns within a cancer type. Our approach may help to uncover the biology underlying tumor progression and response to therapy. We have shown that hydra is more sensitive than standard gene set enrichment approaches for detecting differential pathway expression. Additionally, we applied the unsupervised hydra analysis to pediatric neuroblastoma and osteosarcoma data and discovered distinct tumor microenvironment states. The hydra pipeline is a sensitive unsupervised clustering approach for N-of-1 tumor analysis and will facilitate pediatric precision oncology by discovering expression subtype signatures.

## Conclusion

CO<sub>2</sub> Maecenas convallis mauris sit amet sem ultrices gravida. Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec euismod ligula. Curabitur fringilla pulvinar lectus consectetur pellentesque. Quisque augue sem, tincidunt sit amet feugiat eget, ullamcorper sed velit.

Sed non aliquet felis. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Mauris commodo justo ac dui pretium imperdiet. Sed suscipit iaculis mi at feugiat. Ut neque ipsum, luctus id lacus ut, laoreet scelerisque urna. Phasellus venenatis, tortor nec vestibulum mattis, massa tortor interdum felis, nec pellentesque metus tortor nec nisl. Ut ornare mauris tellus, vel dapibus arcu suscipit sed. Nam condimentum sem eget mollis euismod. Nullam dui urna, gravida venenatis dui et, tincidunt sodales ex. Nunc est dui, sodales sed mauris nec, auctor sagittis leo. Aliquam tincidunt, ex in facilisis elementum, libero lectus luctus est, non vulputate nisl augue at dolor. For more information, see S1 Appendix.

## Supporting information

**S1 Fig. Bold the title sentence.** Add descriptive text after the title of the item (optional).

**S2 Fig. Lorem ipsum.** Maecenas convallis mauris sit amet sem ultrices gravida. Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec euismod ligula. Curabitur fringilla pulvinar lectus consectetur pellentesque.

**S1 File. Lorem ipsum.** Maecenas convallis mauris sit amet sem ultrices gravida. Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec euismod ligula. Curabitur fringilla pulvinar lectus consectetur pellentesque.

**S1 Video. Lorem ipsum.** Maecenas convallis mauris sit amet sem ultrices gravida. 465  
Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec euismod ligula. 466  
Curabitur fringilla pulvinar lectus consectetur pellentesque. 467

**S1 Appendix. Lorem ipsum.** Maecenas convallis mauris sit amet sem ultrices 468  
gravida. Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec 469  
euismod ligula. Curabitur fringilla pulvinar lectus consectetur pellentesque. 470

**S1 Table. Lorem ipsum.** Maecenas convallis mauris sit amet sem ultrices gravida. 471  
Etiam eget sapien nibh. Sed ac ipsum eget enim egestas ullamcorper nec euismod ligula. 472  
Curabitur fringilla pulvinar lectus consectetur pellentesque. 473

## Acknowledgments 474

Cras egestas velit mauris, eu mollis turpis pellentesque sit amet. Interdum et malesuada 475  
fames ac ante ipsum primis in faucibus. Nam id pretium nisi. Sed ac quam id nisi 476  
malesuada congue. Sed interdum aliquet augue, at pellentesque quam rhoncus vitae. 477

## References

## References

1. Conant GC, Wolfe KH. Turning a hobby into a job: how duplicated genes find new functions. Nat Rev Genet. 2008 Dec;9(12):938–950.
2. Ohno S. Evolution by gene duplication. London: George Alien & Unwin Ltd. Berlin, Heidelberg and New York: Springer-Verlag.; 1970.
3. Magwire MM, Bayer F, Webster CL, Cao C, Jiggins FM. Successive increases in the resistance of Drosophila to viral infection through a transposon insertion followed by a Duplication. PLoS Genet. 2011 Oct;7(10):e1002337.