

Anti-Deforestation Policies and Changes in the Clearings Pattern: Brazilian - Concise Report Version

João Pedro Vieira

December 7, 2018

Following Rosa et al. (2012) and Assunção et al. (2017), I am going to analyze how the deforestation pattern changed from 2002 through 2014 in the Legal Amazon. I will focus on three main elements: the trends of the deforestation by polygon size, the heterogeneity across state patterns, and the spatial distribution of the polygons over time. In 2004, the Brazilian Government implemented a satellite-based system that provides near-real-time alerts of deforestation (DETER) to inform law enforcement operations against illegal deforestation of tropical forest in the Amazon. However, this system can only detect clearings of a contiguous area bigger than 25 ha, so patches below this threshold are invisible to DETER and this information is public. Thus, the hypothesis is that DETER created an incentive to deforesters strategically adapt their behavior to deforest in small patches and thus impacted the distribution of the polygons size. Also, in 2008, there was a second turning point on conservation policies with the creation of a blacklist that allowed law enforcement to better target their efforts on municipalities with high rates of recent deforestation. Identifying a causal relationship between these policies changes and the deforestation trends is out of the scope of this report, what we are aiming to show here is a descriptive analysis of the deforestation pattern, considering time, state, and cleared patch size variation.

We will use the following spatial data (shapefiles):

1. Annual Deforestation Polygons [2002-2014]

It contains the annual deforestation series of the Brazilian Amazon. Multiple shapefiles, one for each Landsat mosaic scene.

We use the 2014 file because we can recover all the previous year information from this mask. The shapefiles contains polygons of other land cover categories besides deforestation that won't be used in this analysis, like forest and hydrography.

CRS: LongLat (coordinate system), SAD69_pre96_BRonly (datum), not projected

2. Legal Amazon and State Boundaries

Shapefile with the Legal Amazon Limits and the State Boundaries. Legal Amazon is geopolitical division of Brazil, includes the whole Brazilian Amazon Biome, part of the Cerrado and the Pantanal. More info about it: (https://en.wikipedia.org/wiki/Legal_Amazon)

CRS: LongLat (coordinate system), WGS84 (datum), not projected

1. Deforestation Trends by size of cleared patch

First of all, we are going to explore the evolution of the deforestation across the period of interest (2002-2014), splitting the deforestation increment into two categories: small polygons (contiguous area smaller than 25 hectares) and large polygons (contiguous area greater than 25 hectares). This analysis is inspired by:





Figure 1: Amazon deforestation by the size of cleared forest patch, 2002–2012 - (Assunção et al., 2017)



Notes: Reproducing Fig.1 from Assunção et al. (2017) expanding time period to include 2 more years. The figure illustrates annual Brazilian Amazon deforestation increment decomposed by size of cleared forest patch

As we can see in both graphs above, there was sharp falls in the deforestation after both anti-deforestation policies turning points (2004 and 2008), however as pointed out by (ref1) it seems to have impacted more the large polygons, increasing the relevance of small polygons. Linking this descriptive evidence with the fact that the satellite-based monitoring system can only detect the large cleared patches and that this limitation was public information, there might be a strategic behavior adaptation by deforesters explaining it. Our plot is a reproduction of (ref1) graph, but expanding to include two more years. This inclusion doesn't change the overall interpretation, only shows that the levels are somewhat constant, with a small increase similar





Figure 2: (a) Percentage and (b) area of deforested patches of different sizes in the Brazilian Amazon from 2002 through 2009. - (Rosa et al., 2012)

for both groups.

For completeness we also reproduce:

Deforestation Trends by Polygon Size


Inspired by Figure 2 - (a) - Rosa et al. (2012)



Notes: Reproducing Fig.2 (a) from Rosa et al. (2012) expanding time period to include 5 more years and different patches. The figure illustrates the share of deforested patches of different sizes in the Brazilian Amazon from 2002 through 2014.

Deforestation Trends by Polygon Size

Inspired by Figure 2 - (b) - Rosa et al. (2012)



Notes: Reproducing Fig.2 (b) from Rosa et al. (2012) expanding time period to include 5 more years and different patches. The figure illustrates area of deforested patches of different sizes in the Brazilian Amazon from 2002 through 2014.




Figure 3: Deforestation inside registered properties by property and cleared patch size, 2002–2012. - (Assunção et al., 2017))

These plots allows us to explore the trends for more categories and also look at the shares and not only the nominal increment. In our reproduction we expand it to include 5 more years and use the size intervals based on (ref1) to separate the policy relevant category of (<25ha). Looking at the shares we can see a sistematic increase in the share of small polygons (<25 ha) until 2010, after that there is a decrease but remaining as the majority proportion. The share is calculated as the ratio of the deforestation generated by the polygon size category and the total deforestation, for each year.

2. State Heterogeneity


Secondly, we want to explore the variation of these trends across the states, to check for the presence of outliers and heterogeneity. As argued by (ref1), regional heterogeneities require the policies to be tailored to maximize its efficiency. Inspired by:

In this case, we cannot reproduce the content of Fig. 3 entirely because we do not have assess to registered properties information, on the other hand, we expand their analysis to cover all the Legal Amazon States

instead of only two. In practice, we cannot link the polygon size with the type of agent (small, medium or large properties) but we can have a better notion of the heterogeneity of the deforestation by size.

Simple Plot of State Boundaries and UFs


For starters we do a exploratory plot, to show the Legal Amazon region and its states.



Plot of proportion of polygons by cleared patch through time by state.

For a complete notion of what's happening in each state we plot the proportion of the 4 sizes categories for each year.

Polygons Trends by State and Cleared Patch Size





We see that there are states such as AC and AP with almost all the deforestation coming from small polygons, contrasting with states like MT where larger polygons are more relevant. As expected for most of the states, small polygons increased their relevance becoming the majority, yet the presence of other patterns supports the idea of tailored policy to adjust the policies accordingly to the local pattern instead of treating the whole region as unique and uniform.

Maps of the proportion of small polygons across years

To complement the last plot we also focus only on the share of small polygons and the share of the State Deforested area to have a better visualization of the state-differences in each year.

Polygons Trends by State and Cleared Patch Size









Figure 4: Distribution of deforested patches of different sizes in the Brazilian Amazon for periods of (a) rapidly increasing deforestation (2002 through 2004) and (b) rapidly decreasing deforestation (2005 through 2009).- (Rosa et al., 2012)

From these visualizations we can conclude that there are persistent differences across the states for any given year for both measures, even though there is a common trends of less total deforestation and more presence of small cleared patches.


3. Spatial Distribution of deforestation by cleared patch size across time (2002-2014) - create function to just change the start and end year

Finally, we want to take advantage that we have spatial data at the polygon level to observe the spatial distribution of the deforestation by polygon size in different time periods, first a period of fast increase in deforestation rate (2002-2004), then a period of fast decrease (2005-2009) and lastly a period of relative stabilization (2010-2014). Inspired by:


Distribution of deforested patches by size (2002-2004)



Distribution of deforested patches by size (2005-2009)



Distribution of deforested patches by size (2010-2014)



Conclusion

References

Assunção, Juliano, Clarissa Gandour, Pedro Pessoa, and Romero Rocha. “Property-level assessment of change in forest clearing patterns: The need for tailoring policy in the Amazon.” *Land Use Policy* 66 (2017): 18-27.

Rosa, Isabel MD, Carlos Souza Jr, and Robert M. Ewers. “Changes in size of deforested patches in the Brazilian Amazon.” *Conservation Biology* 26, no. 5 (2012): 932-937.