

1 **A Bayesian Model of the DNA Barcode Gap**

2 Jarrett D. Phillips^{1,2*} (ORCID: 0000-0001-8390-386X), ... (others?)

3 ¹*School of Computer Science, University of Guelph, Guelph, ON., Canada, N1G2W1*

4 ²*Department of Integrative Biology, University of Guelph, Guelph, ON., Canada, N1G2W1*

5 ***Corresponding Author:** Jarrett D. Phillips¹

6 **Email Address:** jphill01@uoguelph.ca

7 **Running Title:** Bayesian inference for DNA barcode gap estimation

Abstract

A simple statistical model of the DNA barcode gap is outlined. Specifically, accuracy of recently introduced nonparametric metrics, inspired by coalescent theory, to characterize the extent of proportional overlap/separation in maximum and minimum pairwise genetic distances within and among species, respectively, is explored in both frequentist and Bayesian contexts. The empirical cumulative distribution function (ECDF) is utilized to estimate probabilities associated with positively skewed extreme tail distribution quantiles bounded on the closed unit interval $[0, 1]$ based on a straightforward binomial distance overlap count. Using R and the probabilistic programming language, Stan, the proposed maximum likelihood estimators and Bayesian model are demonstrated on CYTB gene sequences from two *Agabus* diving beetle species exhibiting limits in the extent of representative taxonomic sampling. Large-sample theory and MCMC simulations show much uncertainty in parameter estimates, particularly when specimen sample sizes for target species are small. Findings herein highlight the promise of the Bayesian approach using a conjugate beta prior for reliable posterior uncertainty estimation when available data are sparse. Obtained results can shed light on foundational and applied research questions concerning DNA-based specimen identification and species delineation for studies in evolutionary biology and ecology, as well as biodiversity conservation and management, of wide-ranging taxa.

Keywords: Bayesian/frequentist inference, DNA barcoding, intraspecific genetic distance, interspecific genetic distance, specimen identification, species discovery

1 Introduction

The routine use of DNA sequences to support broad evolutionary hypotheses and questions concerning demographic processes, like gene flow and speciation, is now ubiquitous. Since the late 1980s, mitochondrial DNA (mtDNA) has been employed to produce a

34 distinctive and measurable pattern of genetic polymorphism in diverse and
35 spatially-distributed taxonomic lineages such as birds, fishes, insects, and arachnids, among
36 other extensively studied groups (Avice et al., 1987). The application of genomic data to
37 applied fields like biodiversity forensics, conservation, and management for the molecular
38 identification of unknown specimen samples came later (*e.g.*, Forensically Informative
39 Nucleotide Sequencing (FINS); Bartlett and Davidson (1992)). Since its inception over 20
40 years ago, DNA barcoding (Hebert et al., 2003a,b) built significantly on earlier work and has
41 emerged as a robust method of specimen identification and species discovery across myriad
42 multicellular eukaryotes which have been sequenced at easily obtained short, standardized
43 gene regions like the cytochrome *c* oxidase subunit I (5'-COI) mitochondrial locus for animals.
44 However, DNA barcoding is not without its controversies.

45 The success of the single-locus approach, particularly for regulatory and forensic
46 applications, depends crucially on two important factors: (1) the availability of high-quality
47 specimen records found in public reference sequence databases such as the Barcode of Life
48 Data Systems (BOLD; <http://www.barcodinglife.org>) (Ratnasingham and Hebert, 2007) and
49 GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>), and (2) the establishment of a DNA
50 barcode gap — the notion that the maximum genetic distance observed within species is
51 much smaller than the minimum degree of marker variation found among species (Meyer
52 and Paulay, 2005; Meier et al., 2008). Early work has demonstrated that the presence of
53 a DNA barcode gap hinges strongly on extant levels of species haplotype diversity gauged
54 from comprehensive specimen sampling at wide geographic and ecological scales (Bergsten
55 et al., 2012; Čandek and Kuntner, 2015). Despite this, many taxonomic groups lack adequate
56 separation in their pairwise intraspecific and interspecific genetic distances due to varying
57 rates of evolution in both genes and taxa (Pentinsaari et al., 2016). Furthermore, it has
58 been well-demonstrated that the presence of a DNA barcode gap becomes less certain with
59 increasing spatial scale of sampling, since interspecific distances increase due to population
60 structuring under limited dispersal through isolation-by-distance, while intraspecific distances

shrink as more closely-related species are sampled (Phillips et al., 2022). This can pose problems in cases of rare species or monotypic taxa, for instance (Ahrens et al., 2016). As a result, rapid matching of unknown samples to expertly-validated references can be compromised, leading to cases of false positives (taxon oversplitting) and false negatives (excessive lumping of taxa) due to incomplete lineage sorting, interspecies hybridization, genome introgression, species synonymy, cryptic species diversity, and misidentifications (Hubert and Hanner, 2015; Phillips et al., 2022). Another culprit includes PCR/sequencing error caused by incorrect nucleotide basecalling, chimeras/heteroplasmies, sample contamination, indels (insertion/deletions), NUMT (nuclear-mitochondrial insert) pseudogene amplification, or *Wolbachia* endosymbiont infection, among other explanations (Phillips et al., 2023). Each of these can artificially magnify levels of genetic diversity, rendering techniques like DNA barcoding unreliable for certain groups like marine invertebrates which are both undersampled and possess slow rates of molecular evolution.

Recent work has argued that DNA barcoding, in its current form, is lacking in statistical rigor (Phillips et al., 2022). Most studies rely strongly on heuristic distance-based thresholds calculated from convenient specimen sample sizes, rather than on optimal sampling designs using collated taxon haplotype information (Phillips et al., 2015, 2020, 2022), to infer taxonomic identity and delineate species boundaries. Of these studies, few report measures of uncertainty, such as standard errors (SEs) and confidence intervals (CIs), around estimates of intraspecific and interspecific variation (*e.g.*, Wiemers and Fiedler (2007)), calling into question the existence of a true species' DNA barcode gap (Čandek and Kuntner, 2015; Phillips et al., 2022). To support this idea, novel nonparametric locus- and species-specific metrics based on the multispecies coalescent (MSC) were recently outlined by Phillips et al. (2024). The basic coalescent (Kingman, 1982a,b) encompasses a backwards continuous-time stochastic Markov process of allelic sampling among lineages under the infinite sites model of mutation within a large neutrally-evolving, panmictic, diploid species population of constant size from the present into the past towards the most recent common ancestor (MRCA).

Although the coalescent sampling process plays a foundational role in evolutionary biology and population genetics theory due to its inherent simplicity and flexibility, the approach has been both under-utilized and under-appreciated as a key player within DNA barcoding (Collins and Cruickshank, 2014; Downton et al., 2014; Hubert and Hanner, 2015; Phillips et al., 2022). Numerous theoretical and empirical investigations have shown that protein-coding sequence variation within haploid loci like COI at nonsynonymous sites is generally subject to strong purifying selection and selective sweeps, rather than the occurrence of functionally silent substitutions within synonymous regions (Stoeckle and Thaler, 2014). In effect, mutations become diagnostic of species at a much faster rate compared to nuclear regions of the genome. Previously proposed MSC algorithmic approaches (of which there are too many to exhaustively list here, and which have mostly been applied to multilocus sequence data), generally assume a strict molecular clock and a simplified model of DNA sequence evolution across closely-related taxa, from which an estimated species phylogeny may be constructed (*e.g.*, with or without use of a guide tree) (*e.g.*, Rannala and Yang (2003, 2017); Yang and Rannala (2010, 2014, 2017)). Even tree-based taxon delimitation methods designed for single loci, such as the Generalized Mixed Yule Coalescent (GMYC) (Pons et al., 2006; Monaghan et al., 2009; Fujisawa and Barraclough, 2013) and Poisson Tree Processes (PTP) (Zhang et al., 2013; Kapli et al., 2017), which have seen much use in DNA barcoding initiatives, have their flaws. Optimal performance of these methods relies heavily on careful parameter selection (*e.g.*, prior specification of branching rates in ultrametric trees in the case of GMYC) (Hubert et al., 2024) within third party software like BEAST (Bouckaert et al., 2019), among other concerns (Talavera et al., 2013; Fonseca et al., 2021). In contrast, Phillips et al.’s (2024) approach is tree-free and does not require judicious parameter setting. Therefore, the method is extremely efficient and fast to run.

The DNA barcode gap statistics have been shown to hold strong promise for reliable DNA barcode gap assessment when applied to predatory *Agabus* (Coleoptera: Dytiscidae) diving beetles (Phillips et al., 2024). Despite their ease of sampling and well-established taxonomy,

this group possesses few morphologically-distinct taxonomic characters that readily facilitate their assignment to the species level (Bergsten et al., 2012). Accurate taxon discrimination within *Agabus* is paramount since several species are currently classified as endangered on the International Union for Conservation of Nature (IUCN) Red List (<https://www.iucnredlist.org>). Further, the proposed metrics indicate that sister species pairs from this taxon are often difficult to distinguish on the basis of their DNA barcode sequences (Phillips et al., 2024), which may be caused by *Agabus*'s increased degree of non-monophyly at wider geospatial scales, occurrence of hybridization/introgression, or presence of synonymy (Bergsten et al., 2012). Using sequence data from three mitochondrial cytochrome markers (5'-COI, 3'-COI, and cytochrome *b* (CYTB)) obtained from BOLD and GenBank, results from Phillips et al. (2024) highlight that DNA barcoding has been a one-sided argument. Phillips et al.'s (2024) findings point to the need to balance both the sufficient collection of specimens, as well as the extensive sampling of species. Not surprisingly, DNA barcode libraries for most taxonomic groups are biased toward the latter effort (Phillips et al., 2015, 2019, 2022).

The estimators from Phillips et al. (2024) represent a clear improvement over simple, yet arbitrary, distance heuristics such as the 2% rule noted by Hebert et al. (2003a) and the 10 $\times$ rule (Hebert et al., 2004) that form the core of threshold-based single-locus species delimitation tools like Automatic Barcode Gap Discovery (ABGD) (Puillandre et al., 2011), Assemble Species by Automatic Partitioning (ASAP) (Puillandre et al., 2021), and the Barcode Index Number (BIN) framework (Ratnasingham and Hebert, 2013). The 2% rule asserts that DNA barcodes differing by at least 2% at sequenced genomic regions should be expected to originate from different biological species, whereas the 10 $\times$ rule suggests that barcode sequences displaying 10 times more genetic variation among species than within taxa is suggestive of a distinct evolutionary origin. However, the lack of adoption of an explicit and universally agreed upon species concept in DNA barcoding studies that readily governs lineage formation and proliferation necessary to establish rigorous taxon definitions for

successful delimitation of hypothesized and heuristic evolutionary units using these well-known criteria, in conjunction with secondary lines of evidence (*e.g.*, morphology, ecology, geography, ontogeny, and behaviour) promised by an integrative framework (Dayrat, 2005), is missing (de Queiroz, 2007; Rannala, 2015; Pante et al., 2015; Wells et al., 2022). This is the case, for instance, with species existing in allopatry/sympatry, where establishment of reproductive isolation on the basis of fixed diagnostic differences using the Biological Species Concept (BSC) (Mayr, 1942), is challenging (Fujita et al., 2012). In addition, the reliance on visualization approaches, such as frequency histograms, dotplots, and quadrant plots, to expose DNA barcoding’s limitations, has also been criticized (Collins and Cruickshank, 2013; Phillips et al., 2022). Up until the work of Phillips et al. (2024), the majority of studies (*e.g.*, Young et al. (2021)) have treated the DNA barcode gap as a binary response. Given poor sampling depth for most taxa, a Yes/No dichotomy is inherently flawed because it can falsely imply a DNA barcode gap is present for a taxon of interest when in fact no such separation in genetic distances exists. On the other hand, in the case of sequence overlap, while the presence a single specimen record is a necessary requirement to nullify the existence of a DNA barcode gap in light of available data, it is not a sufficient condition, since said taxon record could represent a species misidentification or other error. Thus, the MSC represents an ideal model to settle ongoing debates and issues regarding DNA barcoding’s use as both a molecular identification and delineation tool. Notably, Fujita et al. (2012) remarks that the MSC applied to the species delimitation problem harmonizes many species concepts since it identifies independent evolutionary trajectories under the umbrella of the General Lineage Concept. The recently introduced DNA barcode gap metrics go some way into operationalizing this task.

Phillips et al.’s (2024) proposed statistics quantify the extent of asymmetric directionality of proportional distance distribution overlap/separation for species within well-sampled taxonomic genera based on a straightforward distance count, in a similar vein to established measures of statistical similarity such as the Kullback-Leibler (KL) divergence (Kullback

169 and Leibler, 1951) and other related statistics. The metrics can be employed in a variety
 170 of ways, including to validate performance of marker genes for specimen identification to
 171 the species level (using nonparametric bootstrapping, as in Phillips et al. (2024)). The
 172 estimators can further be utilized to assess whether computed values are consistent with
 173 fundamental population genetic-level parameters like effective population size (N_e), mutation
 174 rates (μ), migration rates (m), and divergence times (τ), as well as derived ones, such as the
 175 population mutation rate $\theta = 4N_e\mu$ and the fixation index, F_{ST} , for species under study in
 176 a statistical phylogeographic setting (Knowles and Maddison, 2002; Knowles, 2009; Mather
 177 et al., 2019). Early on, species discovery using DNA barcoding was presumed to only work
 178 for reciprocally monophyletic groups, and thus concerned itself with terminal branches of
 179 generated phylogenies rather than more basal lineages occurring deeper in hypothesized
 180 species trees (Fujita et al., 2012; Mutanen et al., 2016). Furthermore, the occurrence of
 181 short branches and large N_e within resolved phylogenies increases the probability of deep
 182 coalescence, clouding species delimitations, which often fail or are uncertain in broad
 183 parameter space (Carstens et al., 2013; Hickerson et al., 2006; Rannala, 2015). As DNA
 184 barcoding is a single-locus approach, it is problematic for evolutionarily young taxa, wherein
 185 incomplete lineage sorting within gene genealogies is a common phenomenon due to the
 186 ongoing stochastic dynamic of mutation generating population variation, and genetic drift
 187 driving gene variants to fixation (Rannala, 2015). The most promising way forward in
 188 assessing the validity of Phillips et al.’s (2024) method seems to be through the use of
 189 software such as BPP (Bayesian Phylogenetics and Phylogeography), which permits efficient
 190 full Bayesian simulations under various MSC models (*e.g.*, MSC-I (MSC with introgression)
 191 or MSC-M (MSC with migration), among others) using MCMC for tree parameter estimation
 192 (via the A00 option, for instance) (Flouri et al., 2018), or PHRAPL (Phylogeographic
 193 Inference using Approximate Likelihoods) (Jackson et al., 2017a,b), which employs tractable
 194 phylogenetic likelihood calculations via the genealogical divergence index (gdi). Other
 195 hypothesis-driven approaches to species delimitation model selection, like Bayes Factors

(BFs) (Leaché et al., 2014), are also possible.

While introduction of the new metrics is a step in the right direction, what appears to be missing is a rigorous statistical treatment of the DNA barcode gap. This includes an unbiased way to compute the statistical accuracy of Phillips et al.’s (2024) estimators arising through problems inherent in frequentist maximum likelihood estimation for probability distributions having bounded positive support on the closed unit interval $[0, 1]$. To this end, here, a Bayesian model of the DNA barcode gap coalescent is introduced to rectify such issues. The model allows accurate estimation of posterior means, posterior standard deviations (SDs), posterior quantiles, and credible intervals (CrIs) for the statistics given datasets of intraspecific and interspecific distances for species of interest.

2 Methods

2.1 DNA Barcode Gap Metrics

The novel nonparametric maximum likelihood estimators (MLEs) of proportional overlap/separation between intraspecific and interspecific distance distributions for a given species (x) to aid assessment of the DNA barcode gap are as follows:

$$p_x = \frac{\#\{d_{ij} \geq a\}}{\#\{d_{ij}\}} \quad (1)$$

$$q_x = \frac{\#\{d_{XY} \leq b\}}{\#\{d_{XY}\}} \quad (2)$$

$$p'_x = \frac{\#\{d_{ij} \geq a'\}}{\#\{d_{ij}\}} \quad (3)$$

$$q'_x = \frac{\#\{d'_{XY} \leq b\}}{\#\{d'_{XY}\}} \quad (4)$$

where d_{ij} are distances within species, d_{XY} are distances among species for an *entire* genus of concern, and d'_{XY} are combined interspecific distances for a target species and its closest

neighbouring species. The notation $\#$ reflects a count. Quantities a , a' , and b correspond to $\min(d_{XY})$, $\min(d'_{XY})$, and $\max(d_{ij})$, the minimum interspecific distance, the minimum combined interspecific distance, and the maximum intraspecific distance, respectively (Figure 1).




Figure 1: Modified depiction from Meyer and Paulay (2005) and Phillips et al. (2024) of the overlap/separation of intraspecific and interspecific distances (δ) for calculation of the DNA barcode gap metrics (p_x and q_x) for a hypothetical species x . The minimum interspecific distance is denoted by a and the maximum intraspecific distance is indicated by b . The quantity $f(\delta)$ is akin to a kernel density estimate of the probability density function of distances. A similar visualization can be displayed for p'_x and q'_x within the interval $[a', b]$.

Hence, Equations (1)-(4) are simply empirical partial means of distances falling at and below, or at and exceeding, given distribution thresholds. Notice further that a/a' , and b are also the first and n th order statistics, $X_{(1)}$ and $X_{(n)}$, respectively, with $a/a' < b$, which have been pointed out by Phillips et al. (2022) as important for developing a statistical theory to test for the existence of the DNA barcode gap. Equations (1)-(4) can also be expressed in terms of empirical cumulative distribution functions (ECDFs) (see next section). A clear connection between the ECDF and the nonparametric MLE was noted by Efron and Tibshirani (1993) as well as others. Distances form a continuous distribution and are easily computed from a model of DNA sequence evolution, such as uncorrected or corrected p-distances (Jukes and Cantor, 1969; Kimura, 1980) using, for example, the `dist.dna()` function available in the `ape` R package (Paradis et al., 2004). However, calculated values are *not* independent and

228 identically distributed (IID) because estimated standard errors (SEs) will depend on both the
 229 number of species sampled within the genus under study, as well as the number of specimens
 230 sampled within a target species. In Phillips et al. (2024), To tease this out, Phillips et al.
 231 (2024) suggests plotting estimator values against their estimated SEs, along with a simple
 232 random downsampling scheme. In the case of two species comprising a focal genus, one well
 233 sampled and the other poorly sampled, values of the metrics close to zero for the sufficiently
 234 sampled species will likely possess larger SEs following downsizing to match the number of
 235 poorly sampled specimens (Phillips et al., 2024). The approach of Phillips et al. (2024) differs
 236 markedly from the traditional definition of the DNA barcoding gap laid out by Meyer and
 237 Paulay (2005) and Meier et al. (2008) in that the proposed metrics incorporate interspecific
 238 distances which *include* the target species of interest. Furthermore, if a focal species is
 239 found to have multiple nearest neighbours, then the species possessing the smallest average
 240 distance is used (Phillips et al., 2024). These schemes more accurately account for species'
 241 coalescence histories inferred from contemporaneous samples of DNA sequences leading to
 242 instances of barcode sequence sharing, such as interspecific hybridization/introgression events
 243 (Phillips et al., 2024). For example, within Equations (3) and (4), the degree of distance
 244 distribution overlap/separation between a target taxon and its nearest neighbouring species,
 245 gauged from magnitudes of p'_x and q'_x , is directly proportional to the amount of time in
 246 which the two lineages diverged from the MRCA (Phillips et al., 2024). More specifically,
 247 $a \propto t_{XY, \min}$ and $b \propto t_{ij, \max}$, the minimum and maximum (unobserved) divergence time (in
 248 N_e generations), respectively, for two different individuals (i, j) and species (X, Y) within
 249 a well-inventoried genus (Phillips et al., 2024). Figure 1 in Phillips et al. (2024) depicts a
 250 phylogeny for two hypothetical species, A and B , showing paraphyly based on the sequencing
 251 of two mitochondrial loci. The (unobserved) time the two species formed, t_{AB} , is intermediate
 252 between $t_{XY, \min}$ and $t_{ij, \max}$, but theoretically, its maximum ($t_{AB, \max}$) may occur more
 253 distantly in the past closer to the root comprising the MRCA for all sampled taxa (Phillips
 254 et al., 2024). From a coalescence perspective, p_x is the probability two lineages within species

²⁵⁵ x coalesce after $t_{XY, \min}$, whereas q_x represents the probability a lineage within species s
²⁵⁶ coalesces with a lineage in species k (within a sample) after $t_{XY, \min}$, but before $t_{ij, \max}$. Thus,
²⁵⁷ the quantities can be used as a criterion to assess the failure of DNA barcoding in recently
²⁵⁸ radiated taxonomic groups, among other plausible biological explanations mentioned earlier.
²⁵⁹ Note, distances are constrained to the interval $[0, 1]$, whereas the metrics are defined only
²⁶⁰ on the interval $[a/a', b]$. Values of the estimators obtained from Equations (1)-(4) close to
²⁶¹ or equal to zero give evidence for separation between intraspecific and interspecific distance
²⁶² distributions; that is, values suggest the presence of a DNA barcode gap for a target species.
²⁶³ Conversely, values near or equal to one give evidence for distribution overlap; that is, values
²⁶⁴ likely indicate the absence of a DNA barcode gap.

²⁶⁵ **2.2 The Model**

²⁶⁶ Before delving into the derivation of the proposed DNA barcode gap metrics, review of
²⁶⁷ some fundamental statistical theory is necessary.

²⁶⁸ For a given random variable X from a given distribution F , its cumulative distribution
²⁶⁹ function (CDF) is defined by

$$F_X(t) = \mathbb{P}(X \leq t) = 1 - \mathbb{P}(X > t). \quad (5)$$

²⁷⁰ Rearranging Equation (5) gives

$$\mathbb{P}(X > t) = 1 - F_X(t) = 1 - \mathbb{P}(X \leq t), \quad (6)$$

²⁷¹ from which it follows that

$$\mathbb{P}(X \geq t) = 1 - F_X(t) + \mathbb{P}(X = t). \quad (7)$$

272 The CDF has many desirable statistical properties such as boundedness, continuity, and
 273 monotonicity. Consult any graduate level text on mathematical statistics for reference.

274 Equations (1)-(4) can thus be written in terms of ECDFs via Equations (5) and (7) as
 275 follows, since the true underlying F s, are unknown *a priori*, and therefore must be estimated
 276 using available sequence distance data:

$$\begin{aligned} p_x &= \mathbb{P}(d_{ij} \geq a) \\ &= 1 - \hat{F}_{d_{ij}}(a) + \mathbb{P}(d_{ij} = a) \end{aligned} \quad (8)$$

$$\begin{aligned} q_x &= \mathbb{P}(d_{XY} \leq b) \\ &= \hat{F}_{d_{XY}}(b) \end{aligned} \quad (9)$$

$$\begin{aligned} p'_x &= \mathbb{P}(d_{ij} \geq a') \\ &= 1 - \hat{F}_{d_{ij}}(a') + \mathbb{P}(d_{ij} = a') \end{aligned} \quad (10)$$

$$\begin{aligned} q'_x &= \mathbb{P}(d'_{XY} \leq b) \\ &= \hat{F}_{d'_{XY}}(b). \end{aligned} \quad (11)$$

277 Given n increasing-ordered data points, the (discrete) ECDF, $\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{[x_i \leq t]}$,
 278 comprises an increasing step function having jump discontinuities of size $\frac{1}{n}$ at each sample
 279 observation (x_i) , excluding ties (or steps of weight $\frac{i}{n}$ with duplicate observations), where
 280 $\mathbb{1}(x)$ is the indicator function. Note that $1 - \hat{F}_n(t)$ is the complementary ECDF (CECDF),
 281 which is a declining function. Further, notice that $\mathbb{P}(X = t) \neq 0$. Equations (8)-(11)
 282 clearly demonstrate the asymmetric directionality of the proposed metrics. Plots of the
 283 ECDFs/CECDFs for nonzero intraspecific, interspecific, and combined interspecific distances,

superimposed with corresponding values of a/a' and b , provide a new, useful way to visualize the DNA barcode gap. Firstly, since there is no need to tune the output (*e.g.*, number of bins and bin width in the case of histograms). Secondly, probabilities reflecting areas under the distance curves depicted in **Figure 1** can be directly read off the ECDF/CECDF curves. Values for q_x and q'_x can also be determined from direct inspection of the ECDF/CECDF plot by determining the corresponding y -value (probability) for a given value of x (distance). Furthermore, calculation of the DNA barcode gap estimators is convenient as they implicitly account for total distribution area (including overlap).

A major criticism of large sample (frequentist) theory is that it relies on asymptotic properties of the MLE (whose population parameter is assumed to be a fixed but unknown quantity), such as estimator normality and consistency as the sample size approaches infinity. This problem is especially pronounced in the case of binomial proportions (Newcombe, 1998). The estimated Wald SE of the sample proportion, is given by $\widehat{SE}[\hat{p}] = \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$, where $\hat{p} = \frac{Y}{n}$ is the MLE, Y is the total number of successes ($Y = \sum_{i=1}^n y_i$), and n is the total number of trials (*i.e.*, sample size). However, the above formula for the standard error is problematic for several reasons. First, it is a Normal approximation which makes use of the central limit theorem (CLT); thus, large sample sizes are required for reliable estimation. When few observations are available, SEs will be large and inaccurate, leading to low statistical power to detect a true DNA barcode gap when one actually exists. Further, resulting interval estimates could span values less than zero or greater than one, or have zero width, which is practically meaningless. Second, when proportions are exactly equal to zero or one, resulting SEs will be exactly zero, rendering $\widehat{SE}[\hat{p}]$ given above completely useless. In the context of the proposed DNA barcode gap metrics, values obtained at the boundaries of their support are often encountered. Therefore, reliable calculation of SEs is not feasible. Given the importance of sufficient sampling of species genetic diversity for DNA barcoding initiatives, a different statistical estimation approach is necessary.

Bayesian inference offers a natural path forward in this regard since it allows for

straightforward specification of prior beliefs concerning unknown model parameters and permits the seamless propagation of uncertainty, when data are lacking and sample sizes are small, through integration with the likelihood function associated with true generating processes. The posterior distribution ($\pi(\theta|Y)$) is given by Bayes' theorem up to a proportionality $\pi(\theta|Y) \propto \pi(Y|\theta)\pi(\theta)$, where θ are unobserved parameters, Y are known data, $\pi(Y|\theta)$ is the likelihood, and $\pi(\theta)$ is the prior. As a consequence, because parameters are treated as random variables, Bayesian models are much more flexible and generally more easily interpretable compared to frequentist approaches. Under the Bayesian paradigm, entire posterior distributions, along with their summaries (*e.g.*, CrIs) are outputted, rather than just long run behaviour reflected in sampling distributions, p-values, and CIs as in the frequentist case, thus allowing direct probability statements to be made.

Essentially, from a statistical perspective, the goal herein is to nonparametrically estimate probabilities corresponding to extreme tail quantiles for positive highly skewed distributions on the unit interval (or any closed subinterval thereof). Here, it is sought to numerically approximate the extent of proportional overlap/separation of intraspecific and interspecific distance distributions within the subinterval $[a/a', b]$. This is a challenging computational problem within the current study as detailed in subsequent sections. The usual approach employs kernel density estimation (KDE), along with numerical or Monte Carlo integration of complicated probability distribution functions (PDFs), and invocation of extreme value theory (EVT); however, this requires careful selection of the bandwidth parameter, among other considerations. This becomes problematic when fitting finite mixture models where nonidentifiability is rampant. For DNA barcode gap estimation, this would correspond to a two-component mixture (one for intraspecific distance comparisons, and the other for interspecific comparisons), with one or more distribution modes or curve intersection points between components, and the presence of zero distance inflation. This makes parameter estimation difficult using methods like the Expectation-Maximization (EM) algorithm (Dempster et al., 1977) as the algorithm may become stuck in suboptimal regions of the

parameter search space and prematurely converge to local optima. This seems wholly plausible, especially given that the genus-level distribution of genetic distances is a mixture of closely-related (*i.e.*, nearest neighbour) and more distant species. Here, for simplicity, an alternate route is taken to avoid these obstacles.

Counts, y , of overlapping distances (as expressed in the numerator of Equations (1)-(4)) are treated as binomially distributed with expectation $\mathbb{E}[Y|\theta] = k\theta$, and variance $\mathbb{V}[Y|\theta] = k\theta(1 - \theta)$, where $k = \{N, C\}$ are total count vectors of intraspecific and combined interspecific distances, respectively, for a target species along with its nearest neighbour species, and $k = M$ is a total count vector for all interspecific species comparisons. This follows from the fact that the ECDF/CECDF is binomially distributed. The quantity thus being estimated is the parameter vector $\underline{\theta} = \{p_x, q_x, p'_x, q'_x\}$. The metrics encompassing $\underline{\theta}$ are presumed to follow a Beta(α, β) distribution, with real shape parameters α and β , which is a natural choice of prior on probabilities. The beta distribution has a prior mean of $\mathbb{E}[\theta|\alpha, \beta] = \frac{\alpha}{\alpha+\beta}$ and a prior variance equal to $\mathbb{V}[\theta|\alpha, \beta] = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$. In the case where $\alpha = \beta$, all generated Beta(α, β) distributions will possess the same prior expectation, whereas the prior variance will shrink as both α and β increase. Such a scheme is quite convenient since the beta distribution is conjugate to the binomial distribution. Thus, the posterior distribution is also beta distributed, specifically, Beta($\alpha + Y, \beta + n - Y$), having expectation $\mathbb{E}[\theta|Y, \alpha, \beta] = \frac{\alpha+Y}{\alpha+\beta+n}$ and variance $\mathbb{V}[\theta|Y, \alpha, \beta] = \frac{(\alpha+Y)(\beta+n-Y)}{(\alpha+\beta+n)^2(\alpha+\beta+n+1)}$. In the context of DNA barcoding, it is important that the DNA barcode gap metrics effectively differentiate between extremes of no overlap/complete separation and complete overlap/no separation, corresponding to values of the metrics equal to 0 and 1 (equivalent to total distance counts of 0 and n), respectively. These extremes yield a posterior expectation of $\mathbb{E}[\theta|Y = 0, \alpha, \beta] = \frac{\alpha}{\alpha+\beta+n}$ and a posterior variance of $\mathbb{V}[\theta|Y = 0, \alpha, \beta] = \frac{\alpha(\beta+n)}{(\alpha+\beta+n)^2(\alpha+\beta+n+1)}$ and $\mathbb{E}[\theta|Y = n, \alpha, \beta] = \frac{\alpha+n}{\alpha+\beta+n}$ and $\mathbb{V}[\theta|Y = n, \alpha, \beta] = \frac{(\alpha+n)\beta}{(\alpha+\beta+n)^2(\alpha+\beta+n+1)}$. Note, the posterior variances are equivalent at these thresholds for all $\alpha = \beta$.

Parameters were given an uninformative Beta(1, 1) prior, which is identical to a standard

365 uniform ($\text{Uniform}(0, 1)$) prior since it places equal probability on all parameter values within
 366 its support. This distribution has an expected value of $\mu = \frac{1}{2} = 0.500$ and a variance of
 367 $\sigma^2 = \frac{1}{12} \approx 0.083$. Further, the posterior is $\text{Beta}(Y+1, n-Y+1)$, from which various moments
 368 such as the expected value $\mathbb{E}[\theta|Y] = \frac{Y+1}{n+2}$ and variance $\mathbb{V}[\theta|Y] = \frac{(Y+1)(n-Y+1)}{(n+2)^2(n+3)}$, and other
 369 quantities, can be easily calculated. Clearly, $\mathbb{E}[\theta|Y=0] = \frac{1}{n+2}$ and $\mathbb{V}[\theta|Y=0] = \frac{n+1}{(n+2)^2(n+3)}$,
 370 and $\mathbb{E}[\theta|Y=n] = \frac{n+1}{n+2}$ and $\mathbb{V}[\theta|Y=n] = \frac{n+1}{(n+2)^2(n+3)}$. For example, with $n = 2$ sampled
 371 specimens for a given species (the minimum sample size possible) having no overlapping
 372 genetic distances, on average, θ will be equal to $\mu = \frac{1}{4} = 0.250$, with a variance of
 373 $\sigma^2 = \frac{3}{80} \approx 0.0375$. In contrast, with $n = 100$ individuals and no overlap, the mean reduces to
 374 $\mu = \frac{1}{102} \approx 0.001$, with a variance of $\sigma^2 = 9.425 \times 10^{-5}$. When $n = 2$ and both taxon distances
 375 overlap, the mean is $\mu = \frac{3}{4} = 0.750$, having a variance of $\sigma^2 \approx 0.0375$. For $n = 100$ where all
 376 distances overlap, θ has expectation $\mu = \frac{101}{102} \approx 0.990$ and variance $\sigma^2 \approx 9.425 \times 10^{-5}$. These
 377 edge case examples demonstrate that the $\text{Beta}(1, 1)$ distribution serves as a good first-pass
 378 prior for DNA barcode gap estimation using Phillips et al.'s (2024) metrics, though tighter
 379 bounds can likely be achieved. In general however, when possible, it is always advisable
 380 to incorporate prior information, even if only weak, rather than simply imposing complete
 381 ignorance in the form of a flat prior distribution. In the case of unimodal distributions, the
 382 (estimated) posterior mean often possesses the property that it readily decomposes into a
 383 convex linear combination, in the form of a weighted sum, of the (estimated) prior mean
 384 and the MLE. That is $\hat{\mu}_{\text{posterior}} = w\hat{\mu}_{\text{prior}} + (1-w)\hat{\mu}_{\text{MLE}}$, where for the beta distribution,
 385 $w = \frac{\alpha+\beta}{\alpha+\beta+n}$. Therefore, with sufficient data, $w \rightarrow 0$ as $n \rightarrow \infty$, regardless of the values of α
 386 and β , and the choice of prior distribution becomes less important since the posterior will be
 387 dominated by the likelihood. For the $\text{Beta}(1, 1)$, $w = \frac{2}{2+n}$, with $n = 2$ giving $w = \frac{1}{2} = 0.500$;
 388 that is, the posterior is the arithmetic average of the prior and the likelihood. For $n = 100$,
 389 the prior weight is $w = \frac{2}{102} \approx 0.0196$. The full Bayesian model for species x is thus given by

$$\begin{aligned}
y_{\text{lwr}} &\sim \text{Binomial}(N, p_{\text{lwr}}) \\
y_{\text{upr}} &\sim \text{Binomial}(M, p_{\text{upr}}) \\
y'_{\text{lwr}} &\sim \text{Binomial}(N, p'_{\text{lwr}}) \\
y'_{\text{upr}} &\sim \text{Binomial}(C, p'_{\text{upr}}) \\
p_{\text{lwr}}, p_{\text{upr}}, p'_{\text{lwr}}, p'_{\text{upr}} &\sim \text{Beta}(1, 1).
\end{aligned} \tag{12}$$

Note that p_x , q_x , p'_x , and q'_x in Equations (1)-(4) are denoted p_{lwr} , p_{upr} , p'_{lwr} , and q'_{upr} , respectively within Equation (12) for easy distinction between MLEs and Bayesian posterior estimates. The above statistical theory and derivations lay a good foundation for the remainder of this paper.

The proposed model is inherently vectorized to allow processing of multiple species datasets simultaneously. Model fitting was achieved using the Stan probabilistic programming language (Carpenter et al., 2017) framework for Hamiltonian Monte Carlo (HMC) via the No-U-Turn Sampler (NUTS) sampling algorithm (Hoffman and Gelman, 2014) through the **rstan** R package (version 2.32.6) (Stan Development Team, 2023) in R (version 4.4.1) (R Core Team, 2024). Four Markov chains were run for 2000 iterations each in parallel across four cores with random parameter initializations. Within each chain, a total of 1000 samples was discarded as warmup (*i.e.*, burnin) to reduce dependence on starting conditions and to ensure posterior samples are reflective of the equilibrium distribution. Further, 1000 post-warmup draws were utilized per chain during the sampling phase. Because HMC/NUTS results in dependent samples that are minimally autocorrelated, chain thinning is not required. Each of these tuning parameters reflect default Markov Chain Monte Carlo (MCMC) settings in Stan to control both bias and variance, respectively, in the resulting draws. All analyses in the present work were carried out on a 2023 Apple MacBook Pro with M2 chip and 16 GB RAM running macOS Ventura 13.2. A random seed was set to ensure

reproducibility of model results. Outputted estimates were rounded to three decimal places of precision. Posterior distributions were visualized as KDE plots using the `ggplot()` function within the `ggplot2` R package (version 3.5.1) (Wickham, 2016) with the default Gaussian kernel and optimal smoothness selection. To successfully run the Stan program, end users must have installed an appropriate compiler (such as GCC or Clang) which is compatible with their operating system, such as macOS.

Convergence was assessed both visually and quantitatively as follows: (1) through examining parameter traceplots, which depict the trajectory of accepted MCMC draws as a function of the number of iterations, (2) through monitoring the Gelman-Rubin potential scale reduction factor statistic ($\hat{R}$) (Gelman and Rubin, 1992; Vehtari et al., 2021), which measures the concordance of within-chain *versus* between-chain variance, and (3) through calculating the effective sample size (ESS) for each parameter, which quantifies the number of independent samples generated Markov chains are equivalent to. Mixing of chains was deemed sufficient when traceplots looked like “fuzzy caterpillars”, $\hat{R} < 1.01$, and effective sample sizes were reasonably large (Gelman et al., 2020). After sampling, a number of summary quantities were reported, including posterior means, posterior SDs, and posterior quantiles from which 95% CrIs could be computed to make probabilistic inferences concerning true population parameters. To validate the overall correctness of the proposed statistical model given by Equation (12), as a means of comparison, posterior predictive checks (PPCs) were also employed to generate binomial random variates in the form of counts from the posterior predictive distribution; that is $\underline{\gamma} = \{Np_x, Mq_x, Np'_x, Cq'_x\}$ to verify that the model adequately captures relevant features of the observed data. The proposed Bayesian model outlined here has a straightforward interpretation (**Table 1**).

Table 1: Interpretation of the DNA barcode gap estimators within $[a/a', b]$

Parameter	Explanation
p_x/p_{lwr}	When p_x/p_{lwr} is close to 0 (1), it suggests that the probability of intraspecific (interspecific) distances being larger (smaller) than interspecific (intraspecific) distances is low (high) on average, while the probability of interspecific (intraspecific) distances being larger (smaller) than intraspecific (interspecific) distances is high (low) on average; that is, there is (no) evidence for a DNA barcode gap.
q_x/p_{upr}	When q_x/p_{upr} is close to 0 (1), it suggests that the probability of interspecific (intraspecific) distances being larger (smaller) than intraspecific (interspecific) distances is high (low) on average, while the probability of intraspecific (interspecific) distances being larger (smaller) than interspecific (intraspecific) distances is low (high) on average; that is, there is (no) evidence for a DNA barcode gap.
p'_x/p'_{lwr}	When p'_x/p'_{lwr} is close to 0 (1), it suggests that the probability of intraspecific (combined interspecific distances for a target species and its nearest neighbour species) distances being larger (smaller) than combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) is low (high) on average, while the probability of combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) being larger (smaller) than intraspecific distances (combined interspecific distances for a target species and its nearest neighbour species) is high (low) on average; that is, there is (no) evidence for a DNA barcode gap.
q'_x/p'_{upr}	When q'_x/p'_{upr} is close to 0 (1), it suggests that the probability of combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) being larger (smaller) than intraspecific distances (combined interspecific distances for a target species and its nearest neighbour species) is high (low) on average, while the probability of intraspecific distances (combined interspecific distances for a target species and its nearest neighbour species) being larger (smaller) than combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) is low (high) on average; that is, there is (no) evidence for a DNA barcode gap.

3 Results and Discussion

The *Agabus* CYTB dataset analyzed by Phillips et al. (2024) is revisited herein for the species *A. bipustulatus* and *A. nevadensis*, since these taxa were the sole representatives for this locus, with the most and the least specimen records, respectively ($N = 701$ and $N = 2$) across all three assessed molecular markers. Briefly, using the R package **MACER** (Young et al., 2021), DNA sequences were downloaded from GenBank and BOLD and processed using a stringent workflow to obtain a 343-bp high-quality FASTA alignment representing 46 unique haplotypes. Genetic distances were calculated using uncorrected p-distances. Further, *A. bipustulatus* comprised 46 total haplotypes, whereas *A. nevadensis* possessed two haplotypes. Note, DNA barcode gap estimation is only possible for species having at least two specimen records. This dataset is a prime illustrative example highlighting the issue of inadequate taxon sampling, which arises frequently in large-scale phylogenetic and phylogeographic studies, in several respects. First, from a statistical viewpoint, sample sizes reflect extremes in reliable parameter estimation. Second, from a DNA barcoding perspective, *Agabus* currently comprises about 200 extant Linnean species according to the Global Biodiversity Information Facility (GBIF) (<https://www.gbif.org>); yet, due to the level of convenience sampling inherent in taxonomic collection efforts for this genus, adequate representation of species and genetic diversity is far from complete. Indeed, herein species coverage for CYTB is only around 1.000%.

ECDF/CECDF plots were generated using the base R function `ecdf()`, along with `ggplot()`, and are displayed in **Figure 2**. Interpretation is simple: in the case of *A. bipustulatus*, p_x , all distances are greater than $a = 0$, with 36.900% of distances equal to a . Conversely, both q_x and q'_x equal one because all interspecific distances are less than $b \approx 0.026$; for *A. nevadensis*, both metrics are approximately equal to 0.010 since only about 1% of distances fall below $b = 0$. MCMC parameter traceplots showed rapid mixing of chains to the stationary distribution (**Figure 3**). Further, all $\hat{R}$ and ESS values (not shown) were close to their recommended cutoffs of one and thousands of samples, respectively, indicating chains

are both well-mixed and have converged to the posterior distribution. Bayesian posterior estimates were reported alongside frequentist MLEs, in addition to SEs, posterior SDs, 95% CIs, and 95% CrIs (Table 2).




Figure 2: ECDF/CECDF plots of the DNA barcode gap metrics for *A. bipustulatus* and *A. nevadensis* with a/a' and b clearly indicated as dashed vertical lines. When all genetic distances are equal to zero, no step function is shown.




Figure 3: MCMC parameter traceplots applied to *A. bipustulatus* ([1]; $N = 701$) and *A. nevadensis* ([2]; $N = 2$) for CYTB across 1000 post-warmup iterations.

Table 2: Nonparametric frequentist and Bayesian estimates of distance distribution overlap/separation for the DNA barcode gap coalescent model parameters applied to *A. bipustulatus* ($N = 701$) and *A. nevadensis* ($N = 2$) for CYTB, including 95% CIs and CrIs. CrIs are based on 4000 posterior draws. All parameter estimates are reported to three decimal places of precision.

Species	Parameter	MLE (SE, 95% CI)	Bayes Est. (SD; 95% CrI)
<i>A. bipustulatus</i>	p_x/p_{lwr}	1.000 (0.000; 1.000-1.000)	1.000 (0.000; 1.000-1.000)
<i>A. bipustulatus</i>	q_x/p_{upr}	1.000 (0.000; 1.000-1.000)	1.000 (0.000; 0.999-1.000)
<i>A. bipustulatus</i>	p'_x/p'_{lwr}	1.000 (0.000; 1.000-1.000)	1.000 (0.000; 1.000-1.000)
<i>A. bipustulatus</i>	q'_x/p'_{upr}	1.000 (0.000; 1.000-1.000)	1.000 (0.000; 0.999-1.000)
<i>A. nevadensis</i>	p_x/p_{lwr}	1.000 (0.000; 1.000-1.000)	0.835 (0.144; 0.470-0.996)
<i>A. nevadensis</i>	q_x/p_{upr}	0.010 (0.002; 0.006-0.014)	0.010 (0.002; 0.007-0.014)
<i>A. nevadensis</i>	p'_x/p'_{lwr}	1.000 (0.000; 1.000-1.000)	0.834 (0.138; 0.481-0.994)
<i>A. nevadensis</i>	q'_x/p'_{upr}	0.010 (0.070; -0.128-0.148)	0.010 (0.002; 0.007-0.014)

CIs were calculated using the usual large sample $(1 - \alpha)100\%$ -level interval estimate given by $\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$, where $z_{1-\frac{\alpha}{2}} = 1.960$ is the critical value for 95% confidence (*i.e.*, the 97.5th percent quantile from the standard Normal distribution), and α is the stated significance level (here, 5%). Given a $(1 - \alpha)100\%$ CI, with repeated sampling, on average $(1 - \alpha)100\%$ of constructed intervals will contain the true parameter of interest, assuming a correctly specified statistical model; on the other hand, any given CI will either capture or exclude the true parameter with 100% certainty. This in stark contrast to a CrI, where the true parameter is contained within said interval with $(1 - \alpha)100\%$ probability, contingent on the data generating model being correct. Note, by default Stan computes equal-tailed (central) CrIs such that there is equal area situated in the left and right tails of the posterior distribution. For a 95% CrI, this corresponds to the 2.5th and 97.5th percent quantiles. However, constructed intervals are usually only valid for symmetric or nearly symmetric distributions. Given the bounded nature of the DNA barcode gap metrics, whose posterior distributions, as expected, show considerable skewness, an alternative approach to reporting CrIs, such as Highest Posterior Density (HPD) intervals (Chen and Shao, 1999) or shortest probability intervals (SPIIn) (Liu et al., 2015) is warranted. As such asymmetric intervals generally attain greater statistical efficiency (in the form of smaller Mean Squared Error (MSE) or variance) and higher coverage probabilities than more standard interval estimates,

careful in-depth comparison is left for future work.

Findings based on nonparametric MLEs and Bayesian posterior means were quite comparable with one another and show evidence of complete overlap in intraspecific, interspecific, and combined interspecific distances for *A. bipustulatus* in both the $p/q/p_{\text{LWR}}/p_{\text{UPR}}$ and $p'/q'/p'_{\text{LWR}}/p'_{\text{UPR}}$ directions since the metrics attain magnitudes very close to one, with minimal noise (**Table 2**). As a result, this likely indicates that no DNA barcode gap is present for this species. Such findings are strongly reinforced by the very tight clustering of posterior draws (**Figure 4**) and associated interval estimates owing to the large number of specimens sampled for this species.




Figure 4: Scatterplots (black solid points) and distributions (black solid lines) depicting the DNA barcode gap metrics for *A. bipustulatus* ($N = 701$) and *A. nevadensis* ($N = 2$) across CYTB based on 4000 Bayesian posterior draws. MLEs and posterior means are displayed as coloured (red/blue) solid and dashed lines for the metrics, respectively.

On the other hand, the situation for *A. nevadensis* is more nuanced, as posterior values are further spread out (**Table 2** and **Figure 4**), suggesting less overall certainty in true parameter values given the low specimen sampling coverage for this taxon. Of note, calculated SEs and

SDs are large, with the 95% CIs and 95% CrIs being quite wide in most cases, consistent with much uncertainty regarding the computed frequentist and Bayesian posterior means of the DNA barcode gap metrics. For instance, the Bayesian analysis suggests that the data are consistent with both p_{lwr} and p'_{lwr} ranging from approximately 0.500-1.000 due to the large SD of about 0.140. The posterior means associated with these estimates themselves are also far from one. The CrI for p_{upr} and p'_{upr} also spans an order of magnitude, with the lower tail estimate of 0.007 being very close to zero. Further, regarding the frequentist analysis for the same species, the 95% CI for q_x is quite wide, reflecting considerable uncertainty in its true parameter value. Similarly, that for q'_x extends to negative values at the left endpoint, due to the corresponding SE of 0.070 being too high as a result of the extremely low sample size of $n = 2$ individuals sampled (**Table 2**). Since the 95% CI for q'_x truncated at the lower endpoint includes the value of zero, the null hypothesis for the presence of a DNA barcode gap cannot be rejected. Despite this, it is worth noting that truncation is not standard statistical practice and will likely lead to an interval with less than 95% nominal coverage having high relative error. In such cases, more appropriate confidence interval methods like the Wilson score interval, the exact (Clopper-Pearson) interval, or the Agresti-Coull interval should be employed (Newcombe, 1998; Agresti and Coull, 1998). KDEs for *A. bipustulatus* are strongly left (negatively) skewed (**Figure 4**), whereas those for *A. nevadensis* exhibit more symmetry, especially for p_{upr} and p'_{upr} (**Figure 4**). These differences are likely due to the stark contrast in sample sizes for the two examined species. Nevertheless, simulated counts of overlapping specimen records from the posterior predictive distribution (**Table 3**) were found to be very close to observed counts for both species, indicating that the proposed model adequately captures underlying variation.

Table 3: Posterior predictive checks of distance distribution overlap/separation for the DNA barcode gap coalescent model parameters applied to *A. bipustulatus* ($N = 701$) and *A. nevadensis* ($N = 2$) for CYTB. CrIs are based on 4000 posterior draws. All parameter estimates are reported to three decimal places of precision.

Species	Variable	Y	n	Bayes Est. (SD; 95% CrI)
<i>A. bipustulatus</i>	y_{lwr}	491401.000	491401.000	491400.018 (1.378; 491396.000-491401.000)
<i>A. bipustulatus</i>	y_{upr}	2804.000	2804.000	2803.019 (1.433; 2799.000-2804.000)
<i>A. bipustulatus</i>	y'_{jwr}	491401.000	491401.000	491400.008 (1.412; 491396.000-491401.000)
<i>A. bipustulatus</i>	y_{upr}	2804.000	2804.000	2802.992 (1.429; 2799.000-2804.000)
<i>A. nevadensis</i>	y_{lwr}	4.000	4.000	3.355 (0.888; 1.000-4.000)
<i>A. nevadensis</i>	y_{upr}	28.000	2804.000	29.151 (7.620; 16.000-45.000)
<i>A. nevadensis</i>	y'_{jwr}	4.000	4.000	3.325 (0.884; 1.000-4.000)
<i>A. nevadensis</i>	y_{upr}	28.000	2804.000	28.942 (7.409; 15.000-45.000)

Obtained results suggest that use of the Beta(1, 1) prior may not be appropriate given a low number of collected individuals for most taxa in DNA barcoding efforts. This suggests that further consideration of more informative beta priors is worthwhile. This is clear for *A. nevadensis*, where use of a stronger prior will likely eliminate the issue regarding accurate estimation of p_{lwr} and p'_{lwr} and stabilize their uncertainty.

4 Conclusion

Herein, the accuracy of the DNA barcode gap was analyzed from a rigorous statistical lens to expedite both the curation and growth of reference sequence libraries, ensuring they are populated with high quality, statistically defensible specimen records fit for purpose to address standing questions in ecology, evolutionary biology, management, and conservation. To accomplish this, recently proposed, easy to calculate nonparametric MLEs were formally derived using ECDFs/CECDFs and applied to assess the extent of overlap/separation of distance distributions within and among two species of predatory water beetles in the genus *Agabus* sequenced at CYTB using a Bayesian binomial count model with conjugate beta priors. Findings highlight a high level of parameter uncertainty for *A. nevadensis*, whereas posterior estimates of the DNA barcode gap metrics for *A. bipustulatus* are much more certain. Based on these results, it is imperative that specimen sampling be prioritized to better reflect actual species boundaries. More generally, apart from the metrics being

employed to better highlighting the importance of within-species genetic diversity versus between-species divergence, it is expected that the approach developed herein will be of broad utility in applied fields, such as DNA-based detection of seafood fraud within global supply chains, and in the determination of species occupancy/detection probabilities at ecological sites of interest using active and passive environmental DNA (eDNA) methods such as metabarcoding.

Since the DNA barcode gap metrics often attain values very close to zero (suggesting no overlap and complete separation of distance distributions) and/or very near one (indicating no separation and complete overlap), in addition to more intermediate values, a noninformative $\text{Beta}(\frac{1}{2}, \frac{1}{2})$ prior may be more appropriate over complete ignorance imposed by a $\text{Beta}(1, 1)$ prior. The former distribution is U-shaped symmetric and places greater probability density at the extremes of the distribution due to its heavier tails, while still allowing for variability in parameter estimates within intermediate values along its domain. Note that this prior is Jeffreys' prior density (Jeffreys, 1946), which is proportional to the square root of the Fisher information $\mathcal{I}(\theta)$; that is, $\pi(\theta) \propto \theta^{-\frac{1}{2}}(1 - \theta)^{-\frac{1}{2}}$. Jeffreys' prior has several desirable statistical properties as a prior: that it is inversely proportional to the standard deviation of the binomial distribution, and most notably, that it is invariant to model reparameterization (Gelman et al., 2014). However, this prior can lead to divergent transitions, among other pathologies, imposed by complex geometry (*i.e.*, curvature) in the posterior space since many iterative stochastic MCMC sampling algorithms experience difficulties when exploring high density distribution regions. Thus, remedies to resolve them, such as lowering the step size of the HMC/NUTS sampler, should be attempted in future work, along with other approaches such as empirical Bayes estimation to approximate beta prior hyperparameters from observed data through the MLE or other methods of parameter estimation, such as the method of moments. Alternatively, hierarchical modelling could be employed to estimate separate distribution model hyperparameters for each species and/or compute distinct estimates for the directionality/comparison level of the DNA barcode gap metrics (*i.e.*, lower *vs.* upper,

560 non-prime *vs.* prime) separately within the genus under study. This would permit greater
561 flexibility through incorporating more fine-grained structure seen in the data; however, low
562 taxon sample sizes may preclude valid inferences to be reasonably ascertained due to the
563 large number additional parameters which would be introduced through the specification
564 of the hyperprior distributions. Methods outlined in Gelman et al. (2014, 2020), such as
565 dealing with non-exchangeability of observations and alternate model parameterizations like
566 the logit, may prove useful in this regard. Perhaps, the best course of action though is
567 the elicitation of priors from domain experts. Even though more work remains, it is clear
568 that both frequentist and Bayesian inference hold much promise for the future of molecular
569 biodiversity science.

Supplementary Information

None declared

Data Availability Statement

Raw data, R, and Stan code can be accessed via Dryad at:

<http://datadryad.org/stash/share/>

RZIfMixcEODe0RWP7eyXWQewSVbqEIA9UTrH3ZVKyn4.

A GitHub repository can be found at:

<https://github.com/jphill01/Bayesian-DNA-Barcode-Gap-Coalescent>.

Acknowledgements

We wish to recognise the valuable comments and discussions of Daniel (Dan) Gillis, Robert (Bob) Hanner, Robert (Rob) Young, and XXX anonymous reviewers.

We acknowledge that the University of Guelph resides on the ancestral lands of the Attawandaron people and the treaty lands and territory of the Mississaugas of the Credit. We recognize the significance of the Dish with One Spoon Covenant to this land and offer our respect to our Anishinaabe, Haudenosaunee and Métis neighbours as we strive to strengthen our relationships with them.

Funding

None declared.

Conflict of Interest

None declared.

Author Contributions

JDP wrote the manuscript, wrote R and Stan code, as well as analyzed and interpreted all model results.

References

Agresti, A. and B. A. Coull

1998. Approximate is better than ‘exact’ for interval estimation of binomial proportions.

The American Statistician, 52(2):119–126.

Ahrens, D., F. Fujisawa, H.-J. Krammer, J. Eberle, S. Fabrizi, and A. Vogler

2016. Rarity and incomplete sampling in DNA-based species delimitation. *Systematic*

Biology, 65(3):478–494.

Avise, J., J. Arnold, R. Ball, Jr., E. Bermingham, T. Lamb, J. Neigel, C. Reeb, and N. Saunders

1987. Intraspecific phylogeography: The mitochondrial DNA bridge between population genetics and systematics. *Annu. Rev. Ecol. Syst.*, 18:489–522.

Bartlett, S. and W. Davidson

1992. FINS (forensically informative nucleotide sequencing): A procedure for identifying the animal origin of biological specimens. *BioTechniques*, 12(3):408—411.

Bergsten, J., D. Bilton, T. Fujisawa, M. Elliott, M. Monaghan, M. Balke, L. Hendrich,

J. Geijer, J. Herrmann, G. Foster, I. Ribera, A. Nilsson, T. Barraclough, and A. Vogler

2012. The effect of geographical scale of sampling on DNA barcoding. *Systematic Biology*, 61(5):851–869.

Bouckaert, R., T. G. Vaughan, J. Barido-Sottani, S. Duchêne, M. Fourment,

A. Gavryushkina, J. Heled, G. Jones, D. Kühnert, N. De Maio, M. Matschiner, F. K.

Mendes, N. F. Müller, H. A. Ogilvie, L. du Plessis, A. Poppinga, A. Rambaut, D. Rasmussen,
I. Siveroni, M. A. Suchard, C.-H. Wu, D. Xie, C. Zhang, T. Stadler, and A. J. Drummond
2019. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis.
PLOS Computational Biology, 15(4):1–28.

Čandek, K. and M. Kuntner

2015. DNA barcoding gap: Reliable species identification over morphological and
geographical scales. *Molecular Ecology Resources*, 15(2):268–277.

Carpenter, B., A. Gelman, M. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker,
J. Guo, P. Li, and A. Riddell

2017. Stan: A probabilistic programming language. *Journal of Statistical Software*, 76:1.

Carstens, B. C., T. A. Pelletier, N. M. Reid, and J. D. Satler

2013. How to fail at species delimitation. *Molecular Ecology*, 22(17):4369–4383.

Chen, M.-H. and Q.-M. Shao

1999. Monte Carlo estimation of Bayesian credible and HPD intervals. *Journal of
Computational and Graphical Statistics*, 8(1):69–92.

Collins, R. A. and R. H. Cruickshank

2013. The seven deadly sins of DNA barcoding. *Molecular Ecology Resources*,
13(6):969–975.

Collins, R. A. and R. H. Cruickshank

2014. Known knowns, known unknowns, unknown unknowns and unknown knowns in dna
barcoding: A comment on dowton et al. *Systematic Biology*, 63(6):1005–1009.

Dayrat, B.

2005. Towards integrative taxonomy. *Biological Journal of the Linnean Society*,
85(3):407–417.

de Queiroz, K.

2007. Species concepts and species delimitation. *Systematic Biology*, 56(6):879–886.

Dempster, A. P., N. M. Laird, and D. B. Rubin

1977. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–22.

Dowton, M., K. Meiklejohn, S. L. Cameron, and J. Wallman

2014. A preliminary framework for DNA barcoding, incorporating the multispecies coalescent. *Systematic Biology*, 63(4):639–644.

Efron, B. and R. J. Tibshirani

1993. *An Introduction to the Bootstrap*. Chapman & Hall.

Flouri, T., X. Jiao, B. Rannala, and Z. Yang

2018. Species tree inference with BPP using genomic sequences and the multispecies coalescent. *Molecular Biology and Evolution*, 35(10):2585–2593.

Fonseca, E. M., D. J. Duckett, and B. C. Carstens

2021. P2C2M.GMYC: An R package for assessing the utility of the Generalized Mixed Yule Coalescent model. *Methods in Ecology and Evolution*, 12(3):487–493.

Fujisawa, T. and T. Barraclough

2013. Delimiting species using single-locus data and the generalized mixed yule coalescent approach: A revised method and evaluation on simulated data sets. *Systematic Biology*, 62(5):707–724.

Fujita, M. K., A. D. Leaché, F. T. Burbrink, J. A. McGuire, and C. Moritz

2012. Coalescent-based species delimitation in an integrative taxonomy. *Trends in Ecology & Evolution*, 27(9):480–488.

Gelman, A., J. Carlin, H. Stern, D. Duncan, A. Vehtari, and D. Rubin

2014. *Bayesian Data Analysis*, third edition. Chapman and Hall/CRC.

Gelman, A. and D. Rubin

1992. Inference from iterative simulation using multiple sequences. *Statistical Science*, 7(4):457–472.

Gelman, A., A. Vehtari, D. Simpson, C. Margossian, B. Carpenter, Y. Yao, L. Kennedy, J. Gabry, P.-C. Bürkner, and M. Modrák

2020. Bayesian workflow.

Hebert, P., A. Cywinska, S. Ball, and J. deWaard

2003a. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(1512):313–321.

Hebert, P., S. Ratnasingham, and J. de Waard

2003b. Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(Suppl 1):S96–S99.

Hebert, P. D., M. Y. Stoeckle, T. S. Zemlak, and C. M. Francis

2004. Identification of birds through DNA barcodes. *PLoS Biol*, 2(10):e312.

Hickerson, M. J., C. P. Meyer, and C. Moritz

2006. DNA barcoding will often fail to discover new animal species over broad parameter space. *Systematic Biology*, 55(5):729–739.

Hoffman, M. and A. Gelman

2014. The No-U-Turn Sampler: Adaptively setting path lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, 15:1593–1623.

Hubert, N. and R. Hanner

2015. DNA barcoding, species delineation and taxonomy: A historical perspective. *DNA Barcodes*, 3:44–58.

Hubert, N., J. Phillips, and R. Hanner

2024. Delimiting species with single-locus dna sequences. In *DNA Barcoding*, R. DeSalle, ed., volume 2744 of *Methods in Molecular Biology*. New York, NY: Humana.

Jackson, N. D., B. C. Carstens, A. E. Morales, and B. C. O’Meara

2017a. Species delimitation with gene flow. *Systematic Biology*, 66(5):799–812.

Jackson, N. D., A. E. Morales, B. C. Carstens, and B. C. O’Meara

2017b. PHRAPL: Phylogeographic inference using approximate likelihoods. *Systematic Biology*, 66(6):1045–1053.

Jeffreys, H.

1946. An invariant form for the prior probability in estimation problems. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 186(1007):453–461.

Jukes, T. and C. Cantor

1969. Evolution of protein molecules. In *Mammalian Protein Metabolism*, H. N. Munro, ed., Pp. 21–132. New York: Academic Press.

Kapli, P., S. Lutteropp, J. Zhang, K. Kobert, P. Pavlidis, A. Stamatakis, and T. Flouri

2017. Multi-rate poisson tree processes for single-locus species delimitation under maximum likelihood and markov chain monte carlo. *Bioinformatics*, 33(11):1630–1638.

Kimura, M.

1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16(1):111–120.

Kingman, J.

1982a. The coalescent. *Stochastic Processes and Their Applications*, 13:235–248.

710 Kingman, J.
711 1982b. On the genealogy of large populations. *Journal of Applied Probability*, 19(A):27–43.

712 Knowles, L. L.
713 2009. Statistical phylogeography. *Annual Review of Ecology, Evolution, and Systematics*,
714 40(Volume 40, 2009):593–612.

715 Knowles, L. L. and W. P. Maddison
716 2002. Statistical phylogeography. *Molecular Ecology*, 11(12):2623–2635.

717 Kullback, S. and R. Leibler
718 1951. On information and sufficiency. *Annals of Mathematical Statistics*, 22(1):79–86.

719 Leaché, A., M. Fujita, V. Minin, and R. Bouckaert
720 2014. Species delimitation using genome-wide snp data. *Systematic Biology*, 63(4):534–542.

721 Liu, Y., A. Gelman, and T. Zheng
722 2015. Simulation-efficient shortest probability intervals. *Statistical Computing*, 25:809–819.

723 Mather, N., S. Traves, and S. Ho
724 2019. A practical introduction to sequentially Markovian coalescent methods for estimating
725 demographic history from genomic data. *Ecology and Evolution*, 10(1):579–589.

726 Mayr, E.
727 1942. *Systematics and the Origin of Species from the Viewpoint of a Zoologist*. New York:
728 Columbia University Press.

729 Meier, R., G. Zhang, and F. Ali
730 2008. The use of mean instead of smallest interspecific distances exaggerates the size of
731 the “barcoding gap” and leads to misidentification. *Systematic Biology*, 57(5):809–813.

732 Meyer, C. and G. Paulay
733 2005. DNA barcoding: Error rates based on comprehensive sampling. *PLOS Biology*,
734 3(12):e422.

735 Monaghan, M., R. Wild, M. Elliot, T. Fujisawa, M. Balke, D. Inward, D. Lees,
736 R. Ranaivosolo, P. Eggleton, T. Barraclough, and A. Vogler
737 2009. Accelerated species inventory on madagascar using coalescent-based models of species
738 delineation. *Systematic Biology*, 58:298–311.

739 Mutanen, M., S. M. Kivelä, R. A. Vos, C. Doorenweerd, S. Ratnasingham, A. Hausmann,
740 P. Huemer, V. Dincă, E. J. van Nieukerken, C. Lopez-Vaamonde, R. Vila, L. Aarvik,
741 T. Decaëns, K. A. Efetov, P. D. N. Hebert, A. Johnsen, O. Karsholt, M. Pentinsaari,
742 R. Rougerie, A. Segerer, G. Tarmann, R. Zahren, and H. C. J. Godfray
743 2016. Species-level para- and polyphyly in DNA barcode gene trees: Strong operational
744 bias in european lepidoptera. *Systematic Biology*, 65(6):1024–1040.

745 Newcombe, R. G.
746 1998. Two-sided confidence intervals for the single proportion: comparison of seven
747 methods. *Statistics in Medicine*, 17(8):857–872.

748 Pante, E., N. Puillandre, A. Viricel, S. Arnaud-Haond, D. Aurelle, M. Castelin, A. Chenuil,
749 C. Destombe, D. Forcioli, M. Valero, F. Viard, and S. Samadi
750 2015. Species are hypotheses: Avoid connectivity assessments based on pillars of sand.
751 *Molecular Ecology*, 24(3):525–544.

752 Paradis, E., J. Claude, and K. Strimmer
753 2004. Ape: Analyses of phylogenetics and evolution in r language. *Bioinformatics*,
754 20(2):289–290.

755 Pentinsaari, M., H. Salmela, M. Mutanen, and T. Roslin
756 2016. Molecular evolution of a widely-adopted taxonomic marker (COI) across the animal
757 tree of life. *Scientific Reports*, 6:35275.

758 Phillips, J., T. Athey, P. McNicholas, and R. Hanner
759 2023. VLF: An r package for the analysis of very low frequency variants in DNA sequences.
760 *Biodiversity Data Journal*, 11:e96480.

761 Phillips, J., D. Gillis, and R. Hanner
762 2019. Incomplete estimates of genetic diversity within species: Implications for DNA
763 barcoding. *Ecology and Evolution*, 9(5):2996–3010.

764 Phillips, J., D. Gillis, and R. Hanner
765 2020. HACSIm: An R package to estimate intraspecific sample sizes for genetic diversity
766 assessment using haplotype accumulation curves. *PeerJ Computer Science*.

767 Phillips, J., D. Gillis, and R. Hanner
768 2022. Lack of statistical rigor in DNA barcoding likely invalidates the presence of a true
769 species’ barcode gap. *Frontiers in Ecology and Evolution*, 10:859099.

770 Phillips, J., C. Griswold, R. Young, N. Hubert, and H. Hanner
771 2024. *A Measure of the DNA Barcode Gap for Applied and Basic Research*, Pp. 375–390.
772 New York, NY: Springer US.

773 Phillips, J., R. Gwiazdowski, D. Ashlock, and R. Hanner
774 2015. An exploration of sufficient sampling effort to describe intraspecific DNA barcode
775 haplotype diversity: examples from the ray-finned fishes (Chordata: Actinopterygii). *DNA*
776 *Barcodes*, 3:66–73.

777 Pons, J., T. G. Barraclough, J. Gomez-Zurita, A. Cardoso, D. P. Duran, S. Hazell,

778 S. Kamoun, W. D. Sumlin, and A. P. Vogler

779 2006. Sequence-based species delimitation for the DNA taxonomy of undescribed insects.

780 *Systematic Biology*, 55(4):595–609.

781 Puillandre, N., S. Brouillet, and G. Achaz

782 2021. ASAP: Assemble Species by Automatic Partitioning. *Molecular Ecology Resources*,

783 21(2):609–620.

784 Puillandre, N., A. Lambert, S. Brouillet, and G. Achaz

785 2011. ABGD, Automatic Barcode Gap Discovery for primary species delimitation.

786 *Molecular Ecology*, 21(8):1864–1877.

787 R Core Team

788 2024. *R: A Language and Environment for Statistical Computing*. R Foundation for

789 Statistical Computing, Vienna, Austria.

790 Rannala, B.

791 2015. The art and science of species delimitation. *Current Zoology*, 61(5):846–853.

792 Rannala, B. and Z. Yang

793 2003. Bayes estimation of species divergence times and ancestral population sizes using

794 DNA sequences from multiple loci. *Genetics*, 164:1645–1656.

795 Rannala, B. and Z. Yang

796 2017. Efficient Bayesian species tree inference under the multispecies coalescent. *Systematic*

797 *Biology*, 66(5):823–842.

798 Ratnasingham, S. and P. Hebert

799 2007. BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Molecular*

800 *Ecology Notes*, 7(3):355–364.

Ratnasingham, S. and P. D. N. Hebert

2013. A dna-based registry for all animal species: The barcode index number (bin) system. *PLoS One*, 8(7):e66213.

Stan Development Team

2023. RStan: The R interface to Stan. R package version 2.32.6.

Stoeckle, M. and D. Thaler

2014. DNA barcoding works in practice but not in (neutral) theory. *PLoS One*, 9(7):e100755.

Talavera, G., V. Dincă, and R. Vila

2013. Factors affecting species delimitations with the GMYC model: insights from a butterfly survey. *Methods in Ecology and Evolution*, 4(12):1101–1110.

Vehtari, A., A. Gelman, D. Simpson, B. Carpenter, and P.-C. Bürkner

2021. Rank-normalization, folding, and localization: An improved $\hat{R}$ for assessing convergence of MCMC (with discussion). *Bayesian Analysis*, 16(2):667–718.

Wells, T., T. Carruthers, P. Muñoz-Rodríguez, A. Sumadijaya, J. R. I. Wood, and R. W. Scotland

2022. Species as a heuristic: Reconciling theory and practice. *Systematic Biology*, 71(5):1233–1243.

Wickham, H.

2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wiemers, M. and K. Fiedler

2007. Does the DNA barcoding gap exist? - a case study in blue butterflies (Lepidoptera: Lycaenidae). *Frontiers in Zoology*, 4(8).

824 Yang, Z. and B. Rannala
825 2010. Bayesian species delimitation using multilocus sequence data. *Proceedings of the*
826 *National Academy of Sciences*, 107:9264–9269.

827 Yang, Z. and B. Rannala
828 2014. Unguided species delimitation using dna sequence data from multiple loci. *Molecular*
829 *Biology and Evolution*, 31(12):3125–3135.

830 Yang, Z. and B. Rannala
831 2017. Bayesian species identification under the multispecies coalescent provides significant
832 improvements to DNA barcoding analyses. *Molecular Ecology*, 26:3028–3036.

833 Young, R., R. Gill, D. Gillis, and R. Hanner
834 2021. Molecular Acquisition, Cleaning and Evaluation in R (MACER) - A tool to assemble
835 molecular marker datasets from BOLD and GenBank. *Biodiversity Data Journal*, 9:e71378.

836 Zhang, J., P. Kapli, P. Pavlidis, and A. Stamatakis
837 2013. A general species delimitation method with applications to phylogenetic placements.
838 *Bioinformatics*, 29(22):2869–2876.