

1 **A Bayesian Model of the DNA Barcode Gap**

2 Jarrett D. Phillips<sup>1,2\*</sup> (ORCID: 0000-0001-8390-386X), ... (others?)

3 <sup>1</sup>*School of Computer Science, University of Guelph, Guelph, ON., Canada, N1G2W1*

4 <sup>2</sup>*Department of Integrative Biology, University of Guelph, Guelph, ON., Canada, N1G2W1*

5 **\*Corresponding Author:** Jarrett D. Phillips<sup>1</sup>

6 **Email Address:** jphill01@uoguelph.ca

7 **Running Title:** Bayesian inference for DNA barcode gap estimation

## Abstract

A simple statistical model of the DNA barcode gap is outlined. Specifically, accuracy of recently introduced nonparametric metrics, inspired by coalescent theory, to characterize the extent of proportional overlap/separation in maximum and minimum pairwise genetic distances within and among species, respectively, is explored in both frequentist and Bayesian contexts. The empirical cumulative distribution function (ECDF) is utilized to estimate probabilities associated with positively skewed extreme tail distribution quantiles bounded on the closed unit interval  $[0, 1]$  based on a straightforward binomial distance overlap count. Using R and Stan, the proposed maximum likelihood estimators and Bayesian model are demonstrated on cytochrome *b* (CYTB) gene sequences from two *Agabus* diving beetle species exhibiting limits in the extent of representative taxonomic sampling. Large-sample theory and MCMC simulations show much uncertainty in parameter estimates, particularly when specimen sample sizes for target species are small. Findings highlight the promise of the Bayesian approach using a conjugate beta prior for reliable posterior uncertainty estimation when available data are sparse. Obtained results can shed light on foundational and applied research questions concerning DNA-based specimen identification and species delineation for studies in evolutionary biology and ecology, as well as biodiversity conservation and management, of wide-ranging taxa.

**Keywords:** Bayesian/frequentist inference, DNA barcoding, intraspecific genetic distance, interspecific genetic distance, specimen identification, species discovery

## 1 Introduction

The routine use of DNA sequences to support broad evolutionary hypotheses and questions concerning demographic processes, like gene flow and speciation, is now ubiquitous. Since the late 1980s, mitochondrial DNA (mtDNA) has been employed to produce a distinctive and measurable pattern of genetic polymorphism in diverse and

spatially-distributed taxonomic lineages such as birds, fishes, insects, and arachnids, among other extensively studied groups (Avice et al., 1987). The application of genomic data to applied fields like biodiversity forensics, conservation, and management for the molecular identification of unknown specimen samples came later (*e.g.*, Forensically Informative Nucleotide Sequencing (FINS); Bartlett and Davidson (1992)). Since its inception over 20 years ago, DNA barcoding (Hebert et al., 2003a,b) built significantly on earlier work and has emerged as a robust method of specimen identification and species discovery across myriad multicellular eukaryotes which have been sequenced at easily obtained short, standardized gene regions like the cytochrome *c* oxidase subunit I (5'-COI) mitochondrial locus for animals.

The success of the single-locus approach, particularly for regulatory and forensic applications, depends crucially on two important factors: (1) the availability of high-quality specimen records found in public reference sequence databases such as the Barcode of Life Data Systems (BOLD; <http://www.barcodinglife.org>) (Ratnasingham and Hebert, 2007) and GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>), and (2) the establishment of a DNA barcode gap — the notion that the maximum genetic distance observed within species is much smaller than the minimum degree of marker variation found among species (Meyer and Paulay, 2005; Meier et al., 2008). Early work has demonstrated that the presence of a DNA barcode gap hinges strongly on extant levels of species haplotype diversity gauged from comprehensive specimen sampling at wide geographic and ecological scales (Bergsten et al., 2012; Čandek and Kuntner, 2015). Despite this, many taxonomic groups lack adequate separation in their pairwise intraspecific and interspecific genetic distances due to varying rates of evolution in both genes and taxa (Pentinsaari et al., 2016). Furthermore, it has been well demonstrated that the presence of a DNA barcode gap becomes less certain with increasing spatial scale of sampling since interspecific distances increase, while intraspecific distances shrink as more closely-related species are sampled (Phillips et al., 2022). This can pose problems in cases of rare species or monotypic taxa, for instance (Ahrens et al., 2016). As a result, rapid matching of unknown samples to expertly-validated references can

be compromised, leading to cases of false positives (taxon oversplitting) and false negatives (excessive lumping of taxa) due to incomplete lineage sorting, interspecies hybridization, genome introgression, species synonymy, cryptic species diversity, and misidentifications (Hubert and Hanner, 2015; Phillips et al., 2022).

Recent work has argued that DNA barcoding, in its current form, is lacking in statistical rigor, as most studies rely strongly on heuristic distance-based measures to infer taxonomic identity and delineate species boundaries. Of these studies, few report measures of uncertainty, such as standard errors (SEs) and confidence intervals (CIs), around estimates of intraspecific and interspecific variation (*e.g.*, Wiemers and Fiedler (2007)), calling into question the existence of a true species' DNA barcode gap (Čandek and Kuntner, 2015; Phillips et al., 2022). To support this idea, novel nonparametric locus- and species-specific metrics based on the multispecies coalescent (MSC) were recently outlined by Phillips et al. (2024). The basic coalescent (Kingman, 1982a,b) encompasses a backwards continuous-time stochastic Markov process of allelic sampling among lineages under the infinite sites model of mutation within a large neutrally-evolving, panmictic species population of constant size from the present into the past towards the most recent common ancestor (MRCA). Although the coalescent sampling process plays a foundational role in evolutionary and population genetics theory due to its inherent simplicity and flexibility, it has been both under-utilized and under-appreciated as a key player within DNA barcoding (Collins and Cruickshank, 2014; Stoeckle and Thaler, 2014; Hubert and Hanner, 2015; Phillips et al., 2022). Previously proposed MSC algorithmic approaches (of which there are too many to exhaustively list here), generally assume a strict molecular clock and a simplified model of DNA sequence evolution across closely-related taxa, from which an estimated species phylogeny may be constructed (*e.g.*, with or without use of a guide tree) (*e.g.*, Rannala and Yang (2003, 2017); Yang and Rannala (2010, 2014, 2017)). Even taxon delimitation methods designed for single loci, such as the Generalized Mixed Yule Coalescent (GMYC) (Pons et al., 2006; Fujisawa and Barraclough, 2013) and Poisson Tree Processes (PTP) (Zhang et al., 2013;

88 Kapli et al., 2017), which have seen much use in DNA barcoding initiatives, have their flaws.  
89 Optimal performance of these methods relies heavily on careful parameter selection (*e.g.*,  
90 prior specification of branching rates in ultrametric trees in the case of GMYC) (Hubert  
91 et al., 2024) within third party software like BEAST (Bouckaert et al., 2019), among other  
92 concerns (Fonseca et al., 2021). In contrast, Phillips et al.’s (2024) approach is tree-free and  
93 does not require judicious parameter setting. Therefore, the method is extremely efficient  
94 and fast to run.

95 The DNA barcode gap statistics have been shown to hold strong promise for reliable DNA  
96 barcode gap assessment when applied to predatory *Agabus* (Coleoptera: Dytiscidae) diving  
97 beetles (Phillips et al., 2024). Despite their ease of sampling and well-established taxonomy,  
98 this group possesses few morphologically-distinct taxonomic characters that readily facilitate  
99 their assignment to the species level (Bergsten et al., 2012). Accurate taxon discrimination  
100 within *Agabus* is paramount since several species are currently listed as endangered. Further,  
101 the proposed metrics indicate that sister species pairs from this taxon are often difficult  
102 to distinguish on the basis of their DNA barcode sequences (Phillips et al., 2024), which  
103 may be caused by *Agabus*’s increased degree of non-monophyly at wider geospatial scales,  
104 hybridization/introgression, or synonymy (Bergsten et al., 2012). Using sequence data from  
105 three mitochondrial cytochrome markers (5’-COI, 3’-COI, and cytochrome *b* (CYTB))  
106 obtained from BOLD and GenBank, results highlight that DNA barcoding has been a  
107 one-sided argument. Phillips et al.’s (2024) findings point to the need to balance both  
108 the sufficient collection of specimens, as well as the extensive sampling of species. Not  
109 surprisingly, DNA barcode libraries are biased toward the latter effort (Phillips et al., 2022).

110 The estimators from Phillips et al. (2024) represent a clear improvement over simple,  
111 yet arbitrary, distance heuristics such as the 2% rule noted by Hebert et al. (2003a) and  
112 the 10 $\times$  rule (Hebert et al., 2004) that form the basis of single-locus species delimitation  
113 tools like Automatic Barcode Gap Discovery (ABGD) (Puillandre et al., 2011), Assemble  
114 Species by Automatic Partitioning (ASAP) (Puillandre et al., 2021), and the Barcode Index

115 Number (BIN) framework (Ratnasingham and Hebert, 2013). The 2% rule asserts that  
 116 DNA barcodes differing by at least 2% at sequenced genomic regions should be expected  
 117 to originate from different biological species, whereas the 10 $\times$  rule suggests that sequences  
 118 displaying 10 times more genetic variation among species than within taxa is evidence for a  
 119 distinct evolutionary origin. However, the lack of adoption of an explicit, universally agreed  
 120 upon, species concept that readily governs lineage formation and proliferation necessary to  
 121 establish rigorous taxon definitions for successful delimitation of hypothesized and heuristic  
 122 evolutionary units using these well-known criteria, in conjunction with secondary lines of  
 123 evidence (*e.g.*, morphology, ecology, geography, and behaviour) promised by an integrative  
 124 framework, is missing (de Queiroz, 2007; Rannala, 2015; Pante et al., 2015; Wells et al., 2022).  
 125 In addition, the reliance on visualization approaches, such as frequency histograms, dotplots,  
 126 and quadrant plots, to expose DNA barcoding’s limitations, has also been criticized  
 127 (Collins and Cruickshank, 2013; Phillips et al., 2022). Up until the work of Phillips et al.  
 128 (2024), the majority of studies (*e.g.*, Young et al. (2021)) have treated the DNA barcode  
 129 gap as a binary response. However, given poor sampling depth for most taxa, a Yes/No  
 130 dichotomy is inherently flawed because it can falsely imply a DNA barcode gap is present  
 131 for a taxon of interest when in fact no such separation in genetic distances exists. On the  
 132 other hand, in the case of sequence overlap, while a single specimen record is sufficient to  
 133 nullify the existence of a DNA barcode gap in light of available data, it is not a necessary  
 134 condition, since said taxon record could represent a species misidentification or other error.  
 135 The proposed statistics quantify the extent of asymmetric directionality of proportional  
 136 distance distribution overlap/separation for species within well-sampled taxonomic genera  
 137 based on a straightforward distance count, in a similar vein to established measures of  
 138 statistical similarity such as the Kullback-Leibler (KL) divergence (Kullback and Leibler,  
 139 1951) and other related statistics. The metrics can be employed in a variety of ways,  
 140 including to validate performance of marker genes for specimen identification to the species  
 141 level (using nonparametric bootstrapping, as in Phillips et al. (2024)), as well as to assess

whether computed values are consistent with population genetic-level parameters like effective  
 population size ( $N_e$ ), mutation rates ( $\mu$ ) and divergence times ( $\tau$ ) for species under study in a  
 statistical phylogeographic setting (Knowles and Maddison, 2002; Mather et al., 2019). Early  
 on, DNA barcoding was presumed to only work for reciprocally monophyletic groups and  
 thus concerned itself with terminal branches of generated phylogenies rather than more basal  
 lineages occurring deeper in hypothesized species trees (Mutanen et al., 2016). Furthermore,  
 the occurrence of short branches and large  $N_e$  within resolved phylogenies increases the  
 probability of deep coalescence, clouding species delimitations, which often fail or are  
 uncertain in broad parameter space (Carstens et al., 2013; Hickerson et al., 2006; Rannala,  
 2015). As DNA barcoding is a single-locus approach, it is problematic for evolutionarily young  
 taxa, wherein incomplete lineage sorting within gene genealogies is a common phenomenon  
 due to the ongoing stochastic dynamic of mutation generating population variation, and  
 genetic drift driving gene variants to fixation (Rannala, 2015). The most promising way  
 forward in this regard seems to be through the use of software such as BPP (Bayesian  
 Phylogenetics and Phylogeography), which permits efficient full Bayesian simulations under  
 various MSC models (*e.g.*, MSC-I (MSC with introgression) or MSC-M (MSC with  
 migration), among others) using MCMC for tree parameter estimation (via the A00 option,  
 for instance) (Flouri et al., 2018), or PHRAPL (Phylogeographic Inference using Approximate  
 Likelihoods) (Jackson et al., 2017a,b), which employs tractable phylogenetic likelihood  
 calculations via the genealogical divergence index (gdi).

While introduction of the metrics is a step in the right direction, what appears to be  
 missing is a rigorous statistical treatment of the DNA barcode gap. This includes an unbiased  
 way to compute the statistical accuracy of Phillips et al.’s (2024) estimators arising through  
 problems inherent in frequentist maximum likelihood estimation for probability distributions  
 having bounded positive support on the closed unit interval  $[0, 1]$ . To this end, here, a  
 Bayesian model of the DNA barcode gap coalescent is introduced to rectify such issues. The  
 model allows accurate estimation of posterior means, posterior standard deviations (SDs),

posterior quantiles, and credible intervals (CrIs) for the metrics given datasets of intraspecific and interspecific distances for species of interest.

## 2 Methods

### 2.1 DNA Barcode Gap Metrics

The novel nonparametric maximum likelihood estimators (MLEs) of proportional overlap/separation between intraspecific and interspecific distance distributions for a given species ( $x$ ) to aid assessment of the DNA barcode gap are as follows:

$$p_x = \frac{\#\{d_{ij} \geq a\}}{\#\{d_{ij}\}} \quad (1)$$


$$q_x = \frac{\#\{d_{XY} \leq b\}}{\#\{d_{XY}\}} \quad (2)$$

$$p'_x = \frac{\#\{d_{ij} \geq a'\}}{\#\{d_{ij}\}} \quad (3)$$

$$q'_x = \frac{\#\{d'_{XY} \leq b\}}{\#\{d'_{XY}\}} \quad (4)$$

where  $d_{ij}$  are distances within species,  $d_{XY}$  are distances among species for an entire genus of concern, and  $d'_{XY}$  are combined interspecific distances for a target species and its closest neighbouring species. The notation  $\#$  reflects a count. Quantities  $a$ ,  $a'$ , and  $b$  correspond to  $\min(d_{XY})$ ,  $\min(d'_{XY})$ , and  $\max(d_{ij})$ , the minimum interspecific distance, the minimum combined interspecific distance, and the maximum intraspecific distance, respectively (**Figure 1**).





**Figure 1:** Modified depiction from Meyer and Paulay (2005) and Phillips et al. (2024) of the overlap/separation of intraspecific and interspecific distances ( $\delta$ ) for calculation of the DNA barcode gap metrics ( $p_x$  and  $q_x$ ) for a hypothetical species  $x$ . The minimum interspecific distance is denoted by  $a$  and the maximum intraspecific distance is indicated by  $b$ . The quantity  $f(\delta)$  is akin to a kernel density estimate of the probability density function of distances. A similar visualization can be displayed for  $p'_x$  and  $q'_x$  within the interval  $[a', b]$ .

Hence, Equations (1)-(4) are simply empirical partial means of distances falling at and below, or at and exceeding, given distribution thresholds. Notice further that  $a/a'$ , and  $b$  are also the first and  $n$ th order statistics,  $X_{(1)}$  and  $X_{(n)}$ , respectively, with  $a/a' < b$ , which have been pointed out by Phillips et al. (2022) as important for developing a mathematical theory to test the existence of the DNA barcode gap. Equations (1)-(4) can also be expressed in terms of empirical cumulative distribution functions (ECDFs) (see next section). Distances form a continuous distribution and are easily computed from a model of DNA sequence evolution, such as uncorrected or corrected p-distances (Jukes and Cantor, 1969; Kimura, 1980) using, for example, the `dist.dna()` function available in the `ape` R package (Paradis et al., 2004). However, computed values are *not* independent and identically distributed (IID) because estimated standard errors (SEs) will depend on both the number of species sampled with the genus under study, as well as the number of specimens sampled within a target species. In Phillips et al. (2024), To tease this out, Phillips et al. (2024) suggests plotting estimator values against their estimated SEs, along with a simple random downsampling scheme. In the case of two species comprising a focal genus, one well sampled and the other poorly sampled,

values of the metrics close to zero for the sufficiently sampled species will likely possess larger SEs following downsizing to match the number of poorly sampled specimens (Phillips et al., 2024). The approach of Phillips et al. (2024) differs markedly from the traditional definition of the DNA barcoding gap laid out by Meyer and Paulay (2005) and Meier et al. (2008) in that the proposed metrics incorporate interspecific distances which *include* the target species of interest. Furthermore, if a focal species is found to have multiple nearest neighbours, then the species possessing the smallest average distance is used (Phillips et al., 2024). These schemes more accurately account for species' coalescence processes inferred from contemporaneous samples of DNA sequences leading to instances of barcode sequence sharing, such as interspecific hybridization/introgression events (Phillips et al., 2024). Within Equations (3) and (4), the degree of distance distribution overlap/separation between a target taxon and its nearest neighbouring species, gauged from magnitudes of  $p'_x$  and  $q'_x$ , is directly proportional to the amount of time in which the two lineages diverged from the MRCA (Phillips et al., 2024). Thus, the quantities can be used as a criterion to assess the failure of DNA barcoding in recently radiated taxonomic groups, among other plausible biological explanations. Note, distances are constrained to the interval  $[0, 1]$ , whereas the metrics are defined only on the interval  $[a/a', b]$ . Values of the estimators obtained from Equations (1)-(4) close to or equal to zero give evidence for separation between intraspecific and interspecific distance distributions; that is, values suggest the presence of a DNA barcode gap for a target species. Conversely, values near or equal to one give evidence for distribution overlap; that is, values likely indicate the absence of a DNA barcode gap.

## 2.2 The Model

Before delving into the derivation of the proposed DNA barcode gap metrics, review of some fundamental statistical theory is necessary.

For a given random variable  $X$ , its cumulative distribution function (CDF) is defined by

$$F_X(t) = \mathbb{P}(X \leq t) = 1 - \mathbb{P}(X > t). \quad (5)$$

222 Rearranging Equation (5) gives

$$\mathbb{P}(X > t) = 1 - F_X(t) = 1 - \mathbb{P}(X \leq t), \quad (6)$$

223 from which it follows that

$$\mathbb{P}(X \geq t) = 1 - F_X(t) + \mathbb{P}(X = t). \quad (7)$$

224 Equations (1)-(4) can thus be expressed in terms of ECDFs as follows, since the true  
 225 underlying CDFs,  $F(\cdot)$ , are unknown *a priori*, and therefore must be estimated using available  
 226 data:

$$\begin{aligned}
p_x &= \mathbb{P}(d_{ij} \geq a) \\
&= 1 - \hat{F}_{d_{ij}}(a) + \mathbb{P}(d_{ij} = a) \\
&= \hat{F}_{d_{ij}}(b) - \hat{F}_{d_{ij}}(a) + \mathbb{P}(d_{ij} = a)
\end{aligned} \tag{8}$$

$$\begin{aligned}
q_x &= \mathbb{P}(d_{XY} \leq b) \\
&= \hat{F}_{d_{XY}}(b)
\end{aligned} \tag{9}$$

$$\begin{aligned}
p'_x &= \mathbb{P}(d_{ij} \geq a') \\
&= 1 - \hat{F}_{d_{ij}}(a') + \mathbb{P}(d_{ij} = a') \\
&= \hat{F}_{d_{ij}}(b) - \hat{F}_{d_{ij}}(a') + \mathbb{P}(d_{ij} = a')
\end{aligned} \tag{10}$$

$$\begin{aligned}
q'_x &= \mathbb{P}(d'_{XY} \leq b) \\
&= \hat{F}_{d'_{XY}}(b)
\end{aligned} \tag{11}$$

227 From this, it can be seen that  $\hat{F}_{d_{ij}}(b) = 1$  in Equations (8) and (10). Given  $n$   
 228 increasing-ordered data points, the (discrete) ECDF,  $\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{[x_i \leq t]}$ , comprises a step  
 229 function having jump discontinuities of size  $\frac{1}{n}$  at each sample observation ( $x_i$ ), excluding ties  
 230 (or steps of weight  $\frac{i}{n}$  with duplicate observations), where  $\mathbf{1}(x)$  is the indicator function. Note,  
 231  $\mathbb{P}(X = t) \neq 0$ . Equations (8)-(11) clearly demonstrate the asymmetric directionality of the  
 232 proposed metrics. Plots of the ECDFs of intraspecific, interspecific, and combined distances,  
 233 superimposed with corresponding values of  $a/a'$  and  $b$ , provide a useful way to visualize the  
 234 DNA barcode gap. Furthermore, calculation of the DNA barcode gap estimators is convenient  
 235 as they implicitly account for total distribution area (including overlap).

236 A major criticism of large sample (frequentist) theory is that it relies on asymptotic  
 237 properties of the MLE (whose population parameter is assumed to be a fixed but unknown  
 238 quantity), such as estimator normality and consistency as the sample size approaches infinity.  
 239 This problem is especially pronounced in the case of binomial proportions (Newcombe, 1998).

The estimated Wald SE of the sample proportion, is given by  $\widehat{SE}[\hat{p}] = \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$ , where  $\hat{p} = \frac{Y}{n}$  is the MLE,  $Y$  is the total number of successes ( $Y = \sum_{i=1}^n y_i$ ), and  $n$  is the total number of trials (*i.e.*, sample size). However, the above formula for the standard error is problematic for several reasons. First, it is a Normal approximation which makes use of the central limit theorem (CLT); thus, large sample sizes are required for reliable estimation. When few observations are available, SEs will be large and inaccurate, leading to low statistical power to detect a true DNA barcode gap when one actually exists. Further, resulting interval estimates could span values less than zero or greater than one, or have zero width, which is practically meaningless. Second, when proportions are exactly equal to zero or one, resulting SEs will be exactly zero, rendering  $\widehat{SE}[\hat{p}]$  given above completely useless. In the context of the proposed DNA barcode gap metrics, values obtained at the boundaries of their support are often encountered. Therefore, reliable calculation of SEs is not feasible. Given the importance of sufficient sampling of species genetic diversity for DNA barcoding initiatives, a different statistical estimation approach is necessary.

Bayesian inference offers a natural path forward in this regard since it allows for straightforward specification of prior beliefs concerning unknown model parameters and permits the seamless propagation of uncertainty, when data are lacking and sample sizes are small, through integration with the likelihood function associated with true generating processes. The posterior distribution ( $\pi(\theta|Y)$ ) is given by Bayes' theorem up to a proportionality  $\pi(\theta|Y) \propto \pi(Y|\theta)\pi(\theta)$ , where  $\theta$  are unobserved parameters,  $Y$  are known data,  $\pi(Y|\theta)$  is the likelihood, and  $\pi(\theta)$  is the prior. As a consequence, because parameters are treated as random variables, Bayesian models are much more flexible and generally more easily interpretable compared to frequentist approaches. Under the Bayesian paradigm, entire posterior distributions, along with their summaries (*e.g.*, CrIs) are outputted, rather than just long run behaviour reflected in sampling distributions, p-values, and CIs as in the frequentist case, thus allowing direct probability statements to be made.

Essentially, from a statistical perspective, the goal herein is to nonparametrically estimate

267 probabilities corresponding to extreme tail quantiles for positive highly skewed distributions  
 268 on the unit interval (or any closed subinterval thereof). Here, it is sought to numerically  
 269 approximate the extent of proportional overlap/separation of intraspecific and interspecific  
 270 distance distributions within the subinterval  $[a/a', b]$ . This is a challenging computational  
 271 problem within the current study as detailed in subsequent sections. The usual approach  
 272 employs kernel density estimation (KDE), along with numerical or Monte Carlo integration  
 273 of complicated probability distribution functions (PDFs), and invocation of extreme value  
 274 theory (EVT); however, this requires careful selection of the bandwidth parameter, among  
 275 other considerations. This becomes problematic when fitting finite mixture models where  
 276 nonidentifiability is rampant. For DNA barcode gap estimation, this would correspond  
 277 to a two-component mixture (one for intraspecific distance comparisons, and the other for  
 278 interspecific comparisons), with one or more curve intersection points between components,  
 279 and the presence of zero distance inflation. This makes parameter estimation difficult using  
 280 methods like the Expectation-Maximization (EM) algorithm (Dempster et al., 1977) as  
 281 the algorithm may become stuck in suboptimal regions of the parameter search space and  
 282 prematurely converge to local optima. This seems wholly plausible, especially given that  
 283 the genus-level distribution of genetic distances is a mixture of closely-related (*i.e.*, nearest  
 284 neighbour) and more distant species. Here, for simplicity, an alternate route is taken to  
 285 avoid these obstacles. Counts,  $y$ , of overlapping distances (as expressed in the numerator  
 286 of Equations (1)-(4)) are treated as binomially distributed with expectation  $\mathbb{E}[Y|\theta] = k\theta$ ,  
 287 and variance  $\mathbb{V}[Y|\theta] = k\theta(1 - \theta)$ , where  $k = \{N, C\}$  are total count vectors of intraspecific  
 288 and combined interspecific distances, respectively, for a target species along with its nearest  
 289 neighbour species, and  $k = M$  is a total count vector for all interspecific species comparisons.  
 290 This follows from the fact that the ECDF is binomially distributed. The quantity thus being  
 291 estimated is the parameter vector  $\underline{\theta} = \{p_x, q_x, p'_x, q'_x\}$ .

292 The metrics encompassing  $\underline{\theta}$  are presumed to follow a Beta( $\alpha, \beta$ ) distribution, with real  
 293 shape parameters  $\alpha$  and  $\beta$ , which is a natural choice of prior on probabilities. The beta

294 distribution has a prior mean of  $\mathbb{E}[\theta|\alpha, \beta] = \frac{\alpha}{\alpha+\beta}$  and a prior variance equal to  
 295  $\mathbb{V}[\theta|\alpha, \beta] = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$ . In the case where  $\alpha = \beta$ , all generated  $\text{Beta}(\alpha, \beta)$  distributions  
 296 will possess the same prior expectation, whereas the prior variance will shrink as both  $\alpha$  and  
 297  $\beta$  increase. Such a scheme is quite convenient since the beta distribution is conjugate to the  
 298 binomial distribution. Thus, the posterior distribution is also beta distributed, specifically,  
 299  $\text{Beta}(\alpha + Y, \beta + n - Y)$ , having expectation  $\mathbb{E}[\theta|Y, \alpha, \beta] = \frac{\alpha+Y}{\alpha+\beta+n}$  and variance  
 300  $\mathbb{V}[\theta|Y, \alpha, \beta] = \frac{(\alpha+Y)(\beta+n-Y)}{(\alpha+\beta+n)^2(\alpha+\beta+n+1)}$ . In the context of DNA barcoding, it is important that the  
 301 DNA barcode gap metrics effectively differentiate between extremes of no overlap/complete  
 302 separation and complete overlap/no separation, corresponding to values of the metrics equal  
 303 to 0 and 1 (equivalent to total distance counts of 0 and  $n$ ), respectively. These extremes yield  
 304 a posterior expectation of  $\mathbb{E}[\theta|Y = 0, \alpha, \beta] = \frac{\alpha}{\alpha+\beta+n}$  and a posterior variance of  
 305  $\mathbb{V}[\theta|Y = 0, \alpha, \beta] = \frac{\alpha(\beta+n)}{(\alpha+\beta+n)^2(\alpha+\beta+n+1)}$  and  $\mathbb{E}[\theta|Y = n, \alpha, \beta] = \frac{\alpha+n}{\alpha+\beta+n}$  and  
 306  $\mathbb{V}[\theta|Y = n, \alpha, \beta] = \frac{(\alpha+n)\beta}{(\alpha+\beta+n)^2(\alpha+\beta+n+1)}$ . Note, the posterior variances are equivalent at these  
 307 thresholds for all  $\alpha = \beta$ .

308 Parameters were given an uninformative  $\text{Beta}(1, 1)$  prior, which is equivalent to a standard  
 309 uniform ( $\text{Uniform}(0, 1)$ ) prior since it places equal probability on all parameter values within  
 310 its support. This distribution has an expected value of  $\mu = \frac{1}{2} = 0.500$  and a variance of  
 311  $\sigma^2 = \frac{1}{12} \approx 0.083$ . Further, the posterior is  $\text{Beta}(Y+1, n-Y+1)$ , from which various moments  
 312 such as the expected value  $\mathbb{E}[\theta|Y] = \frac{Y+1}{n+2}$  and variance  $\mathbb{V}[\theta|Y] = \frac{(Y+1)(n-Y+1)}{(n+2)^2(n+3)}$ , and other  
 313 quantities, can be easily calculated. Clearly,  $\mathbb{E}[\theta|Y = 0] = \frac{1}{n+2}$  and  $\mathbb{V}[\theta|Y = 0] = \frac{n+1}{(n+2)^2(n+3)}$ ,  
 314 and  $\mathbb{E}[\theta|Y = n] = \frac{n+1}{n+2}$  and  $\mathbb{V}[\theta|Y = n] = \frac{n+1}{(n+2)^2(n+3)}$ . For example, with  $n = 2$  sampled  
 315 specimens for a given species (the minimum sample size possible) having no overlapping  
 316 genetic distances, on average,  $\theta$  will be equal to  $\mu = \frac{1}{4} = 0.250$ , with a variance of  
 317  $\sigma^2 = \frac{3}{80} \approx 0.0375$ . In contrast, with  $n = 100$  and no overlap, the mean reduces to  
 318  $\mu = \frac{1}{102} \approx 0.001$ , with a variance of  $\sigma^2 = 9.425 \times 10^{-5}$ . When  $n = 2$  and both taxon distances  
 319 overlap, the mean is  $\mu = \frac{3}{4} = 0.750$ , having a variance of  $\sigma^2 \approx 0.0375$ . For  $n = 100$  where  
 320 all distances overlap,  $\theta$  has expectation  $\mu = \frac{101}{102} \approx 0.990$  and variance  $\sigma^2 \approx 9.425 \times 10^{-5}$ .

321 These edge case examples demonstrate that the Beta(1, 1) distribution serves as a good  
 322 first-pass prior for DNA barcode gap estimation using Phillips et al.'s (2024) metrics, though  
 323 tighter bounds can still be achieved. In general however, when possible, it is always advisable  
 324 to incorporate prior information, even if only weak, rather than simply imposing complete  
 325 ignorance in the form of a flat prior distribution. In the case of unimodal distributions, the  
 326 (estimated) posterior mean often possesses the property that it readily decomposes into a  
 327 convex linear combination, in the form of a weighted sum, of the (estimated) prior mean  
 328 and the MLE. That is  $\hat{\mu}_{\text{posterior}} = w\hat{\mu}_{\text{prior}} + (1 - w)\hat{\mu}_{\text{MLE}}$ , where for the beta distribution,  
 329  $w = \frac{\alpha + \beta}{\alpha + \beta + n}$ . Therefore, with sufficient data,  $w \rightarrow 0$  as  $n \rightarrow \infty$ , regardless of the values of  $\alpha$   
 330 and  $\beta$ , and the choice of prior distribution becomes less important since the posterior will be  
 331 dominated by the likelihood. For the Beta(1, 1),  $w = \frac{2}{2+n}$ , with  $n = 2$  giving  $w = \frac{1}{2} = 0.500$ ;  
 332 that is, the posterior is the arithmetic average of the prior and the likelihood. For  $n = 100$ ,  
 333 the prior weight is  $w = \frac{2}{102} \approx 0.0196$ . The full Bayesian model for species  $x$  is thus given by

$$\begin{aligned}
 y_{\text{lwr}} &\sim \text{Binomial}(N, p_{\text{lwr}}) \\
 y_{\text{upr}} &\sim \text{Binomial}(M, p_{\text{upr}}) \\
 y'_{\text{lwr}} &\sim \text{Binomial}(N, p'_{\text{lwr}}) \\
 y'_{\text{upr}} &\sim \text{Binomial}(C, p'_{\text{upr}}) \\
 p_{\text{lwr}}, p_{\text{upr}}, p'_{\text{lwr}}, p'_{\text{upr}} &\sim \text{Beta}(1, 1).
 \end{aligned} \tag{12}$$

334 Note that  $p_x$ ,  $q_x$ ,  $p'_x$ , and  $q'_x$  in Equations (1)-(4) are denoted  $p_{\text{lwr}}$ ,  $p_{\text{upr}}$ ,  $p'_{\text{lwr}}$ ,  $q'_{\text{upr}}$  within  
 335 Equation (12) for easy distinction between MLEs and Bayesian posterior estimates. The  
 336 above statistical theory and derivations lay a good foundation for the remainder of this  
 337 paper.

338 The proposed model is inherently vectorized to allow processing of multiple species  
 339 datasets simultaneously. Model fitting was achieved using the Stan probabilistic



programming language (Carpenter et al., 2017) framework for Hamiltonian Monte Carlo (HMC) via the No-U-Turn Sampler (NUTS) sampling algorithm (Hoffman and Gelman, 2014) through the `rstan` R package (version 2.32.6) (Stan Development Team, 2023) in R (version 4.4.1) (R Core Team, 2024). Four Markov chains were run for 2000 iterations each in parallel across four cores with random parameter initializations. Within each chain, a total of 1000 samples was discarded as warmup (*i.e.*, burnin) to reduce dependence on starting conditions and to ensure posterior samples are reflective of the equilibrium distribution. Further, 1000 post-warmup draws were utilized per chain during the sampling phase. Because HMC/NUTS results in dependent samples that are minimally autocorrelated, chain thinning is not required. Each of these tuning parameters reflect default Markov Chain Monte Carlo (MCMC) settings in Stan to control both bias and variance respectively in the resulting draws. All analyses in the present work were carried out on a 2023 Apple MacBook Pro with M2 chip and 16 GB RAM running macOS Ventura 13.2. A random seed was set to ensure reproducibility of model results. Outputted estimates were rounded to three decimal places of precision. Posterior distributions were visualized as KDE plots using the `ggplot2` R package (version 3.5.1) (Wickham, 2016) with the default Gaussian kernel and optimal smoothness selection. To successfully run the Stan program, end users must have installed an appropriate compiler (such as GCC or Clang) which is compatible with their operating system, such as macOS.

Convergence was assessed both visually and quantitatively as follows: (1) through examining parameter traceplots, which depict the trajectory of accepted MCMC draws as a function of the number of iterations, (2) through monitoring the Gelman-Rubin potential scale reduction factor statistic ( $\hat{R}$ ) (Gelman and Rubin, 1992; Vehtari et al., 2021), which measures the concordance of within-chain *versus* between-chain variance, and (3) through calculating the effective sample size (ESS) for each parameter, which quantifies the number of independent samples generated Markov chains are equivalent to. Mixing of chains was deemed sufficient when traceplots looked like “fuzzy caterpillars”,  $\hat{R} < 1.01$ , and effective

sample sizes were reasonably large (Gelman et al., 2020). After sampling, a number of summary quantities were reported, including posterior means, posterior SDs, and posterior quantiles from which 95% CrIs could be computed to make probabilistic inferences concerning true population parameters. To validate the overall correctness of the proposed statistical model given by Equation (12), as a means of comparison, posterior predictive checks (PPCs) were also employed to generate binomial random variates in the form of counts from the posterior predictive distribution; that is  $\gamma = \{Np_x, Mq_x, Np'_x, Cq'_x\}$  to verify that the model adequately captures relevant features of the observed data. The proposed Bayesian model outlined here has a straightforward interpretation (**Table 1**).

### 3 Results and Discussion

The *Agabus* CYTB dataset analyzed by Phillips et al. (2024) is revisited herein for the species *A. bipustulatus* and *A. nevadensis*, since these taxa were the sole representatives for this locus, with the most and the least specimen records, respectively ( $N = 701$  and  $N = 2$ ) across all three assessed molecular markers. Briefly, using the R package **MACER** (Young et al., 2021), DNA sequences were downloaded from GenBank and BOLD and processed to obtain a 343 bp FASTA alignment representing 46 unique haplotypes. Genetic distances were calculated using uncorrected p-distances. Further, *A. bipustulatus* comprised 46 total haplotypes, whereas *A. nevadensis* possessed two haplotypes. Note, DNA barcode gap estimation is only possible for species having at least two specimen records. This dataset is a prime illustrative example highlighting the issue of inadequate taxon sampling, which arises frequently in large-scale phylogenetic and phylogeographic studies, in several respects. First, from a statistical viewpoint, sample sizes reflect extremes in reliable parameter estimation. Second, from a DNA barcoding perspective, *Agabus* currently comprises about 200 extant species according to the Global Biodiversity Information Facility (GBIF) (<https://www.gbif.org>); yet, due to the level of convenience sampling inherent in taxonomic

**Table 1:** Interpretation of the DNA barcode gap estimators within  $[a/a', b]$

| Parameter              | Explanation                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $p_x/p_{\text{lwr}}$   | When $p_{\text{lwr}}$ is close to 0 (1), it suggests that the probability of intraspecific (interspecific) distances being larger (smaller) than interspecific (intraspecific) distances is low (high) on average, while the probability of interspecific (intraspecific) distances being larger (smaller) than intraspecific (interspecific) distances is high (low) on average; that is, there is (no) evidence for a DNA barcode gap.                                                                                                                                                                                                                                                                                      |
| $q_x/p_{\text{upr}}$   | When $p_{\text{upr}}$ is close to 0 (1), it suggests that the probability of interspecific (intraspecific) distances being larger (smaller) than intraspecific (interspecific) distances is high (low) on average, while the probability of intraspecific (interspecific) distances being larger (smaller) than interspecific (intraspecific) distances is low (high) on average; that is, there is (no) evidence for a DNA barcode gap.                                                                                                                                                                                                                                                                                      |
| $p'_x/p'_{\text{lwr}}$ | When $p'_{\text{lwr}}$ is close to 0 (1), it suggests that the probability of intraspecific (combined interspecific distances for a target species and its nearest neighbour species) distances being larger than combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) is low (high) on average, while the probability of combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) being larger than intraspecific distances (combined interspecific distances for a target species and its nearest neighbour species) is high (low) on average; that is, there is (no) evidence for a DNA barcode gap. |
| $q'_x/p'_{\text{upr}}$ | When $p'_{\text{upr}}$ is close to 0 (1), it suggests that the probability of combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) being larger than intraspecific distances (combined interspecific distances for a target species and its nearest neighbour species) is high (low) on average, while the probability of intraspecific distances (combined interspecific distances for a target species and its nearest neighbour species) being larger than combined interspecific distances for a target species and its nearest neighbour species (intraspecific distances) is low (high) on average; that is, there is (no) evidence for a DNA barcode gap. |

collection efforts for this genus, adequate representation of species and genetic diversity is far from complete.

MCMC parameter traceplots showed rapid mixing of chains to the stationary distribution (**Supplementary Figure 1**). Further, all  $\hat{R}$  and ESS values (not shown) were close to their recommended cutoffs of one and thousands of samples, respectively, indicating chains are both well-mixed and have converged to the posterior distribution.

Bayesian posterior estimates were reported alongside frequentist MLEs, in addition to SEs, posterior SDs, 95% CIs and 95% CrIs (**Table 2**).


**Table 2:** Nonparametric frequentist and Bayesian estimates of distance distribution overlap/separation for the DNA barcode gap coalescent model parameters applied to *A. bipustulatus* ( $N = 701$ ) and *A. nevadensis* ( $N = 2$ ) for CYTB, including 95% CIs and CrIs. CrIs are based on 4000 posterior draws. All parameter estimates are reported to three decimal places of precision.

| Species                | Parameter              | MLE (SE, 95% CI)            | Bayes Est. (SD; 95% CrI)   |
|------------------------|------------------------|-----------------------------|----------------------------|
| <i>A. bipustulatus</i> | $p_x/p_{\text{lwr}}$   | 1.000 (0.000; 1.000-1.000)  | 1.000 (0.000; 1.000-1.000) |
| <i>A. bipustulatus</i> | $q_x/p_{\text{upr}}$   | 1.000 (0.000; 1.000-1.000)  | 1.000 (0.000; 0.999-1.000) |
| <i>A. bipustulatus</i> | $p'_x/p'_{\text{lwr}}$ | 1.000 (0.000; 1.000-1.000)  | 1.000 (0.000; 1.000-1.000) |
| <i>A. bipustulatus</i> | $q'_x/p'_{\text{upr}}$ | 1.000 (0.000; 1.000-1.000)  | 1.000 (0.000; 0.999-1.000) |
| <i>A. nevadensis</i>   | $p_x/p_{\text{lwr}}$   | 1.000 (0.000; 1.000-1.000)  | 0.835 (0.144; 0.470-0.996) |
| <i>A. nevadensis</i>   | $q_x/p_{\text{upr}}$   | 0.010 (0.002; 0.006-0.014)  | 0.010 (0.002; 0.007-0.014) |
| <i>A. nevadensis</i>   | $p'_x/p'_{\text{lwr}}$ | 1.000 (0.000; 1.000-1.000)  | 0.834 (0.138; 0.481-0.994) |
| <i>A. nevadensis</i>   | $q'_x/p'_{\text{upr}}$ | 0.010 (0.070; -0.128-0.148) | 0.010 (0.002; 0.007-0.014) |

CIs were calculated using the usual large sample  $(1 - \alpha)100\%$ -level interval estimate given by  $\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$ , where  $z_{1-\frac{\alpha}{2}} = 1.960$  is the critical value for 95% confidence (*i.e.*, the 97.5th percent quantile from the standard Normal distribution), and  $\alpha$  is the stated significance level (here, 5%). Given a  $(1 - \alpha)100\%$  CI, with repeated sampling, on average  $(1 - \alpha)100\%$  of constructed intervals will contain the true parameter of interest, assuming a correctly specified statistical model; on the other hand, any given CI will either capture or exclude the true parameter with 100% certainty. This in stark contrast to a CrI, where the true parameter is contained within said interval with  $(1 - \alpha)100\%$  probability, contingent on the data generating model being correct. Note, by default Stan computes equal-tailed

(central) CrIs such that there is equal area situated in the left and right tails of the posterior distribution. For a 95% CrI, this corresponds to the 2.5th and 97.5th percent quantiles. However, constructed intervals are usually only valid for symmetric or nearly symmetric distributions. Given the bounded nature of the DNA barcode gap metrics, whose posterior distributions, as expected, show considerable skewness, an alternative approach to reporting CrIs, such as Highest Posterior Density (HPD) intervals (Chen and Shao, 1999) or shortest probability intervals (SPIn) (Liu et al., 2015) is warranted. As such asymmetric intervals generally attain greater statistical efficiency (in the form of smaller Mean Squared Error (MSE) or variance) and higher coverage probabilities than more standard interval estimates, careful in-depth comparison is left for future work.

Findings based on nonparametric MLEs and Bayesian posterior means were quite comparable with one another and show evidence of complete overlap in intraspecific, interspecific, and combined interspecific distances for *A. bipustulatus* in both the  $p/q/p_{\text{lwr}}/p_{\text{upr}}$  and  $p'/q'/p'_{\text{lwr}}/p'_{\text{upr}}$  directions since the metrics attain magnitudes very close to one, with minimal noise (**Table 2**). As a result, this likely indicates that no DNA barcode gap is present for this species. Such findings are strongly reinforced by the very tight clustering of posterior draws (**Figure 2**) and associated interval estimates owing to the large number of specimens sampled for this species.



**Figure 2:** Scatterplots (black solid points) and distributions (black solid lines) depicting the DNA barcode gap metrics for *A. bipustulatus* ( $N = 701$ ) and *A. nevadensis* ( $N = 2$ ) across CYTB based on 4000 Bayesian posterior draws. MLEs and posterior means are displayed as coloured (red/blue) solid and dashed lines for the metrics, respectively.

On the other hand, the situation for *A. nevadensis* is more nuanced, as posterior values are further spread out (**Table 2** and **Figure 2**), suggesting less overall certainty in true parameter values given the low specimen sampling coverage for this taxon. Of note, calculated SEs and SDs are large, with the 95% CIs and 95% CrIs being quite wide in most cases, consistent with much uncertainty regarding the computed frequentist and Bayesian posterior means of the DNA barcode gap metrics. For instance, the Bayesian analysis suggests that the data are consistent with both  $p_{\text{low}}$  and  $p'_{\text{low}}$  ranging from approximately 0.500-1.000 due to the large SD of about 0.140. The posterior means associated with these estimates themselves are also far from one. The CrI for  $p_{\text{upr}}$  and  $p'_{\text{upr}}$  also spans an order of magnitude. Further, regarding the frequentist analysis for the same species, the 95% CI for  $q_x$  is quite wide, reflecting considerable uncertainty in its true parameter value. Similarly, that for  $q'_x$  extends to negative values at the left endpoint, due to the corresponding SE of 0.070 being too high as

a result of the extremely low sample size of  $n = 2$  individuals sampled (**Table 2**). Since the 95% CI for  $q'_x$  truncated at the lower endpoint includes the value of zero, the null hypothesis for the presence of a DNA barcode gap cannot be rejected. Despite this, it is worth noting that truncation is not standard statistical practice and will likely lead to an interval with less than 95% nominal coverage. In such cases, more appropriate confidence interval methods like the Wilson score interval, the exact (Clopper-Pearson) interval, or the Agresti-Coull interval should be employed (Newcombe, 1998; Agresti and Coull, 1998). KDEs for *A. bipustulatus* are strongly left (negatively) skewed (**Figure 2**), whereas those for *A. nevadensis* exhibit more symmetry, especially for  $p_{\text{upr}}$  and  $p'_{\text{upr}}$  (**Figure 2**). These differences are likely due to the stark contrast in sample sizes for the two examined species. Nevertheless, simulated counts of overlapping specimen records from the posterior predictive distribution (**Supplementary Table 1**) were found to be very close to observed counts for both species, indicating that the proposed model adequately captures underlying variation. Obtained results suggest that use of the Beta(1, 1) prior may not be appropriate given a low number of collected individuals for most taxa in DNA barcoding efforts. This suggests that further consideration of more informative beta priors is worthwhile. This is clear for *A. nevadensis*, where use of a stronger prior will likely eliminate the issue regarding accurate estimation of  $p_{\text{lwr}}$  and  $p'_{\text{lwr}}$  and stabilize their uncertainty.

## 4 Conclusion

Herein, the accuracy of the DNA barcode gap was analyzed from a rigorous statistical lens to expedite both the curation and growth of reference sequence libraries, ensuring they are populated with high quality, statistically defensible specimen records fit for purpose to address standing questions in ecology, evolutionary biology, management, and conservation. To accomplish this, recently proposed, easy to calculate nonparametric MLEs were formally derived using ECDFs and applied to assess the extent of overlap/separation of distance

distributions within and among two species of predatory water beetles in the genus *Agabus* sequenced at CYTB using a Bayesian binomial count model with conjugate beta priors. Findings highlight a high level of parameter uncertainty for *A. nevadensis*, whereas posterior estimates of the DNA barcode gap metrics for *A. bipustulatus* are much more certain. Based on these results, it is imperative that specimen sampling be prioritized to better reflect actual species boundaries. More generally, apart from the metrics being employed to better highlighting the importance of within-species genetic diversity versus between-species divergence, it is expected that the approach developed herein will be of broad utility in applied fields, such as DNA-based detection of seafood fraud within global supply chains, and in the determination of species occupancy/detection probabilities at ecological sites of interest using active and passive environmental DNA (eDNA) methods such as metabarcoding.

Since the DNA barcode gap metrics often attain values very close to zero (suggesting no overlap and complete separation of distance distributions) and/or very near one (indicating no separation and complete overlap), in addition to more intermediate values, a noninformative  $\text{Beta}(\frac{1}{2}, \frac{1}{2})$  prior may be more appropriate over complete ignorance imposed by a  $\text{Beta}(1, 1)$  prior. The former distribution is U-shaped symmetric and places greater probability density at the extremes of the distribution due to its heavier tails, while still allowing for variability in parameter estimates within intermediate values along its domain. Note that this prior is Jeffreys' prior density (Jeffreys, 1946), which is proportional to the square root of the Fisher information  $\mathcal{I}(\theta)$ ; that is,  $\pi(\theta) \propto \theta^{-\frac{1}{2}}(1 - \theta)^{-\frac{1}{2}}$ . Jeffreys' prior has several desirable statistical properties as a prior: that it is inversely proportional to the standard deviation of the binomial distribution, and most notably, that it is invariant to model reparameterization (Gelman et al., 2014). However, this prior can lead to divergent transitions, among other pathologies, imposed by complex geometry (*i.e.*, curvature) in the posterior space since many iterative stochastic MCMC sampling algorithms experience difficulties when exploring high density distribution regions. Thus, remedies to resolve them, such as lowering the step size of the HMC/NUTS sampler, should be attempted in future work, along with other approaches



such as empirical Bayes estimation to approximate beta prior hyperparameters from observed data through the MLE or other methods of parameter estimation, such as the method of moments. Alternatively, hierarchical modelling could be employed to estimate separate distribution model hyperparameters for each species and/or compute distinct estimates for the directionality/comparison level of the DNA barcode gap metrics (*i.e.*, lower *vs.* upper, non-prime *vs.* prime) separately within the genus under study. This would permit greater flexibility through incorporating more fine-grained structure seen in the data; however, low taxon sample sizes may preclude valid inferences to be reasonably ascertained due to the large number additional parameters which would be introduced through the specification of the hyperprior distributions. Methods outlined in Gelman et al. (2014), such as dealing with non-exchangeability of observations and alternate model parameterizations like the logit, may prove useful in this regard. Perhaps, the best course of action though is the elicitation of priors from domain experts. Even though more work remains, it is clear that both frequentist and Bayesian inference hold much promise for the future of molecular biodiversity science.

## Supplementary Information

None declared.

## Data Availability Statement

Raw data, R, and Stan code can be accessed via Dryad at:

<http://datadryad.org/stash/share/>

RZIfMixcEODe0RWP7eyXWQewSVbqEIA9UTrH3ZVKyn4.

A GitHub repository can be found at:

<https://github.com/jphill01/Bayesian-DNA-Barcode-Gap-Coalescent>.

## Acknowledgements

We wish to recognise the valuable comments and discussions of Daniel (Dan) Gillis, Robert (Bob) Hanner, Robert (Rob) Young, and XXX anonymous reviewers.

We acknowledge that the University of Guelph resides on the ancestral lands of the Attawandaron people and the treaty lands and territory of the Mississaugas of the Credit. We recognize the significance of the Dish with One Spoon Covenant to this land and offer our respect to our Anishinaabe, Haudenosaunee and Métis neighbours as we strive to strengthen our relationships with them.

## Funding

None declared.

## Conflict of Interest

None declared.

## Author Contributions

JDP wrote the manuscript, wrote R and Stan code, as well as analyzed and interpreted all model results.

## References

Agresti, A. and B. A. Coull

1998. Approximate is better than ‘exact’ for interval estimation of binomial proportions. *The American Statistician*, 52(2):119–126.

Ahrens, D., F. Fujisawa, H.-J. Krammer, J. Eberle, S. Fabrizi, and A. Vogler

2016. Rarity and incomplete sampling in DNA-based species delimitation. *Systematic Biology*, 65(3):478–494.

Avise, J., J. Arnold, R. Ball, Jr., E. Bermingham, T. Lamb, J. Neigel, C. Reeb, and N. Saunders

1987. Intraspecific phylogeography: The mitochondrial DNA bridge between population genetics and systematics. *Annu. Rev. Ecol. Syst.*, 18:489–522.

Bartlett, S. and W. Davidson

1992. FINS (forensically informative nucleotide sequencing): A procedure for identifying the animal origin of biological specimens. *BioTechniques*, 12(3):408–411.

Bergsten, J., D. Bilton, T. Fujisawa, M. Elliott, M. Monaghan, M. Balke, L. Hendrich,

J. Geijer, J. Herrmann, G. Foster, I. Ribera, A. Nilsson, T. Barraclough, and A. Vogler

2012. The effect of geographical scale of sampling on DNA barcoding. *Systematic Biology*, 61(5):851–869.

Bouckaert, R., T. G. Vaughan, J. Barido-Sottani, S. Duchêne, M. Fourment,

A. Gavryushkina, J. Heled, G. Jones, D. Kühnert, N. De Maio, M. Matschiner, F. K.

Mendes, N. F. Müller, H. A. Ogilvie, L. du Plessis, A. Poppinga, A. Rambaut, D. Rasmussen,  
I. Siveroni, M. A. Suchard, C.-H. Wu, D. Xie, C. Zhang, T. Stadler, and A. J. Drummond  
2019. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis.  
*PLOS Computational Biology*, 15(4):1–28.

Čandek, K. and M. Kuntner

2015. DNA barcoding gap: Reliable species identification over morphological and  
geographical scales. *Molecular Ecology Resources*, 15(2):268–277.

Carpenter, B., A. Gelman, M. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker,  
J. Guo, P. Li, and A. Riddell

2017. Stan: A probabilistic programming language. *Journal of Statistical Software*, 76:1.

Carstens, B. C., T. A. Pelletier, N. M. Reid, and J. D. Satler

2013. How to fail at species delimitation. *Molecular Ecology*, 22(17):4369–4383.

Chen, M.-H. and Q.-M. Shao

1999. Monte Carlo estimation of Bayesian credible and HPD intervals. *Journal of  
Computational and Graphical Statistics*, 8(1):69–92.

Collins, R. A. and R. H. Cruickshank

2013. The seven deadly sins of DNA barcoding. *Molecular Ecology Resources*,  
13(6):969–975.

Collins, R. A. and R. H. Cruickshank

2014. Known knowns, known unknowns, unknown unknowns and unknown knowns in dna  
barcoding: A comment on dowton et al. *Systematic Biology*, 63(6):1005–1009.

de Queiroz, K.

2007. Species concepts and species delimitation. *Systematic Biology*, 56(6):879–886.

- Dempster, A. P., N. M. Laird, and D. B. Rubin  
1977. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–22.
- Flouri, T., X. Jiao, B. Rannala, and Z. Yang  
2018. Species tree inference with BPP using genomic sequences and the multispecies coalescent. *Molecular Biology and Evolution*, 35(10):2585–2593.
- Fonseca, E. M., D. J. Duckett, and B. C. Carstens  
2021. P2C2M.GMYC: An R package for assessing the utility of the Generalized Mixed Yule Coalescent model. *Methods in Ecology and Evolution*, 12(3):487–493.
- Fujisawa, T. and T. Barraclough  
2013. Delimiting species using single-locus data and the generalized mixed yule coalescent approach: A revised method and evaluation on simulated data sets. *Systematic Biology*, 62(5):707–724.
- Gelman, A., J. Carlin, H. Stern, D. Duncan, A. Vehtari, and D. Rubin  
2014. *Bayesian Data Analysis*, third edition. Chapman and Hall/CRC.
- Gelman, A. and D. Rubin  
1992. Inference from iterative simulation using multiple sequences. *Statistical Science*, 7(4):457–472.
- Gelman, A., A. Vehtari, D. Simpson, C. Margossian, B. Carpenter, Y. Yao, L. Kennedy, J. Gabry, P.-C. Bürkner, and M. Modrák  
2020. Bayesian workflow.
- Hebert, P., A. Cywinska, S. Ball, and J. deWaard  
2003a. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(1512):313–321.

Hebert, P., S. Ratnasingham, and J. de Waard

2003b. Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(Suppl 1):S96–S99.

Hebert, P. D., M. Y. Stoeckle, T. S. Zemlak, and C. M. Francis

2004. Identification of birds through DNA barcodes. *PLoS Biol*, 2(10):e312.

Hickerson, M. J., C. P. Meyer, and C. Moritz

2006. DNA barcoding will often fail to discover new animal species over broad parameter space. *Systematic Biology*, 55(5):729–739.

Hoffman, M. and A. Gelman

2014. The No-U-Turn Sampler: Adaptively setting path lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, 15:1593–1623.

Hubert, N. and R. Hanner

2015. DNA barcoding, species delineation and taxonomy: A historical perspective. *DNA Barcodes*, 3:44–58.

Hubert, N., J. Phillips, and R. Hanner

2024. Delimiting species with single-locus dna sequences. In *DNA Barcoding*, R. DeSalle, ed., volume 2744 of *Methods in Molecular Biology*. New York, NY: Humana.

Jackson, N. D., B. C. Carstens, A. E. Morales, and B. C. O’Meara

2017a. Species delimitation with gene flow. *Systematic Biology*, 66(5):799–812.

Jackson, N. D., A. E. Morales, B. C. Carstens, and B. C. O’Meara

2017b. PHRAPL: Phylogeographic inference using approximate likelihoods. *Systematic Biology*, 66(6):1045–1053.

Jeffreys, H.

1946. An invariant form for the prior probability in estimation problems. *Proceedings*

620 of the Royal Society of London. *Series A, Mathematical and Physical Sciences*,  
621 186(1007):453–461.

622 Jukes, T. and C. Cantor  
623 1969. Evolution of protein molecules. In *Mammalian Protein Metabolism*, H. N. Munro,  
624 ed., Pp. 21–132. New York: Academic Press.

625 Kapli, P., S. Lutteropp, J. Zhang, K. Kobert, P. Pavlidis, A. Stamatakis, and T. Flouri  
626 2017. Multi-rate poisson tree processes for single-locus species delimitation under maximum  
627 likelihood and markov chain monte carlo. *Bioinformatics*, 33(11):1630–1638.

628 Kimura, M.  
629 1980. A simple method for estimating evolutionary rates of base substitutions  
630 through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*,  
631 16(1):111–120.

632 Kingman, J.  
633 1982a. The coalescent. *Stochastic Processes and Their Applications*, 13:235–248.

634 Kingman, J.  
635 1982b. On the genealogy of large populations. *Journal of Applied Probability*, 19(A):27–43.

636 Knowles, L. L. and W. P. Maddison  
637 2002. Statistical phylogeography. *Molecular Ecology*, 11(12):2623–2635.

638 Kullback, S. and R. Leibler  
639 1951. On information and sufficiency. *Annals of Mathematical Statistics*, 22(1):79–86.

640 Liu, Y., A. Gelman, and T. Zheng  
641 2015. Simulation-efficient shortest probability intervals. *Statistical Computing*, 25:809–819.

642 Mather, N., S. Traves, and S. Ho

643 2019. A practical introduction to sequentially Markovian coalescent methods for estimating  
644 demographic history from genomic data. *Ecology and Evolution*, 10(1):579–589.

645 Meier, R., G. Zhang, and F. Ali

646 2008. The use of mean instead of smallest interspecific distances exaggerates the size of  
647 the “barcoding gap” and leads to misidentification. *Systematic Biology*, 57(5):809–813.

648 Meyer, C. and G. Paulay

649 2005. DNA barcoding: Error rates based on comprehensive sampling. *PLOS Biology*,  
650 3(12):e422.

651 Mutanen, M., S. M. Kivelä, R. A. Vos, C. Doorende, S. Ratnasingham, A. Hausmann,  
652 P. Huemer, V. Dincă, E. J. van Nieukerken, C. Lopez-Vaamonde, R. Vila, L. Aarvik,  
653 T. Decaëns, K. A. Efetov, P. D. N. Hebert, A. Johnsen, O. Karsholt, M. Pentinsaari,  
654 R. Rougerie, A. Segerer, G. Tarmann, R. Zahiri, and H. C. J. Godfray

655 2016. Species-level para- and polyphyly in DNA barcode gene trees: Strong operational  
656 bias in european lepidoptera. *Systematic Biology*, 65(6):1024–1040.

657 Newcombe, R. G.

658 1998. Two-sided confidence intervals for the single proportion: comparison of seven  
659 methods. *Statistics in Medicine*, 17(8):857–872.

660 Pante, E., N. Puillandre, A. Viricel, S. Arnaud-Haond, D. Aurelle, M. Castelin, A. Chenuil,  
661 C. Destombe, D. Forcioli, M. Valero, F. Viard, and S. Samadi

662 2015. Species are hypotheses: Avoid connectivity assessments based on pillars of sand.  
663 *Molecular Ecology*, 24(3):525–544.

664 Paradis, E., J. Claude, and K. Strimmer

665 2004. Ape: Analyses of phylogenetics and evolution in r language. *Bioinformatics*,  
666 20(2):289–290.



- Pentinsaari, M., H. Salmela, M. Mutanen, and T. Roslin  
2016. Molecular evolution of a widely-adopted taxonomic marker (COI) across the animal  
tree of life. *Scientific Reports*, 6:35275.
- Phillips, J., D. Gillis, and R. Hanner  
2022. Lack of statistical rigor in DNA barcoding likely invalidates the presence of a true  
species' barcode gap. *Frontiers in Ecology and Evolution*, 10:859099.
- Phillips, J., C. Griswold, R. Young, N. Hubert, and H. Hanner  
2024. *A Measure of the DNA Barcode Gap for Applied and Basic Research*, Pp. 375–390.  
New York, NY: Springer US.
- Pons, J., T. G. Barraclough, J. Gomez-Zurita, A. Cardoso, D. P. Duran, S. Hazell,  
S. Kamoun, W. D. Sumlin, and A. P. Vogler  
2006. Sequence-based species delimitation for the DNA taxonomy of undescribed insects.  
*Systematic Biology*, 55(4):595–609.
- Puillandre, N., S. Brouillet, and G. Achaz  
2021. ASAP: Assemble Species by Automatic Partitioning. *Molecular Ecology Resources*,  
21(2):609–620.
- Puillandre, N., A. Lambert, S. Brouillet, and G. Achaz  
2011. ABGD, Automatic Barcode Gap Discovery for primary species delimitation.  
*Molecular Ecology*, 21(8):1864–1877.
- R Core Team  
2024. *R: A Language and Environment for Statistical Computing*. R Foundation for  
Statistical Computing, Vienna, Austria.
- Rannala, B.  
2015. The art and science of species delimitation. *Current Zoology*, 61(5):846–853.

- Rannala, B. and Z. Yang  
2003. Bayes estimation of species divergence times and ancestral population sizes using DNA sequences from multiple loci. *Genetics*, 164:1645–1656.
- Rannala, B. and Z. Yang  
2017. Efficient Bayesian species tree inference under the multispecies coalescent. *Systematic Biology*, 66(5):823–842.
- Ratnasingham, S. and P. Hebert  
2007. BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, 7(3):355–364.
- Ratnasingham, S. and P. D. N. Hebert  
2013. A dna-based registry for all animal species: The barcode index number (bin) system. *PLoS One*, 8(7):e66213.
- Stan Development Team  
2023. RStan: The R interface to Stan. R package version 2.32.6.
- Stoeckle, M. and D. Thaler  
2014. Dna barcoding works in practice but not in (neutral) theory. *PLoS One*, 9(7):e100755.
- Vehtari, A., A. Gelman, D. Simpson, B. Carpenter, and P.-C. Bürkner  
2021. Rank-normalization, folding, and localization: An improved  $\hat{R}$  for assessing convergence of MCMC (with discussion). *Bayesian Analysis*, 16(2):667–718.
- Wells, T., T. Carruthers, P. Muñoz-Rodríguez, A. Sumadijaya, J. R. I. Wood, and R. W. Scotland  
2022. Species as a heuristic: Reconciling theory and practice. *Systematic Biology*, 71(5):1233–1243.
- Wickham, H.  
2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

716 Wiemers, M. and K. Fiedler  
717 2007. Does the DNA barcoding gap exist? - a case study in blue butterflies (Lepidoptera:  
718 Lycaenidae). *Frontiers in Zoology*, 4(8).

719 Yang, Z. and B. Rannala  
720 2010. Bayesian species delimitation using multilocus sequence data. *Proceedings of the*  
721 *National Academy of Sciences*, 107:9264–9269.

722 Yang, Z. and B. Rannala  
723 2014. Unguided species delimitation using dna sequence data from multiple loci. *Molecular*  
724 *Biology and Evolution*, 31(12):3125–3135.

725 Yang, Z. and B. Rannala  
726 2017. Bayesian species identification under the multispecies coalescent provides significant  
727 improvements to DNA barcoding analyses. *Molecular Ecology*, 26:3028–3036.

728 Young, R., R. Gill, D. Gillis, and R. Hanner  
729 2021. Molecular Acquisition, Cleaning and Evaluation in R (MACER) - A tool to assemble  
730 molecular marker datasets from BOLD and GenBank. *Biodiversity Data Journal*, 9:e71378.

731 Zhang, J., P. Kapli, P. Pavlidis, and A. Stamatakis  
732 2013. A general species delimitation method with applications to phylogenetic placements.  
733 *Bioinformatics*, 29(22):2869–2876.