



HIP/MP Comparisons and HIP Results

Jim Pivarski

Alexei Safonov

Texas A&M University

25 May, 2009




- ▶ Pablo and I discussed implementation details early in the week
- ▶ We synchronized all of the inputs and many of the parameters/cuts, and set up parallel CRAFT alignments and tests in cosmic ray MC
- ▶ We didn't finish our comparison— I have not seen all of the MP results— and we never got a chance to discuss how we might converge (schedule conflicts? a bad week?)
- ▶ What I know is the following:
 - ▶ even with the same inputs and similar cuts, HIP and MP differ in data by 2–4 mm, 0.5–2 mrad, systematic trend in y , z
 - ▶ HIP yields high-accuracy results in Monte Carlo
- ▶ What I'll present here:
 - ▶ comparative study of HIP and MP implementations and results
 - ▶ possible explanations for the discrepancy
 - ▶ my HIP recommendation, and arguments to support it as a good sub-millimeter alignment

Comparison of HIP and MP (1/3) Jim Pivarski 3/40



- Inputs to the procedures (differences between HIP and MP in blue)
 - V11_StreamMuAIGlobalCosmics_227_Tosca090216_FromTrackerPointing_v5 (RunReg 3.8 T only)
 - newest tracker alignment (c. May 19)
 - last month's tracker APEs (c. Apr 24): update not available
 - tracker hits ≥ 15 , $\chi^2/N_{\text{dof}} < 10$, TIB and TOB only
 - high momentum: $100 < p_T < 200$ GeV
 - latest magnetic field: "grid_1103l_090322_3.8t"
 - latest internal DT alignment (agreement between tracks and survey)
 - HIP: CMSSW_2_2_11, MP: CMSSW_2_2_10 (very likely no difference for our purposes)



- Newest tracker alignment is the light blue one
 - New χ^2/N_{dof} peaks at 1.4 without APEs, old peaks at 1.6
 - Tracker alignment still needs to be centered and APEs need to be produced
- Final muon alignment will need to be consistent with final tracker alignment









- ▶ Alignables/parameters (differences between HIP and MP in blue)
 - ▶ only wheels $-1, 0, +1$, all sectors except 1 and 7
 - ▶ stations 1–3: 6 degrees of freedom
 - ▶ station 4: HIP: x, ϕ_y, ϕ_z , MP: x, ϕ_y (we'll compare only stations 1–3)
- ▶ Algorithmic implementation
 - ▶ no treatment of \vec{B} -field, dE/dx effects (HIP and MP implementations differ, so we turned them both off; note that $p_T > 100$ GeV)
 - ▶ residuals calculated from standard CMSSW track-refits with muon chamber APEs $\rightarrow \infty$ (1000 cm)
 - ▶ segment residuals: HIP: linear fit to (extrapolated track – hits), MP: (extrapolated track) – (linear fit to hits) (negligible: $\vec{B} \approx 0$ inside DTs)
 - ▶ cut on muon residuals: HIP: keep residuals tails (cut only unphysical values), MP: cut 1σ symmetrically around the peak
 - ▶ treatment of residuals tails: HIP: fit tail shape and misalignment together, MP: calculate misalignment from matrix inversion of hits
 - ▶ residuals weights: HIP: segment residual $(\chi^2/N_{\text{dof}})^{-1}$, MP: the same?
 - ▶ iteration: HIP: 2 required, 3 applied, MP: 1 applied

We'll discuss the algorithmic differences soon

Comparison of HIP and MP (3/3) Jim Pivarski 5/40



- ▶ Absolute differences (MP – HIP) on the order of 0.5–2.5 mm, 0.5–2 mrad
- ▶ In this comparison, station 4 is excluded, as are 3 failed fits (too few hits):
wh–1, st2, sec08 wh+1, st2, sec02 wh+1, st3, sec08
- ▶ Orientation of local x , y , z directions are ideal (for symmetric comparison):




This plot now uses the latest tracker for both HIP and MillePede!

Comparison of HIP and MP (3/3) Jim Pivarski 6/40



- ▶ Absolute differences (MP – HIP) on the order of 0.5–2.5 mm, 0.5–2 mrad
- ▶ In this comparison, station 4 is excluded, as are 3 failed fits (too few hits):
wh–1, st2, sec08 wh+1, st2, sec02 wh+1, st3, sec08
- ▶ Orientation of local x , y , z directions are ideal (for symmetric comparison):



This plot now uses the latest tracker for both HIP and MillePede!



- ▶ Most of the implementation differences that couldn't be easily synchronized are deep in the algorithms
- ▶ These probably aren't responsible for the discrepancy in results:
 - ▶ \vec{B} -field, dE/dx controls (we turned them off!)
 - ▶ calculation of segment residuals
 - ▶ weighting of residuals
- ▶ These might have something to do with it:
 - ▶ treatment of tails in residuals
 - ▶ residuals \leftrightarrow alignment corrections matrix
- ▶ I'll present each implementation difference in order

Turning off \vec{B} correction







Jim Pivarski 8/40



- ▶ To avoid differences in our \vec{B} and dE/dx controls, we simply turned them off for this alignment

- ▶ Correction is irrelevant at $100 < p_T < 200$ GeV, anyway


(Difference between corrected and uncorrected HIP shown below)






- ▶ Both algorithms combine all hits in one chamber on one track to take advantage of 4 independent residuals: Δx , Δy , $\Delta \frac{dx}{dz}$, $\Delta \frac{dy}{dz}$
- ▶ The implementations differ:
 - ▶ HIP: linear fit to (track extrapolation – hits) error $\propto \Delta \text{curvature}$
 - ▶ MP: (track extrapolation) – (linear fit to hits) error $\propto \text{curvature}$

linear fit to (track extrapolation – hits)




(track extrapolation) – (linear fit to hits)



- ▶ Only matters when the muon is expected to curve significantly inside the chamber, e.g. ME1/1 and ME1/2
 - ▶ probably negligible the DTs



- ▶ “Bad” segments distort residuals distribution, so weight segment residuals by their $(\chi^2/N_{\text{dof}})^{-1}$ (good fits get the most weight)
- ▶ Normalized with $\langle(\chi^2/N_{\text{dof}})^{-1}\rangle = 1.0$, so that errors are meaningful
- ▶ The usual dangers with weights are:
 - ▶ parameterized hit uncertainties *might* be unrepresentative
 - ▶ unphysically low-weight events can bias a distribution (they appear as spikes in a histogram)
- ▶ Not an issue with these weights because:
 - ▶ consistency with a line is a geometric attribute, not very sensitive to the precise hit uncertainties
 - ▶ we exclude the lowest 1% of weights



- ▶ Necessary for good fits in wheel ± 2 , station 1 $\Delta \frac{dy}{dz}$ (above)



- ▶ Residuals are influenced by misalignment geometry and propagation/instrumental effects:
 - ▶ misalignments distort residuals according to an exact 6×4 matrix
 - ▶ some propagation errors are Gaussian (statistical error, multiple scattering, etc.) and some power-law (single-scattering)
 - ▶ propagation correlates Δx with $\Delta \frac{dx}{dz}$ and Δy with $\Delta \frac{dy}{dz}$



- ▶ sources of systematic error to be quantified or controlled: tracker misalignment, imperfect $\vec{B}(\vec{x})$ and material maps, internal DT misalignment
- ▶ All of the above are convoluted together in an 9 dimensional space:
 $(\Delta x, \Delta \frac{dx}{dz}, \Delta y, \Delta \frac{dy}{dz}, x \text{ position}, \frac{dx}{dz} \text{ angle}, y \text{ position}, \frac{dy}{dz} \text{ angle}, q/p_T)$




- ▶ Incorporate misalignment, Gaussian and Lorentzian propagation effects, with $\Delta x - \Delta \frac{dx}{dz}$ correlations, into a **single ansatz**
 - ▶ 6 alignment parameters, σ and Γ for each of the 4 residuals, 2 correlation parameters = 16 parameters for each DT_{station 1-3}
 - ▶ 3-way convolution:
$$\text{residuals} = [\text{Gaussian} \otimes \text{Lorentzian}](\text{misalignment})$$
- ▶ **Fit all chamber variables simultaneously** (unbinned log-likelihood)
 - ▶ include all the physical tails (cut unphysical $|\Delta x_i|, |\Delta y_i| < 1000$ cm)
 - ▶ seed fit with truncated means and standard deviations for stability
 - ▶ project the fit results on all axes to make sure it's working
- ▶ Usually separate two q/p_T bins to account for the $\vec{B}(\vec{x})$ and $dE/dx(\vec{x})$ instrumental effects (but turned off for this alignment)
- ▶ Tracker and DT internal alignment are external inputs, assumed to be correct and worth investigating as systematics studies
 - ▶ quantified misalignment scenarios test sensitivity in MC (e.g. CSA08)
 - ▶ non-pointing cosmic rays reveal global distortions (e.g. TEC)
 - ▶ p_T dependence provides some information (e.g. tracker curl study)

What MillePede does (3/4)

Jim Pivarski 13/40





1. Fit peaks of residuals distributions to Lorentzian (or Gaussian) to identify a [peak - σ , peak + σ] window
2. Compute the residuals \rightarrow alignment corrections inversion for all hits in the window



- All alignment parameters are fitted simultaneously, but residuals tails from propagation effects are explicitly cut out
 - this is a matrix-mean: in a 1-DOF case (e.g. aligning δ_x only), it is equivalent to a computation of the mean within the selected window
 - trade-off: loose cut lets in non-Gaussian tails, tight cut introduces a dependence on its value
- Control $\vec{B}(\vec{x})$ and $dE/dx(\vec{x})$ instrumental effects with a linear q/p_T fit (but turned off for this alignment)
- Same systematics issues as HIP



- ▶ The mean of a distribution with such a tight cut (deep in the bulk of the distribution) would depend on the placement of the cut boundary
- ▶ *However*, the perfect symmetry of $[\text{peak} - \sigma, \text{peak} + \sigma]$ should balance the residuals on each side of the peak, as long as the pre-fit is a good fit \checkmark
- ▶ *However*, the matrix inversion depends on trends in the residuals distributions, which may not be as well balanced
 - ▶ for example, a term contributing to $\delta\phi_z$ alignment is the slope of Δx residuals versus y position (there are many other terms like this)
 - ▶ the slope can be distorted by a tight cut:



- ▶ The example I drew has misalignment \sim residuals σ , which would be about 4 mrad for ϕ_z
- ▶ Monte Carlo can verify whether the MillePede alignment is sensitive to these sorts of effects

Error in the MP matrix? (1/2)

Jim Pivarski

15/40



- HIP alignment matrix, presented last month with a suite of MC tests (including turning off measurement error to observe pure geometric effects)

$$\begin{pmatrix} \Delta x \\ \Delta y \\ \Delta \frac{dx}{dz} \\ \Delta \frac{dy}{dz} \end{pmatrix} = \begin{pmatrix} -1 & 0 & \frac{dx}{dz} & y \frac{dx}{dz} & -x \frac{dx}{dz} & y \\ 0 & -1 & \frac{dy}{dz} & y \frac{dy}{dz} & -x \frac{dy}{dz} & -x \\ 0 & 0 & 0 & \frac{dx}{dz} \frac{dy}{dz} & -1 - \left(\frac{dx}{dz} \right)^2 & \frac{dy}{dz} \\ 0 & 0 & 0 & 1 + \left(\frac{dy}{dz} \right)^2 & -\frac{dx}{dz} \frac{dy}{dz} & -\frac{dx}{dz} \end{pmatrix} \begin{pmatrix} \delta_x \\ \delta_y \\ \delta_z \\ \delta_{\phi_x} \\ \delta_{\phi_y} \\ \delta_{\phi_z} \end{pmatrix}$$

- MP alignment matrix in `SegmentAlignmentDerivatives4D.cc`

$$\begin{pmatrix} \Delta x \\ \Delta y \\ \Delta \frac{dx}{dz} \\ \Delta \frac{dy}{dz} \end{pmatrix} = \begin{pmatrix} -1 & 0 & \frac{dx}{dz} & y \frac{dx}{dz} & -x \frac{dx}{dz} & y \\ 0 & -1 & \frac{dy}{dz} & y \frac{dy}{dz} & -x \frac{dy}{dz} & -x \\ 0 & 0 & \frac{dx}{dz} & 0 & -1 & \frac{dy}{dz} \\ 0 & 0 & \frac{dy}{dz} & 1 & 0 & -\frac{dx}{dz} \end{pmatrix} \begin{pmatrix} \delta_x \\ \delta_y \\ \delta_z \\ \delta_{\phi_x} \\ \delta_{\phi_y} \\ \delta_{\phi_z} \end{pmatrix}$$

- First column of differences (blue) say that δ_z misalignment causes angle residuals (e.g. $\Delta \frac{dx}{dz} = \delta_z \frac{dx}{dz}$)
- Second two columns are an approximation that $\frac{dx}{dz}$ and $\frac{dy}{dz}$ are small ($\left| \frac{dx}{dz} \right|$ reaches 0.25 in every chamber, $\left| \frac{dy}{dz} \right|$ reaches 0.7 in wheel ± 1 and 0.9 in ± 2)

Error in the MP matrix? (2/2)


Jim Pivarski

16/40



- ▶ Biggest discrepancy between HIP and MP is in local δ_z , and the biggest difference between the matrices contributes to δ_z

- ▶ δ_z translations do not change segment angles:









- ▶ A Monte Carlo test would catch this immediately


Cosmic ray Monte Carlo

- ▶ Summer08 tracker-pointing AICaReco skims available
- ▶ Everything is the same as in CRAFT data except:
 - ▶ ideal tracker alignment, magnetic field, and DT internal alignment
 - ▶ about 4 times the sample size
- ▶ All HIP/MP implementation differences should be modeled by MC
 - ▶ if not, we can add realistic tracker, field, and DTs to diagnose

- ▶ Same convergence as in data (2 iterations necessary, 3 taken)
- ▶ 200 μm in $r\phi$ (with about 4 times as many tracks)
- ▶ More accurate in y and z than collisions MC test because of high p_T
(Local z is degenerate with q -antisymmetric effects like \vec{B} and dE/dx)









- Statistical uncertainties are $3\text{--}4\times$ smaller than accuracy





- Define systematic error = $\sqrt{(\text{absolute error})^2 - (\text{statistical})^2}$
 - this “systematic” doesn’t include tracker misalignments, imperfect \vec{B} map, DT internal misalignments
 - we expect statistical to scale with \sqrt{N} , but not systematic





- ▶ Pablo and I did not get to talk about proposing constants, and I have not seen the MillePede Monte Carlo results
- ▶ What I know is:
 - ▶ the constants from the two implementations differ
 - ▶ the $\pm 1 \sigma$ cut in residuals can distort results derived from slopes
 - ▶ the error matrix in `SegmentAlignmentDerivatives4D.cc` will cause errors in z (and propagate to y for wheels ± 1)
 - ▶ the HIP procedure is well-behaved in cosmic ray Monte Carlo
 - ▶ if there are global distortions in the tracker, it would not affect HIP and MP differently
- ▶ Therefore, I will propose the HIP results, show the validation, and list all of the known mysteries

(All cuts and parameters for this alignment were listed on pages 3–4)


DB comparisons (1/5)

CRAFT_ALL_V4 – proposed constants

Jim Pivarski 21/40



- ▶ CRAFT_ALL_V4 is before global alignment
- ▶ We have seen these large absolute δ_x corrections before: they preserve local alignment within sectors (like moving fingers)



This plot now uses the latest tracker!


DB comparisons (2/5)

Jim Pivarski 22/40



CRAFT_ALL_V4 – proposed constants

- ▶ CRAFT_ALL_V4 is before global alignment
- ▶ We have seen these large absolute δ_x corrections before: they preserve local alignment within sectors (like moving fingers)



This plot now uses the latest tracker!


DB comparisons (3/5)

Jim Pivarski 23/40



CRAFT_ALL_V11 – proposed constants

- ▶ CRAFT_ALL_V11 is the first global alignment
- ▶ The wheel rotation is a known effect, related solely to difference in momentum cuts (and also preserves local alignments):
 - ▶ old: $20 < p_T < 100$ GeV, new: $100 < p_T < 200$ GeV



This plot now uses the latest tracker!


DB comparisons (4/5)

Jim Pivarski 24/40



CRAFT_ALL_V11 – proposed constants

- ▶ CRAFT_ALL_V11 is the first global alignment
- ▶ The wheel rotation is a known effect, related solely to difference in momentum cuts (and also preserves local alignments):
 - ▶ old: $20 < p_T < 100$ GeV, new: $100 < p_T < 200$ GeV









This plot now uses the latest tracker!

Rotation with high p_T (5/5)

Jim Pivarski 25/40



- These plots show the difference between low p_T and high p_T (everything else is the same; this is a systematics study)
 - global coordinates: $\Delta\phi$ (top) is rotation around beamline, $r\phi$ (bottom) is the same orientation for all chambers
- 0.35 mrad rotation, 0.04 mrad/m twist, and 3.2 mm spread





Studies of p_T effect (1/2)

Jim Pivarski 26/40



Alignments from low and high p_T differ: how do we know which is right?

- ▶ Ratio of tracker p_T and globalMuon p_T vs. momentum in CRAFT
- ▶ Study repeated with low and high p_T alignments (from prev page)
- ▶ Alignment made with high p_T (solid red and black) yields more correct ratios (1.0) at all momenta



N. Tran


Studies of p_T effect (2/2)

Jim Pivarski 27/40



Alignments from low and high p_T differ: how do we know which is right?



- ▶ Cosmic track splitting: difference between top and bottom half of cosmic ray
- ▶ Tracks with station 1 muon hits (TPFMS and TMR) yield better resolution than **tracker only (TO)** for the first time!
 - ▶ this was not the case for the low- p_T alignment, or any previous alignments
- ▶ Highest- p_T bin in this study is statistically independent of 100–200 GeV alignment




J. Tucker

p_T effect in raw residuals

Jim Pivarski 28/40




- ▶ Single chamber (wheel 0, station 1, sector 10, bottom of barrel)
- ▶ μ^+/μ^- splitting at low p_T is due to $\vec{B}(\vec{x})$ and dE/dx errors
- ▶ Charge-independent variation with p_T is not
- ▶ Not the same shape in all chambers




- ▶ Still looking at only one chamber, note that $\frac{dx}{dz}$ and p_T are related
- ▶ Expected because low- p_T muons are more vertically collimated by the Earth
- ▶ Unique to cosmic rays: in ϕ -symmetric collisions, p_T and $\frac{dx}{dz}$ will be independent

- ▶ Low- p_T band is sloped because of \vec{B}









- ▶ Residuals (greyscale, mm) are a function of both p_T and $\frac{dx}{dz}$
 - ▶ sawtooth effect is vertical trend from dark to light
 - ▶ p_T effect is horizontal darkening in center
- ▶ Sawtooth and p_T effect may be related



Verification with segments (1/2) Jim Pivarski 31/40






RPhi YB0,YB+1,YB-1




- Segment-matching among stations 1–3 yields $\sim 800 \mu\text{m}$
- Station 4 RMS is dominated by a few outliers, mostly in sector 4

A. Calderon

Verification with segments (2/2) Jim Pivarski 32/40




Phi YB0,YB+1,YB-1




► Segment angle-matching is
~0.7 mrad

Segment-extrapolation plots:


@3.8T RPhi YB1







@3.8T RPhi YB1




@3.8T RPhi YB+1



- ▶ Station 4, sector 4 is always an outlier
 - ▶ it has an apparent internal misalignment (next slide)
 - ▶ detailed fit results on next slide
- ▶ Wheel +1, station 4, sector 9 became an outlier at high p_T
 - ▶ low statistics + non-uniform coverage?
 - ▶ detailed fit results on in two slides



x residuals (mm)





dx/dz residuals (mrad)

x position (cm) y position (cm) dx/dz angle (rad) dy/dz angle (rad)





DT wheel 1, station 4, sector 4



- ▶ This is what wheel 0, station 4, sector 4 looked like with high statistics (low- p_T alignment)
- ▶ It *really* looks like a misalignment between the two halves of the chambers in this sector



DT wheel 0, station 4, sector 4




DT wheel 1, station 4, sector 9

Sawtooth effect (1/3)

Jim Pivarski 37/40



- ▶ Slope in $\Delta \frac{dx}{dz}$ versus $\frac{dx}{dz}$ which feeds into other diagnostic plots due to the correlation between Δx and $\Delta \frac{dx}{dz}$ and the correlation between x and $\frac{dx}{dz}$
- ▶ Not a rigid body misalignment of the whole chamber, but can be related to internal layer δ_x corrections




Sawtooth effect (2/3)

Jim Pivarski 38/40



- ▶ Slope in $\Delta \frac{dx}{dz}$ versus $\frac{dx}{dz}$ which feeds into other diagnostic plots due to the correlation between Δx and $\Delta \frac{dx}{dz}$ and the correlation between x and $\frac{dx}{dz}$
- ▶ Not a rigid body misalignment of the whole chamber, but can be related to internal layer δ_x corrections
- ▶ Largest in wheel -1, with a sinusoidal structure




Sawtooth effect (3/3)

Jim Pivarski 39/40



- ▶ Slope in $\Delta \frac{dx}{dz}$ versus $\frac{dx}{dz}$ which feeds into other diagnostic plots due to the correlation between Δx and $\Delta \frac{dx}{dz}$ and the correlation between x and $\frac{dx}{dz}$
- ▶ Not a rigid body misalignment of the whole chamber, but can be related to internal layer δ_x corrections
- ▶ Largest in wheel -1, with a sinusoidal structure





- ▶ Simultaneous solution of alignment and propagation effects (which are not easily separable due to convolution)
- ▶ Method verified in cosmics Monte Carlo with high accuracy ($200\ \mu\text{m}$), including geometry-only test of residuals \leftrightarrow alignment corrections matrix
- ▶ Alignment in data verified by:
 1. ratio of tracker p_T to globalmuon p_T : discovered source of rotation
 2. cosmic ray track splitting: first alignment which improves upon tracker
 3. segment extrapolation: $800\ \mu\text{m}$ in stations 1–3
- (1) Nhan Tran, (2) Jordan Tucker and Nhan Tran, (3) Alicia Calderon
- ▶ What we don't understand:
 - ▶ origin of charge-independent variation of residuals with p_T (though we know that high p_T is more correct)
 - ▶ sawtooth effect (non rigid-body variation of residuals with $\frac{dx}{dz}$)
- ▶ These conditions are external to the alignment algorithm and should affect MillePede equally
- ▶ Note that alignment will need to be repeated, with exactly the same parameters and the new tracker alignment and APEs when they become available

Timeline of updates

Jim Pivarski 41/40



All dates correspond to talks where I presented the updates

Nov 11, 2008	First CRAFT globalMuon alignment attempt: combining residuals from all chambers in each wheel, observed rotation + twist Control \vec{B} with q/p_T extrapolation
Nov 24	Used muon residuals to x-ray tracker, observed TEC misalignment
Jan 21, 2009	Split residuals by chamber, rotation + twist disappeared (low p_T) Gaussian \otimes Lorentzian convolution to model residuals tails Detailed residuals maps show discontinuities at chamber boundaries: only evidence of global structure in tracks from TEC Excluded tracks from TID/TEC, control \vec{B} with two-bin method
Jan 29	Signed-off x , y , ϕ_z chamber-by-chamber constants from 1-D fits with tails, only about half the chambers (now known as CRAFT_ALL_V11)
Jan 28 and Feb 5	First study of sawtooth: not a rigid-body misalignment (z or ϕ_y)
Feb 20	Detailed MC scenario based on CRAFT_ALL_V11 alignment
Mar 31, Apr 2, Apr 14	Updates in expanding 1-D fits into a combined 6-DOF fit Wrote 40-page track-based alignment note
Apr 27	Presented complete 6-DOF alignment with systematics studies and MC study (but low p_T), not accepted for sign-off
May 11	Discovered q -symmetric p_T dependence and origin of old rotation + twist, followed-up with resolution studies in POG
May 18	Discovered global structure in sawtooth distribution
May 25	Proposing a high- p_T HIP alignment for sign-off



CRAFT_ALL_V11 global alignment	This HIP alignment
Tight criteria for accepting an aligned chamber; only half of DTs passed	Loose criteria (5 hits and a successful fit), only 3 fail within selected region: wheels $-1, 0, +1$, all sectors except 1 and 7
x, y, ϕ_z only from independent 1-D fits	All 6 DOF from a combined fit, though station 4 is only x, ϕ_y, ϕ_z
Combining hits into segment-residuals only for their statistical properties	Also using angular information for tighter control of DOF
$20 < p_T < 100$ GeV	$100 < p_T < 200$ GeV
Control \vec{B} with two-bin method	Temporarily turn off \vec{B} control (not needed at this high momentum)
No residuals weights	Weight residuals by $(\chi^2/N_{\text{dof}})^{-1}$
2 iterations	3 iterations (third not needed)
Require 10 tracker hits, no TID/TEC	Require 15 tracker hits (for synchronization with MillePede), no TID/TEC
CMSSW_2_2_7, tracker alignment, APEs, and \vec{B} map of that time	CMSSW_2_2_11, new tracker alignment, APEs, \vec{B} map
Internal DT alignment from CRAFT_ALL_V4	New internal DT alignment (with tracks-survey agreement)
Data source: first CRAFT reprocessing	Latest CRAFT tracker-pointing reprocessing


Tracker curl hypothesis

Jim Pivarski

43/40



p_T -dependent rotation could be curl in tracker, if large enough





Tracker curl constraints

Black= MP starting object

Blue= misaligned

Red= aligned on top of misalignment

Jim Pivarski 44/40



- ▶ Studies performed in CRAFT data Zijin Guo, Roberto Castello
- ▶ Left: tracker tracks are sensitive to 300 μrad curl (blue: adding curl worsens χ^2 and red: re-aligning restores it)
- ▶ Right: also restores wafer positions within 150 μrad except TEC
 - ▶ TEC not used in muon alignment; not relevant here
 - ▶ restored chamber positions randomly distributed around zero: no *systematic* trend on the scale of 86 μrad

All the numbers

Jim Pivarski 45/40



Relative to ideal in mm and mrad

wheel station sector	CRAFT_ALL_V4 (before global alignment) $x, y, z, \phi_x, \phi_y, \phi_z$	CRAFT_ALL_V11 x, y, ϕ_z (unaligned in italics)	This HIP alignment $x, y, z, \phi_x, \phi_y, \phi_z$ (unaligned in italics)
-1, 1, 2	-5.28, -1.02, -3.54, 1.10, -1.65, -0.50,	(0.19), -1.27, (0.44),	0.36, 1.02, -4.09, 3.51, 1.06, 0.89,
-1, 1, 3	-3.14, -0.74, 2.09, 0.77, -1.17, -0.00,	1.33, -0.01, 0.48,	0.62, 2.67, 3.00, 0.60, -2.35, 0.24,
-1, 1, 4	-0.32, 0.28, 4.89, 1.98, 0.93, -0.29,	1.54, 1.94, -0.32,	1.70, 3.16, 6.14, 2.03, -0.04, -0.18,
-1, 1, 5	4.51, -0.38, 1.78, 0.29, 1.63, -0.35,	3.82, (1.78), -0.88,	5.70, 1.36, 7.08, -0.96, 2.74, -1.14,
-1, 1, 6	0.64, -1.24, -2.95, 0.50, 1.83, -0.80,	(-0.26), -1.24, (-0.80),	4.10, 0.64, -0.58, -0.82, 4.51, -0.00,
-1, 1, 8	-1.90, -0.04, -3.37, -0.95, -2.13, 0.18,	0.38, 1.39, -1.38,	1.79, 3.19, -1.58, 0.38, -1.13, -1.16,
-1, 1, 9	-2.24, -1.16, 1.84, -0.58, -1.78, 0.44,	0.27, 0.18, -1.05,	2.15, 1.01, 5.32, -0.15, -2.31, -0.93,
-1, 1, 10	-0.08, 0.36, 5.31, 1.91, 0.69, -0.02,	2.29, 0.24, 0.13,	4.94, -0.38, 7.86, 1.09, 0.11, 0.29,
-1, 1, 11	3.60, -0.46, 0.96, 1.48, 2.49, -0.11,	7.45, -2.79, 0.20,	9.82, -3.91, 7.10, -1.09, 2.06, 0.30,
-1, 1, 12	2.63, 0.45, -3.90, 1.44, 2.54, 0.02,	6.57, 0.56, 0.92,	(2.63, 0.45, -3.90, 1.44, 2.54, 0.02),
-1, 2, 2	-5.29, 2.12, -4.56, 0.22, -1.78, 0.27,	1.11, 0.91, 1.19,	1.52, 2.40, -4.81, -1.05, -4.05, 1.02,
-1, 2, 3	-4.37, -0.74, 0.77, -0.49, -1.67, -0.67,	(1.97), -0.81, (0.19),	0.80, 4.06, -0.23, -3.41, -2.30, 0.06,
-1, 2, 4	-1.63, -0.26, 3.75, -0.08, -1.36, -0.15,	1.15, 1.03, -0.16,	1.67, 2.75, 5.08, -1.21, -1.95, -0.11,
-1, 2, 5	4.03, -2.28, 1.60, 0.06, 0.61, -0.31,	(1.81), (-0.55), (0.37),	4.66, -2.73, 12.06, -3.06, 2.15, 0.22,
-1, 2, 6	3.06, -0.73, -3.87, -0.53, 1.10, 0.29,	1.40, -4.08, -1.17,	4.48, -3.19, 1.95, -3.13, 3.39, -0.25,
-1, 2, 8	-0.83, -1.06, -4.94, -0.98, -2.36, -0.65,	(0.97), -1.06, (-0.65),	(-0.83, -1.06, -4.94, -0.98, -2.36, -0.65),
-1, 2, 9	-1.99, -1.28, 0.81, -1.20, -1.92, 0.25,	(-1.99), (-1.28), (0.25),	2.45, 1.49, 5.03, -2.11, -3.18, 0.26,
-1, 2, 10	-0.15, -0.83, 4.97, -0.10, 0.10, 0.26,	1.51, (-0.83), 0.26,	5.86, -1.09, 7.61, -0.54, -0.54, 0.47,
-1, 2, 11	2.40, -0.35, 1.30, -0.67, 1.67, -0.13,	6.55, -4.02, 0.68,	11.30, -3.82, 5.98, -2.59, 1.14, -0.62,
-1, 2, 12	0.90, -0.44, -4.58, 1.13, 1.79, -0.77,	5.17, -1.26, 0.51,	10.54, -2.42, 3.88, -2.78, 0.62, 0.87,
-1, 3, 2	-4.15, -2.75, -4.37, -0.17, -1.19, -0.26,	3.60, 0.68, 1.23,	3.18, 1.50, -2.31, -0.70, -3.07, 0.76,
-1, 3, 3	-3.17, -1.33, 1.20, 0.36, -1.20, 0.03,	(-3.60), -1.33, (0.03),	2.18, 0.64, 6.04, -0.39, -2.07, -0.15,
-1, 3, 4	-0.19, -0.11, 2.82, 0.06, -0.82, 0.18,	1.99, 2.38, 0.29,	2.63, 1.72, 9.03, 1.39, -1.30, 0.34,
-1, 3, 5	3.53, -0.76, 1.61, -0.05, 1.71, -0.03,	1.05, (0.43), -0.64,	3.16, 1.18, 7.12, -1.24, 1.57, 0.80,
-1, 3, 6	7.54, -1.53, -3.92, -0.03, 1.49, 0.62,	3.79, -2.86, -0.38,	7.39, -2.56, 2.45, -2.27, 2.35, 0.35,

The last column now uses the latest tracker!

All the numbers

Jim Pivarski 46/40



Relative to ideal in mm and mrad

wheel station sector	CRAFT_ALL_V4 (before global alignment) $x, y, z, \phi_x, \phi_y, \phi_z$	CRAFT_ALL_V11 x, y, ϕ_z (unaligned in italics)	This HIP alignment $x, y, z, \phi_x, \phi_y, \phi_z$ (unaligned in italics)
-1, 3, 8	1.10, -1.21, -3.72, -0.59, -1.95, -0.54,	(1.77), -0.01, (-1.96),	3.75, 1.14, -0.54, 0.99, -3.00, -0.76,
-1, 3, 9	-1.81, -2.27, 2.31, -0.16, -1.79, 0.48,	0.47, -1.34, -0.70,	4.18, 0.28, 5.06, 0.43, -3.89, -0.67,
-1, 3, 10	0.10, 0.85, 5.31, -0.22, 0.71, 0.42,	1.43, 2.83, 0.19,	6.29, 2.08, 8.80, 0.66, -0.72, 0.74,
-1, 3, 11	1.42, 0.14, 1.81, -0.16, 2.00, 0.16,	5.17, -2.79, 0.89,	10.38, -1.99, 5.81, -1.17, 0.25, -0.86,
-1, 3, 12	0.24, -0.06, -4.60, 0.26, 2.44, -0.75,	4.52, -1.48, 1.42,	10.36, -0.77, 0.86, -0.70, 2.51, 0.09,
-1, 4, 2	-2.42, -0.81, -5.07, -0.34, -0.50, -0.54,	5.74, (-0.81), -0.54,	6.95, -0.82, -5.04, -0.34, -4.82, -1.18,
-1, 4, 3	-1.15, -0.26, -1.27, -0.40, -0.86, 0.30,	8.48, (-0.26), 0.30,	5.85, -0.26, -1.19, -0.39, -7.10, -0.30,
-1, 4, 4	1.24, 0.04, 1.88, 0.19, -0.28, 0.51,	7.19, (0.05), 0.51,	5.93, 0.05, 1.92, 0.19, -4.35, 1.13,
-1, 4, 5	4.05, -0.66, 1.04, 1.04, 0.89, -0.56,	0.70, (-0.67), -0.56,	3.77, -0.66, 1.05, 1.04, -1.85, -1.59,
-1, 4, 6	3.84, -1.47, -4.56, -0.08, 0.77, -0.76,	(3.84), (-1.47), (-0.76),	1.96, -1.47, -4.55, -0.08, -1.56, -1.09,
-1, 4, 8	-0.39, -2.54, -3.70, -1.04, -0.97, -0.58,	0.77, (-2.54), -0.58,	3.05, -2.54, -3.70, -1.03, -1.28, -0.92,
-1, 4, 9	-1.20, 0.22, 1.68, -1.05, -1.19, 0.02,	-0.21, (0.22), 0.02,	3.81, 0.22, 1.69, -1.05, 0.68, -0.35,
-1, 4, 10	-0.95, 1.05, 4.22, -0.93, 0.10, 0.03,	-2.11, (1.06), 0.03,	5.02, 1.05, 4.24, -0.93, -3.27, 0.34,
-1, 4, 11	1.92, -0.74, 2.13, 0.27, 2.02, -0.36,	4.90, (-0.73), -0.36,	11.73, -0.74, 2.15, 0.27, 6.37, -0.38,
-1, 4, 12	-2.51, -2.00, -2.32, -0.03, 1.19, 0.01,	2.58, (-1.97), 0.01,	9.10, -1.99, -2.30, -0.04, -3.27, 1.33,
-1, 4, 13	2.01, -1.05, 2.90, -0.22, 0.22, -0.31,	1.90, (-1.07), -0.31,	3.76, -1.05, 2.92, -0.22, -3.01, 0.67,
-1, 4, 14	-1.84, 1.59, 5.28, 0.19, 0.28, 0.05,	1.66, (1.59), 0.05,	6.88, 1.60, 5.29, 0.19, -2.18, 1.52,
0, 1, 2	6.69, 0.83, -2.23, -0.81, 2.41, 0.44,	(1.27), 1.23, (-0.25),	1.46, 1.74, -1.42, 1.26, 6.13, 0.49,
0, 1, 3	5.68, 1.11, 5.83, -0.28, 1.80, 0.67,	0.74, 1.53, -0.69,	1.75, 2.17, 7.08, -0.11, 3.98, -0.53,
0, 1, 4	-0.96, -2.22, 8.49, 0.39, -0.21, -0.32,	0.65, -3.82, -0.12,	1.47, -3.66, 9.71, 1.28, 2.56, -0.10,
0, 1, 5	5.69, 1.21, 5.02, 0.26, 3.00, -0.31,	2.79, 2.02, 0.02,	5.25, 0.14, 6.93, 0.69, 2.69, -0.50,
0, 1, 6	-5.34, 1.27, -4.36, -0.00, -2.92, -0.48,	-2.24, 0.26, -1.24,	-4.55, 2.40, 2.14, -0.24, -4.36, -1.27,
0, 1, 8	-3.84, 0.30, -6.65, 0.28, -3.81, -0.35,	-4.11, (-1.36), -1.05,	-2.45, -2.38, -2.70, -0.46, -3.12, -1.46,
0, 1, 9	-3.22, -0.14, 3.21, 0.08, -2.97, -0.14,	(-4.37), -0.13, (-0.14),	-0.17, -1.36, 7.21, -0.17, -3.55, -0.65,
0, 1, 10	-0.70, 0.73, 5.46, -0.18, 1.04, -0.54,	-3.18, 2.76, -0.68,	-6.04, 3.46, 11.47, 0.99, 2.30, -0.83,

The last column now uses the latest tracker!

All the numbers

Jim Pivarski 47/40



Relative to ideal in mm and mrad

wheel station sector	CRAFT_ALL_V4 (before global alignment) $x, y, z, \phi_x, \phi_y, \phi_z$	CRAFT_ALL_V11 x, y, ϕ_z (unaligned in italics)	This HIP alignment $x, y, z, \phi_x, \phi_y, \phi_z$ (unaligned in italics)
0, 1, 11	-2.79, -0.49, 3.52, -0.96, -3.73, 0.06,	-6.42, 1.35, 0.24,	-9.54, 1.98, 7.04, -0.36, -3.15, 0.10,
0, 1, 12	4.18, -0.13, -7.80, 0.83, 3.91, 0.16,	11.81, -1.33, -0.56,	12.97, -1.02, -2.73, 0.55, 3.62, -0.60,
0, 2, 2	5.63, 0.31, -4.74, -0.26, 2.21, 0.16,	1.62, 0.11, -0.09,	0.23, 0.79, -0.21, -1.01, 1.74, 0.16,
0, 2, 3	5.27, -1.41, 4.60, -0.45, 2.57, 0.41,	-0.52, -1.35, 0.13,	0.16, -0.63, 5.97, -2.03, 2.68, 1.36,
0, 2, 4	-0.21, -2.01, 8.59, 0.19, -0.97, -0.39,	2.25, (-2.00), -0.39,	3.07, -5.89, 7.10, 1.13, -3.58, -0.06,
0, 2, 5	5.57, 1.21, 5.73, 0.48, 2.51, -0.14,	2.35, 1.80, -0.24,	5.74, 0.42, 9.51, -0.34, 0.07, 0.74,
0, 2, 6	-5.24, 1.27, -1.75, -0.57, -3.03, 0.54,	-0.72, -1.25, -0.22,	-4.23, 1.29, 1.04, -2.31, -4.20, 0.46,
0, 2, 8	-2.73, 0.93, -5.64, 0.12, -3.72, 0.13,	(-3.28), -1.61, (-0.61),	-0.94, -2.68, -1.70, 0.40, -3.70, -0.67,
0, 2, 9	-2.67, 1.34, 2.36, 0.00, -3.37, -0.23,	-2.71, 0.89, -0.44,	0.74, -0.00, 8.07, 0.26, -3.78, 0.42,
0, 2, 10	0.55, 1.32, 6.66, -0.68, 0.93, -0.12,	0.72, (2.97), -0.29,	-4.40, 3.73, 14.97, -1.30, 0.33, 0.04,
0, 2, 11	-1.98, 0.06, 3.29, -0.31, -3.19, -0.01,	-3.56, 2.39, 0.13,	-9.24, 3.42, 6.89, -0.67, -4.98, 1.12,
0, 2, 12	4.28, 0.74, -6.53, 0.09, 3.66, -0.08,	11.77, -1.38, -1.00,	14.69, -0.63, 0.20, -0.14, 2.03, 0.63,
0, 3, 2	5.43, 0.59, -4.23, -0.73, 1.88, 1.13,	(-1.53), -0.30, (0.15),	-3.11, 0.34, -6.04, 2.19, 3.90, 1.11,
0, 3, 3	4.16, 0.69, 5.01, 0.10, 2.10, 0.26,	-3.66, (0.67), 0.26,	-2.90, 0.41, 4.92, 2.60, 3.65, -0.04,
0, 3, 4	0.43, 0.96, 10.25, -0.05, -0.49, -0.35,	2.37, -2.63, 0.12,	3.71, -1.54, 10.89, -0.82, -0.03, -0.47,
0, 3, 5	6.16, 1.81, 6.29, 0.45, 2.54, -0.42,	0.59, (4.62), -0.32,	4.95, 1.93, 8.42, -0.85, 4.08, -0.46,
0, 3, 6	-4.73, 1.27, -3.45, -0.25, -2.39, -0.30,	0.98, -4.13, -0.88,	-2.50, -1.13, -2.53, -0.14, -3.32, -0.37,
0, 3, 8	-3.13, 1.31, -4.89, -0.27, -2.61, 0.11,	-3.88, 0.66, -0.66,	-2.58, -1.30, 4.03, 0.33, -4.30, -0.72,
0, 3, 9	-1.69, 0.42, 3.22, 0.32, -2.59, 0.30,	-2.02, -0.09, 0.00,	1.47, -0.45, 6.59, 0.10, -2.67, 0.21,
0, 3, 10	-1.09, 1.03, 7.39, -0.19, 0.74, -0.22,	-1.34, 2.55, -0.10,	-7.25, 3.25, 14.10, 1.70, 1.18, -0.16,
0, 3, 11	-3.81, 1.16, 4.06, -0.41, -2.28, -0.16,	-5.75, (1.54), -0.38,	-12.08, 2.96, 11.54, -0.06, 1.85, -0.09,
0, 3, 12	4.24, -0.99, -5.93, 0.28, 3.19, 0.16,	12.47, -3.75, -0.98,	15.51, -3.67, -1.82, 1.69, -0.13, 0.06,
0, 4, 2	3.17, -0.48, -6.37, -0.54, 1.46, -0.20,	-3.58, (-0.48), -0.20,	-5.49, -0.48, -6.34, -0.54, 3.87, -0.29,
0, 4, 3	2.79, 1.37, 1.50, -0.17, 1.55, 0.40,	-6.40, (1.36), 0.40,	-6.17, 1.37, 1.51, -0.17, 2.08, -0.15,
0, 4, 4	2.06, 0.16, 6.59, 0.08, -0.64, -0.35,	7.95, (0.15), -0.35,	8.06, 0.16, 6.60, 0.08, -1.84, -0.67,

The last column now uses the latest tracker!

All the numbers

Jim Pivarski 48/40



Relative to ideal in mm and mrad

wheel station sector	CRAFT_ALL_V4 (before global alignment) $x, y, z, \phi_x, \phi_y, \phi_z$	CRAFT_ALL_V11 x, y, ϕ_z (unaligned in italics)	This HIP alignment $x, y, z, \phi_x, \phi_y, \phi_z$ (unaligned in italics)
0, 4, 5	5.43, 0.61, 4.83, -0.09, 1.47, -0.36,	-1.01, (<i>0.61</i>), -0.36,	3.43, 0.61, 4.85, -0.09, -1.18, -0.53,
0, 4, 6	-4.91, 1.27, -4.91, -0.70, -1.35, -0.73,	1.28, (<i>1.25</i>), -0.73,	-2.14, 1.26, -4.91, -0.70, -1.53, -0.76,
0, 4, 8	-2.13, 1.19, -9.72, -0.14, 0.94, 0.10,	(-2.13), (<i>1.19</i>), (<i>0.10</i>),	-0.38, 1.20, -9.72, -0.14, 2.43, -2.19,
0, 4, 9	-2.27, -2.96, 4.94, 0.05, -2.96, -0.04,	-1.86, (<i>-2.97</i>), -0.04,	1.15, -2.96, 4.95, 0.05, -1.92, -1.22,
0, 4, 10	-2.10, 1.21, 7.99, -0.74, 0.51, -0.23,	0.25, (<i>1.22</i>), -0.23,	-6.26, 1.22, 7.99, -0.74, -0.62, 0.73,
0, 4, 11	-3.99, 1.26, 4.84, 0.62, -2.24, -0.38,	-3.22, (<i>1.26</i>), -0.38,	-12.61, 1.26, 4.82, 0.62, -0.89, -0.46,
0, 4, 12	3.65, -0.13, -5.10, 1.31, 1.26, -0.43,	10.82, (<i>-0.14</i>), -0.43,	16.56, -0.13, -5.11, 1.31, 0.50, -1.55,
0, 4, 13	2.69, 1.15, 6.88, -0.15, 0.22, 0.08,	(<i>2.69</i>), (<i>1.15</i>), (<i>0.08</i>),	4.95, 1.16, 6.88, -0.15, 1.49, -0.62,
0, 4, 14	-1.28, -0.60, 3.34, -0.27, -4.58, -0.16,	-1.75, (<i>-0.61</i>), -0.16,	-9.28, -0.61, 3.31, -0.25, -6.94, -2.69,
1, 1, 2	5.27, -0.75, -3.42, 1.67, 1.55, -1.00,	0.55, (<i>-0.22</i>), -1.93,	-0.00, 3.50, -3.11, 2.07, 3.14, -1.72,
1, 1, 3	5.27, -2.40, 2.94, -0.22, 1.35, 0.04,	0.89, -1.66, -0.41,	1.66, 1.13, 4.84, -2.18, 2.30, -0.68,
1, 1, 4	0.01, -1.72, 5.88, 1.44, -0.57, -0.18,	-1.10, -0.66, 0.11,	-1.60, 3.84, 5.76, -1.11, -1.35, -0.15,
1, 1, 5	-3.54, -2.43, 2.43, 1.11, -2.14, 0.64,	-1.35, -4.40, 0.62,	-2.04, -0.19, 3.75, -0.10, -2.75, 0.40,
1, 1, 6	-2.93, -2.37, -3.06, 1.38, -2.00, -0.25,	(-2.93), (-2.37), (-0.25),	-0.02, -1.68, 2.56, -1.39, -4.74, -0.34,
1, 1, 8	4.27, -1.79, -3.36, 0.31, 2.01, 0.38,	5.32, -0.97, 0.93,	5.46, 0.98, 2.14, 0.63, 1.34, 0.64,
1, 1, 9	2.82, -1.67, 2.76, 0.40, 1.65, 0.01,	2.88, 0.24, -0.09,	2.96, 1.82, 8.26, -0.79, 3.74, 0.20,
1, 1, 10	0.10, -1.10, 4.61, -0.07, 0.41, 0.11,	0.00, 1.83, 0.24,	-2.38, 3.33, 10.54, -0.49, 1.55, 0.64,
1, 1, 11	-2.51, -0.45, 2.29, -0.31, -2.23, 0.02,	-6.07, 1.42, -0.48,	-7.57, 3.72, 6.30, -1.06, -0.52, -0.49,
1, 1, 12	-3.94, -1.75, -4.34, 0.32, -1.96, 0.60,	-8.19, (<i>-1.76</i>), -0.33,	-10.69, -0.59, 0.81, -0.20, -1.44, 0.83,
1, 2, 2	5.93, -2.42, -3.38, 0.16, 1.92, -0.04,	1.17, -3.03, -0.75,	(<i>5.93</i>), (-2.42), (-3.38), (<i>0.16</i>), (<i>1.92</i>), (<i>-0.04</i>),
1, 2, 3	4.80, -1.09, 2.01, -0.10, 1.62, -0.35,	0.46, -1.52, -0.63,	-0.43, 1.09, 6.90, -2.74, 1.65, 0.58,
1, 2, 4	0.24, -1.16, 3.55, 0.78, -0.45, -0.76,	-0.88, -0.70, -0.89,	-1.28, 4.62, 5.82, -2.10, 1.38, -0.55,
1, 2, 5	-2.76, -0.59, 2.97, 0.17, -1.92, 0.03,	0.46, -4.36, 0.03,	-0.76, 1.64, 1.70, -3.61, -2.71, -0.44,
1, 2, 6	-2.73, -0.33, -3.25, 0.30, -1.88, 0.07,	2.77, -1.11, -0.01,	1.13, 2.51, 0.18, -3.04, -3.64, -0.50,
1, 2, 8	4.19, 0.10, -3.94, -0.37, 2.08, 0.25,	5.98, (<i>-0.55</i>), 1.09,	5.30, 0.05, 6.72, -3.19, 3.02, -0.01,

The last column now uses the latest tracker!

All the numbers

Jim Pivarski 49/40









Relative to ideal in mm and mrad

wheel station sector	CRAFT_ALL.V4 (before global alignment) $x, y, z, \phi_x, \phi_y, \phi_z$	CRAFT_ALL.V11 x, y, ϕ_z (unaligned in italics)	This HIP alignment $x, y, z, \phi_x, \phi_y, \phi_z$ (unaligned in italics)
1, 2, 9	4.09, -1.50, 1.52, 0.26, 1.90, 0.13,	4.28, -1.17, 0.47,	3.30, -0.29, 9.90, -2.24, 2.62, 0.01,
1, 2, 10	0.38, 0.63, 3.81, -0.44, 0.21, 0.34,	2.46, 3.04, 0.36,	-1.87, 5.13, 8.95, -1.45, -0.59, 0.84,
1, 2, 11	1.96, -0.12, 1.39, -0.40, -2.13, 0.48,	-0.86, 1.19, -0.11,	-4.48, 2.45, 7.07, -1.61, -2.49, 1.02,
1, 2, 12	0.22, -1.23, -3.77, 0.35, -2.15, -0.05,	<i>(-3.05), (-1.88), (-0.79),</i>	-6.64, 1.25, -0.31, -1.35, -1.25, 0.11,
1, 3, 2	5.76, -2.34, -2.53, -0.35, 1.34, 0.57,	-1.83, -3.80, -0.66,	-1.84, -0.04, -0.15, 0.77, 3.09, 0.37,
1, 3, 3	3.41, -1.59, 1.37, -0.41, 1.57, -0.03,	-3.10, <i>(-2.31)</i> , 0.07,	-2.50, 1.03, 5.55, -1.35, 2.71, -0.43,
1, 3, 4	-0.03, 0.16, 3.85, -0.51, -0.37, 0.06,	-1.25, -2.59, -1.21,	-1.58, 4.16, 5.23, -1.90, 0.91, 0.15,
1, 3, 5	-3.07, -0.88, 2.43, 0.59, -1.01, -0.23,	1.04, -6.71, -0.26,	-0.61, 0.28, 2.86, -1.65, -2.61, -0.97,
1, 3, 6	-4.47, -0.49, -3.18, -0.16, -1.44, 0.01,	1.79, -2.24, -0.12,	0.32, 2.44, -2.08, -3.10, -2.02, -1.06,
1, 3, 8	3.06, -0.64, -4.56, -0.42, 1.95, 0.21,	<i>(2.29)</i> , -0.64, <i>(0.21)</i> ,	<i>(3.06, -0.64, -4.56, -0.42, 1.95, 0.21)</i> ,
1, 3, 9	2.66, -1.81, 2.12, 0.57, 1.59, -1.05,	1.37, <i>(-0.28)</i> , 0.23,	1.18, 1.06, 9.29, 0.61, 2.69, -0.94,
1, 3, 10	-1.64, -0.20, 4.15, -0.42, 0.58, 0.25,	0.21, 1.49, -0.29,	-4.17, 3.48, 12.63, -3.12, -1.16, -0.69,
1, 3, 11	0.17, -0.62, 1.71, -0.62, -2.08, 0.23,	-3.56, 1.00, -0.28,	-7.24, 1.95, 9.40, 0.36, -1.40, -0.02,
1, 3, 12	-1.43, -2.02, -4.96, -0.62, -2.04, 0.29,	-6.06, -3.10, -1.87,	-9.83, 1.47, -1.35, 1.88, 0.33, -0.96,
1, 4, 2	2.87, -2.72, -4.73, 0.00, 0.55, -0.80,	-5.09, <i>(-2.72)</i> , -0.80,	-7.29, -2.71, -4.73, 0.00, 0.28, 0.49,
1, 4, 3	1.74, -1.78, 0.54, 0.07, 0.89, -0.52,	-5.71, <i>(-1.78)</i> , -0.52,	-5.38, -1.78, 0.56, 0.07, 3.34, -0.57,
1, 4, 4	-0.29, 0.38, 3.63, -0.19, 0.96, 0.23,	-6.17, <i>(0.37)</i> , 0.23,	-6.34, 0.38, 3.64, -0.19, 2.44, 0.13,
1, 4, 5	-2.73, -0.13, 2.02, 0.41, -0.48, -0.26,	2.08, <i>(-0.13)</i> , -0.26,	0.26, -0.13, 2.04, 0.41, -3.77, 0.22,
1, 4, 6	-3.33, -1.17, -3.87, 0.48, -0.98, -0.54,	4.37, <i>(-1.17)</i> , -0.54,	2.59, -1.18, -3.86, 0.48, -0.89, -0.82,
1, 4, 8	0.22, -0.18, -3.92, 0.09, 0.61, -0.21,	2.92, <i>(-0.16)</i> , -0.21,	0.89, -0.18, -3.92, 0.09, 0.03, 1.51,
1, 4, 9	2.60, -1.45, 1.74, -0.30, 1.10, 0.23,	0.94, <i>(-1.43)</i> , 0.23,	0.76, -1.45, 1.74, -0.30, 2.45, -0.08,
1, 4, 10	-0.53, -1.67, 3.56, 0.64, -1.33, -0.43,	2.79, <i>(-1.67)</i> , -0.43,	-2.59, -1.67, 3.56, 0.63, -2.06, 1.78,
1, 4, 11	0.17, -0.62, 1.71, -0.62, -2.08, 0.23,	-5.77, <i>(-0.62)</i> , 0.23,	-11.93, -0.62, 1.69, -0.62, -0.51, -0.87,
1, 4, 12	-2.47, -1.36, -3.98, 0.31, -2.05, -0.78,	-6.48, <i>(-1.37)</i> , -0.78,	-10.99, -1.35, -3.98, 0.31, 1.57, -1.34,
1, 4, 13	-1.18, -0.33, 3.31, -0.00, -0.59, -0.31,	0.06, <i>(-0.32)</i> , -0.31,	-1.89, -0.33, 3.31, -0.00, -0.52, 0.78,
1, 4, 14	-0.57, -1.05, 2.15, -1.92, 1.85, -0.39,	<i>(-0.57), (-1.05), (-0.39),</i>	-5.16, -1.05, 2.18, -1.92, -1.24, -0.53,

The last column now uses the latest tracker!



- This the muon alignment produced with the CRAFT_ALL_V11 tracker alignment minus the muon alignment produced with the latest tracker, the one which will be signed-off this week.
 - Translational differences are significant and explains part of the previous discrepancy between HIP and MillePede, page 5 (MillePede had been using the right tracker)
 - Track-fitting efficiency with the new tracker is 80% of track-fitting efficiency with the old tracker (note that pattern recognition used the old tracker, so efficiency cannot be better with the new one)





- ▶ This the muon alignment produced with the CRAFT_ALL_V11 tracker alignment minus the muon alignment produced with the latest tracker, the one which will be signed-off this week.
 - ▶ Translational differences are significant and explains part of the previous discrepancy between HIP and MillePede, page 5 (MillePede had been using the right tracker)
 - ▶ Track-fitting efficiency with the new tracker is 80% of track-fitting efficiency with the old tracker (note that pattern recognition used the old tracker, so efficiency cannot be better with the new one)

