



HW-track Comparison and Tracker Global Shape



Jim Pivarski

Texas A&M University

4 December, 2009



- ▶ Hardware alignment as seen by tracks
- ▶ Resolving tracker global distortions with muon residuals



► Nov 19 Link-fixed HW barrel alignment

► $\phi = x/R$







► Much more consistent with a single offset of the tracker

► $x \rightarrow 1.2$ mm

► $y \rightarrow 4.5$ mm

► $\phi_z \rightarrow 0.58$ mrad



► See <https://hypernews.cern.ch/HyperNews/CMS/get/muon-alignment/423/1.html> or Nov 20 Indico page for the full set of plots



- RMS of x' deviations (top-left) is only 1.8 mm after removing global offset in transverse plane
- Similar to photogrammetry, which had an RMS of 1.6 mm


Other translational d.o.f.

Jim Pivarski 5/14



- ▶ Top: y' differences (direction parallel to beamline)
- ▶ Bottom: z' differences (radial)
- ▶ These are more consistent within sector groups
- ▶ y' has a clear trend with respect to wheel


wheel +0, station 1



- ▶ Map plots show that the remaining differences are not due to tracker distortions
- ▶ For example, the statement “relative misalignment of sectors 4 and 5 is 6 mm” is tracker-independent and propagation-independent
- ▶ When the hardware geometry correctly describes the muon system as a rigid body, the differences with respect to track-based will be a *smooth* function in these plots (e.g. between sectors 11 and 12)



- ▶ We can identify tracker global distortions using muon chamber data
- ▶ Without introducing circularity when we later align the muon system with tracker tracks




- ▶ Independence from muon alignment: plot residuals in *only one muon layer* (wh 0, st 1, sec 10, lay 2) and disregard global position of that layer
- ▶ Look at muon residuals as a function of p_T
- ▶ We only assume that the muon layer is in one location that can't be a function of the p_T of the tracks used to measure it

p_T -dependent muon residuals


Jim Pivarski 8/14




CRAFT-09 aligned with $p_T > 100$ GeV tracks



CRAFT-09 aligned with $p_T > 100$ GeV tracks



CRAFT-09 aligned with $p_T > 100$ GeV tracks



- ▶ Three ways of looking at it:
 - ▶ as a muon residual (Δx)
 - ▶ tracker curvature error ($\Delta\kappa = \Delta x \frac{d\kappa}{dx}$, $\kappa = q/p_T$)
 - ▶ tracker momentum error ($\Delta p_T = \Delta x \frac{dp_T}{dx}$)

where $\frac{d\kappa}{dx}$ and $\frac{dp_T}{dx}$ come from track propagator, numerically


- ▶ $\frac{d\kappa}{dx}$ is nearly constant for a single DT layer (depends on distance from tracker)
- ▶ From muon hits, we learn something which is purely about the tracker's shape
- ▶ If errors were from $\vec{B}(\vec{x})$ or dE/dx , top plot would be antisymmetric, not symmetric

Ambiguity from muon alignment


Jim Pivarski 9/14




CRAFT-09 aligned with low- p_T tracks



CRAFT-09 aligned with low- p_T tracks



CRAFT-09 aligned with low- p_T tracks




- ▶ To avoid a circular argument, we should keep the position of the muon chamber as a free parameter in this study
- ▶ Knowledge of tracker shape will later be used to determine positions of muon chambers (track-based alignment)
- ▶ Freedom to make low- p_T region “correct” and high- p_T region “wrong”
- ▶ Still, difference in curvature between low- and high- p_T regions is the same: this should constrain models of the tracker’s shape

Does this analysis make sense?


Jim Pivarski 10/14




ideal Monte Carlo



ideal Monte Carlo



ideal Monte Carlo




- ▶ When applied to ideal MC, everything is perfect
- ▶ That's good (not a software problem or anything)


Straw-man global distortions (1) Jim Pivarski 11/14






Monte Carlo with layerRotation in the tracker (ruled out by track χ^2)



Monte Carlo with layerRotation in the tracker (ruled out by track χ^2)



Monte Carlo with layerRotation in the tracker (ruled out by track χ^2)




- ▶ $r\phi$ rotation of tracker layers as a function of r
- ▶ Tracker track χ^2 is highly sensitive to this, so it has been ruled out
- ▶ Nevertheless, it would produce a similar effect


Straw-man global distortions (2) Jim Pivarski 12/14






Monte Carlo with sagitta in the tracker (plausible)



Monte Carlo with sagitta in the tracker (plausible)



Monte Carlo with sagitta in the tracker (plausible)




- $r\phi$ rotation of tracker layers as a function of ϕ
- Cosmic ray tracker tracks are not very sensitive to this
- It also produces a similar effect
- That doesn't mean that it's the only explanation

Why it's not sagitta




- ▶ Plot tracker curvature error $\Delta(q/p_T)$ on the color scale (GeV^{-1})
- ▶ Horizontal axes as indicated

CRAFT-09 aligned with $p_T > 100$ GeV tracks



Monte Carlo with sagitta in the tracker (plausible)



- ▶ Real distortion is more a function of q/p_T than ϕ
- ▶ Sagitta error is more a function of ϕ than q/p_T



- ▶ New hardware geometry is much more internally consistent
- ▶ There are still observable discrepancies that are not due to tracker global distortions or track propagation
- ▶ Muon residuals can be used to identify (and eventually constrain) tracker global distortions
 - ▶ turning the problem discovered in May into an asset
 - ▶ I'm trying to get people interested in using this method as a tool for tracker diagnosis: two people are possibly interested