



Update on Dimuon Resonances in “Lepton Jets”


Jim Pivarski

Alexei Safonov

Aysen Tatarinov

Texas A&M University

21 January, 2011



Analysis strategy (reminder)


Motivated by phenomenology:

- ▶ Hidden sector's spectrum is unknown, but weak coupling to the Standard Model implies that it predominantly passes through the lightest hidden state
→ search for low-mass dimuons
- ▶ Muon pairs may overlap, but cascades of light particles would be collimated in groups by their boost
→ first identify well-separated groups, then resolve combinatorics
- ▶ $\mathcal{B}(m_1 \rightarrow \mu\mu)$ is likely to be high, but not necessarily 100%
→ look for muons, but neither require nor exclude other particles (e.g. by applying an isolation cut)

Jim Pivarski 2/22






mass scale






- ▶ Unless the hidden spectrum has a very close degeneracy, the lightest state (m_1) will be on-shell
→ search for a resonance peak
- ▶ Groups of four or more muons represent $m_2 \rightarrow m_1 m_1$ cascades
→ split them into the most consistent assignment of dimuon masses (assumes on-shell m_1)
- ▶ Different event topologies have different backgrounds
→ partition signal region by number of collimated groups (“mu-jets”) and number of dimuons within each mu-jet





- ▶ Look for new resonance in all channels that have small backgrounds
- ▶ Any one of the following reduce Standard Model backgrounds to $\mathcal{O}(1 \text{ pb})$:

- ▶ $p_T \gtrsim 80 \text{ GeV}/c$
- ▶ ≥ 2 mu-jets in an event
- ▶ ≥ 4 muons in a mu-jet



- ▶ Define non-overlapping signal regions:

(a) exactly one mu-jet per event

- (a-1) $p_T > 80 \text{ GeV}/c$ mu-jet containing two muons ($m_1 \rightarrow 2\mu$)
- (a-2) any mu-jet containing four muons ($m_2 \rightarrow m_1 m_1 \rightarrow 4\mu$)
- (a-3) more than four



(b) two mu-jets per event

- (b-1) $2\mu, 2\mu$ ($M \rightarrow m_1 m_1$, which is the NMSSM signature)
- (b-2) $2\mu, 4\mu$ ($M \rightarrow m_1 m_2$)
- (b-3) $4\mu, 4\mu$ ($M \rightarrow m_2 m_2$)
- (b-4) either has more than four

(c) more than two mu-jets per event



- ▶ Unlike jets, our signals have a well-defined but unknown mass
 - ▶ all on-shell m_1 particles in an event have the same mass
 - ▶ backgrounds fill the space of dimuon masses more uniformly
- ▶ Measure signal and backgrounds in a single “fit with sidebands”
 - ▶ topologies with n dimuons per event form an n -dimensional space of observables
 - ▶ signal is a sharp peak somewhere along the diagonal
 - ▶ background distribution is a Cartesian product of shapes derived from data





- ▶ Opening the box: results unblinded last weekend
- ▶ Signal and background shape templates
- ▶ Implications for benchmark models






Opening the Box

Results: all consistent with SM

Jim Pivarski 8/22



- 137 events with a single, high- p_T dimuon (SM-like distribution)
- 1 event with a 4- μ mu-jet
- 11 events with two 2- μ mu-jets
- 0 events within 5σ (detector resolution) of a 2-D diagonal
- 0 events with 3 or more dimuons






Event displays of double-dimuons (which have unequal masses)

(a-2) two dimuons in one mu-jet

CMS Experiment at LHC, CERN
Data recorded: Mon Oct 11 16:03:58 2010 CDT
Run/Event: 147754 / 142156381
Lumi section: 115
Orbit/Crossing: 30005881 / 1255




(b-1) two well-separated dimuons

CMS Experiment at LHC, CERN
Data recorded: Thu Oct 14 00:41:01 2010 CDT
Run/Event: 147927 / 135595064
Lumi section: 128
Orbit/Crossing: 33553173 / 1449

mass = 1.46 GeV/c²


MET = 26.6 GeV






- ▶ Limit-setting in RooFit/RooStats using the MCMC method
- ▶ Luminosity, efficiency, and background shape systematics folded in (to be described later)
- ▶ Set limits on $\sigma \times \mathcal{B} \times \alpha_{\text{gen}}$, where α_{gen} is the acceptance for a given model in a given signal region
 - ▶ signature-based limit, applicable to future theories


(a-1) one high- p_T
dimuon



(a-2) two dimuons in
one mu-jet



(b-1) two well-separated
dimuons



All other signal regions (3 or more dimuons per event) are also excluded at the level of 0.1 pb, independent of m_1 mass $< 5 \text{ GeV}/c^2$







Fit Shape Templates

Signal shape (the easy part)

Jim Pivarski 12/22



- ▶ Since the hidden sector couples weakly to the Standard Model, the m_1 resonance width must be dominated by detector resolution
- ▶ We have four Standard Model resonances in our background-enriched dataset (single dimuon, $p_T < 80$ GeV/c)




Signal shape (the easy part)


Jim Pivarski 13/22



Barrel ($|\eta| < 0.9$)






Endcap ($|\eta| > 0.9$)




- Different resolution in the barrel and endcap
- Filled in resolution vs. mass and p_T with MC pair-gun
- Data agree well with the minimum- p_T curve
- Modeled in final fit as double-Gaussian (barrel and endcap) with a Crystal Ball radiative tail, p_T variation is a systematic uncertainty

- ▶ Different physics sources contribute to each signal region, so the background shape templates must be individually constructed
- ▶ For (a-1): prompt & isolated, $b\bar{b}$ -like, and J/ψ components all scale as $\exp(-p_T/10 \text{ GeV})$ above 40 GeV/c
- ▶ Derive shape from high-statistics low- p_T data, test in medium- p_T , and use in high- p_T signal search



- ▶ Region (a-2): 4 muons in one mu-jet
- ▶ Dominant Standard Model backgrounds: decays-in-flight and misreconstruction (fakes)
- ▶ Simulate fake muons by putting non-muon tracks into mu-jets:

CMS Experiment at LHC, CERN
Data recorded: Mon Oct 11 10:05:08 2010 CDT
Run/Event: 187704 / 140168381
Lumi section: 115
CMSSW version: 30068801 / 1255



Background-enriched

Control



Signal Region

2 muons, 2 tracks

3 muons, 1 track

4 muons

- ▶ Plots of control region with template shape overlaid:






Background shape

Jim Pivarski 16/22



- ▶ Region (b-1): 2 mu-jets with 2 muons each
- ▶ Dominant Standard Model backgrounds: $b\bar{b}$ with both b -quarks producing $\mu\mu X$ by double-semileptonic decay, resonances, etc.
- ▶ Assume that each b -quark decays independently and construct 2-D distribution from Cartesian product of 1-D $b \rightarrow \mu\mu X$ distributions
- ▶ Background-enriched: selected dimuons (left);
control region: off-diagonal part of signal (right; already seen)






Implications for Benchmark Models



SUSY dark matter + extra $\mathcal{U}(1)_{\text{dark}}$
(inspired by PAMELA)

$$z_d \rightarrow 2\mu \text{ and } h_d \rightarrow z_d z_d \rightarrow 4\mu$$


$$m_{z_d} = 1 \text{ GeV}/c^2, m_{h_d} = 3 \text{ GeV}/c^2$$



NMSSM Higgs
(inspired by hidden Higgs)

$$h \rightarrow aa \rightarrow 2\mu, 2\mu$$

$$m_h = 100 \text{ GeV}/c^2, m_a = 2 \text{ GeV}/c^2$$



- ▶ Extra- $\mathcal{U}(1)$ model produces complicated events that have high acceptance but fall into many signal regions
- ▶ Because NMSSM is a “heavy \rightarrow light” model, it produces 2 well-separated mu-jets, each with exactly 2 muons: (b-1)



- Below is the Dark SUSY model used in the Princeton studies





- we're using the same MC samples (both types, $\gamma_{\text{dark}} \rightarrow 2\mu$ and $h_{\text{dark}} \rightarrow 4\mu$, with 11 $m_{\tilde{q}}$ mass points; three shown below)

$m_{\tilde{q}} = 200 \text{ GeV}/c^2$





$m_{\tilde{q}} = 400 \text{ GeV}/c^2$

$m_{\tilde{q}} = 600 \text{ GeV}/c^2$

$\tilde{n}_2 \rightarrow \tilde{n}_1 \gamma_{\text{dark}} \rightarrow 2\mu$




$\tilde{n}_2 \rightarrow \tilde{n}_1 h_{\text{dark}} \rightarrow 4\mu$






- ▶ $\mathcal{B}(\gamma_{\text{dark}} \rightarrow \mu\mu) = 100\%$ in the Dark SUSY MC samples
- ▶ Varying the branching fraction changes the topologies of events (because there are multiple γ_{dark} per event)
- ▶ We varied the effective branching fractions by dropping muon pairs


$\mathcal{B}(\gamma_{\text{dark}} \rightarrow \mu\mu) = 100\%$



$\mathcal{B}(\gamma_{\text{dark}} \rightarrow \mu\mu) = 50\%$



$\mathcal{B}(\gamma_{\text{dark}} \rightarrow \mu\mu) = 33\%$




- ▶ Transitions from multimuons to mostly single-dimuon (a-1)






- Produced by combining the model-independent limits on each signal region (page 10) with the appropriate model acceptances

$$\tilde{n}_2 \rightarrow \tilde{n}_1 \gamma_{\text{dark}} \rightarrow 2\mu$$



$$\tilde{n}_2 \rightarrow \tilde{n}_1 h_{\text{dark}} \rightarrow \gamma_{\text{dark}} \gamma_{\text{dark}} \rightarrow 4\mu$$



Conclusions

Jim Pivarski

22/22



Switch to Alexei's slides




Backup 1: Detector Issues and Acceptance




- ▶ Some triggers' efficiencies depend strongly on whether the muon trajectories cross in the muon system
 - ▶ this is a problem for a model-independent study because different models have different fractions of muon crossing
 - ▶ parameterizing it would make the results too complicated
- ▶ Trigger efficiency vs. η for crossing and **non-crossing** muons:


single-muon (Mu15)



isolated (IsoMu9)





double-muon (DMu3)




- ▶ Only the single-muon barrel trigger is highly efficient for nearby muons, regardless of crossing, so we define acceptance:
each event must have at least one $p_T > 15 \text{ GeV}/c$, $|\eta| < 0.9$ muon



- ▶ GlobalMuons are also inefficient for crossing muons ($\varepsilon \sim 50\%$)
- ▶ TrackerMuons are much less sensitive to crossing, so all muons in the analysis must be:
 - ▶ TrackerMuons with at least 2 arbitrated segments,
 - ▶ $p_T > 5 \text{ GeV}/c$, $|\eta| < 2.4$,
 - ▶ track normalized $\chi^2 < 4$, at least 8 hits.
- ▶ Quality TrackerMuon efficiency for crossing and **non-crossing** muons:




Backup 2: Understanding the Low-Mass Dimuon Spectrum





- ▶ “Raw” mass spectrum has several components:



- ▶ resonances (prompt and from b decays)
- ▶ double-semileptonic $b \rightarrow \mu\mu X$ continuum
- ▶ low-mass Drell-Yan



- ▶ MC isn't perfect: study data/MC differences using isolation (defined such that $\mu\mu$ doesn't self-veto) and distance of flight (L_{xy})




- Split sample into $b\bar{b}$ and Drell-Yan/prompt resonances with this cut:
 $l_{so} > 4.5 \text{ GeV}/c$ **or** $L_{xy} > 2 \text{ mm}$



- Much of the low-mass spectrum is Drell-Yan (not in any official samples, so we generated it with Pythia 8)
- Some resonances are not in inclusive-muon MC: $\omega(782)$, $\psi'(3686)$
- There's also a low-mass excess (red circle), too wide to be a resonance peak, and too low in mass to be $\eta \rightarrow \mu\mu\gamma$



- ▶ Part of the explanation: the cut $l_{so} > 4.5 \text{ GeV}/c$ **or** $L_{xy} > 2 \text{ mm}$ depends on L_{xy} , which becomes imprecise for nearly collinear tracks: Drell-Yan leaks into the “ $b\bar{b}$ ” sample



- ▶ This mass region is below the Pythia generator-level mass cut-off. . .
- ▶ About $1/10^{\text{th}}$ of what remains passes $l_{so} > 4.5 \text{ GeV}/c$ alone: misreconstructed Drell-Yan?
- ▶ History of this study is a geometric progression of explaining 90%, then explaining 90% of what's left, etc..