

Introduction
oooo

Prediction and uncertainty
oooooooo

Case study
oooooo

Drug discovery
oooo

Final thoughts
oo


Solving material design problems with simulation and ML

Jon Paul Janet

01.01.0001

Machine learning and simulation for chemical discovery

How can we design new materials using computers?




The space of possible chemistry is incredibly vast, with $\mathcal{O}(10^{60})$ small organic molecules alone.

All potential medicines, catalysts and materials are somewhere, out in this huge space.

Machine learning and simulation for chemical discovery

How can we design new materials using computers?



The space of possible chemistry is incredibly vast, with $\mathcal{O}(10^{60})$ small organic molecules alone.

All potential medicines, catalysts and materials are somewhere, out in this huge space.

Machine learning and simulation for chemical discovery

How can we design new materials using computers?



The space of possible chemistry is incredibly vast, with $\mathcal{O}(10^{60})$ small organic molecules alone.




All potential medicines, catalysts and materials are somewhere, out in this huge space.




Machine learning and simulation for chemical discovery

How can we design new materials using computers?



Machine learning and simulation for chemical discovery


Inorganic materials design




Machine learning and simulation for chemical discovery

Inorganic materials design

few training points




Machine learning and simulation for chemical discovery



Inorganic materials design

- surrogate modeling & integration with simulation


Machine learning and simulation for chemical discovery



Inorganic materials design

- surrogate modeling & integration with simulation


Machine learning and simulation for chemical discovery



Inorganic materials design

- surrogate modeling & integration with simulation


Machine learning and simulation for chemical discovery



Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification


Machine learning and simulation for chemical discovery



Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification
- Bayesian design

Machine learning and simulation for chemical discovery




Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification
- Bayesian design

Drug discovery

Machine learning and simulation for chemical discovery

Alternative: learn a generative distribution over chemical space




Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification
- Bayesian design


Drug discovery

- reinforcement & generative learning


Machine learning and simulation for chemical discovery



Machine learning and simulation for chemical discovery



Machine learning and simulation for chemical discovery




Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification
- Bayesian design

Drug discovery

- reinforcement & generative learning
- 3D/protein conditioning
- curriculum & transfer learning

Machine learning and simulation for chemical discovery




Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification
- Bayesian design

Drug discovery

- reinforcement & generative learning
- 3D/protein conditioning
- curriculum & transfer learning

Machine learning and simulation for chemical discovery



Inorganic materials design

- surrogate modeling & integration with simulation
- uncertainty quantification
- Bayesian design

Drug discovery


- reinforcement & generative learning
- 3D/protein conditioning
- curriculum & transfer learning

Why transition metal complexes?

Technology requires ever greater control of material properties.

Why transition metal complexes?


Technology requires ever greater control of material properties.



Transition metal complexes offer precisely **tunable electronic properties**

Why transition metal complexes?


Technology requires ever greater control of material properties.



Transition metal complexes offer precisely **tunable electronic properties**


Why transition metal complexes?

Technology requires ever greater control of material properties.



molecular devices


Aravena, D. et al., *Inorg. Chem.*, 55(13):6405–6413, 2016.



Transition metal complexes offer precisely **tunable electronic properties**


Why transition metal complexes?

Technology requires ever greater control of material properties.




Rohde, J. U. et al., *Science*,
299(5609):1037–1039, 2003.

catalysis



molecular devices


Aravena, D. et al., *Inorg. Chem.*, 55(13):6405–6413, 2016.



Transition metal complexes offer precisely **tunable electronic properties**

Why transition metal complexes?

Technology requires ever greater control of material properties.




Rohde, J. U. et al., *Science*,
299(5609):1037–1039, 2003.

catalysis




energy capture

Michael Grätzel, photo credit
Alain Herzog/EPFL



molecular devices


Aravena, D. et al., *Inorg. Chem.*, 55(13):6405–6413, 2016.



Transition metal complexes offer precisely **tunable electronic properties**


Why transition metal complexes?

Technology requires ever greater control of material properties.




Rohde, J. U. et al., *Science*,
299(5609):1037–1039, 2003.

catalysis




energy capture

Michael Grätzel, photo credit
Alain Herzog/EPFL



molecular devices


Aravena, D. et al., *Inorg. Chem.*, 55(13):6405–6413, 2016.



Transition metal complexes offer precisely **tunable electronic properties**


Why transition metal complexes?

Technology requires ever greater control of material properties.




Rohde, J. U. et al., *Science*,
299(5609):1037–1039, 2003.

catalysis




energy capture

Michael Grätzel, photo credit
Alain Herzog/EPFL




[commons.wikimedia.org/
Redox_Flow_Battery](https://commons.wikimedia.org/wiki/File:Redox_Flow_Battery_diagram.svg)
energy storage



molecular devices


Aravena, D. et al., *Inorg.
Chem.*, 55(13):6405–6413, 2016.



Transition metal complexes offer precisely **tunable electronic properties**


Why transition metal complexes?

Technology requires ever greater control of material properties.




Rohde, J. U. et al., *Science*,
299(5609):1037–1039, 2003.

catalysis




energy capture

Michael Grätzel, photo credit
Alain Herzog/EPFL




[commons.wikimedia.org/
Redox_Flow_Battery](https://commons.wikimedia.org/wiki/File:Redox_Flow_Battery_diagram.jpg)

energy storage



molecular devices


Aravena, D. et al., *Inorg.
Chem.*, 55(13):6405–6413, 2016.





Transition metal complexes offer precisely **tunable electronic properties**

Tunable electronic structure of transition metal complexes


Tunable electronic structure of transition metal complexes





Tunable electronic structure of transition metal complexes





Tunable electronic structure of transition metal complexes



Tunable electronic structure of transition metal complexes





Tunable electronic structure of transition metal complexes





Kulik, H.J., *WIREs Comput Mol Sci.*, e1439, 2019





Tunable electronic structure of transition metal complexes





Tunable electronic structure of transition metal complexes



Tunable electronic structure of transition metal complexes






Tunable electronic structure of transition metal complexes






Seredyuk, M et al., *Chem. Mater.*, 18(10):2513–2519, 2006.




Tunable electronic structure of transition metal complexes



Tunable electronic structure of transition metal complexes



Tunable electronic structure of transition metal complexes



Overview: algorithmic surrogate-drive chemical optimization


design
space

Overview: algorithmic surrogate-drive chemical optimization


design
space

candidate
materials


Overview: algorithmic surrogate-drive chemical optimization




Overview: algorithmic surrogate-drive chemical optimization




Overview: algorithmic surrogate-drive chemical optimization




Overview: algorithmic surrogate-drive chemical optimization



Overview: algorithmic surrogate-drive chemical optimization




Overview: algorithmic surrogate-drive chemical optimization



Featurization with RACs


Graph-based features designed for TM complexes:



Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs


Graph-based features designed for TM complexes:



Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs


Graph-based features designed for TM complexes:



Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs


Graph-based features designed for TM complexes:



Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs


Graph-based features designed for TM complexes:



Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs


Graph-based features designed for TM complexes:




Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs


Graph-based features designed for TM complexes:



Coulomb Matrix




RACs




Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs



Graph-based features designed for TM complexes:



Coulomb Matrix




RACs




leads to $\sim 20\times$ more accurate property prediction without 3D geometric information

Featurization with RACs



Graph-based features designed for TM complexes:



Coulomb Matrix




RACs




leads to $\sim 20\times$ more accurate property prediction without 3D geometric information

Featurization with RACs

Graph-based features designed for TM complexes:



spin splitting ΔE_{H-L}




redox ΔG_{solv}

Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs

Graph-based features designed for TM complexes:

spin splitting ΔE_{H-L}




redox ΔG_{solv}


Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.

Featurization with RACs

Graph-based features designed for TM complexes:



spin splitting ΔE_{H-L}





Janet, J.P., and Kulik, H.J., *J. Phys. Chem. A*, 121(46):8939–8954, 2017.


Janet, J.P. et al., *Inorg. Chem.*, 58(16):10592–10606, 2019.

Featurization with RACs

Graph-based features designed for TM complexes:



spin splitting ΔE_{H-L}




redox ΔG_{solv}

Property inference with ML

We can accurately estimate DFT properties using simple neural networks


Property inference with ML

We can accurately estimate DFT properties using simple neural networks





Property inference with ML

We can accurately estimate DFT properties using simple neural networks




Property inference with ML

We can accurately estimate DFT properties using simple neural networks




Property inference with ML

We can accurately estimate DFT properties using simple neural networks




spin splitting energies

- negligible cost relative to DFT (20 GPU-hours)
- errors of 1–3 kcal/mol on similar unseen cases




Property inference with ML

We can accurately estimate DFT properties using simple neural networks




frontier orbital properties

- negligible cost relative to DFT (20 GPU-hours)
- errors of 0.2 eV on similar unseen cases




Property inference with ML


We can accurately estimate DFT properties using simple neural networks



DFT equilibrium bond lengths



- negligible cost relative to DFT (20 GPU-hours)
- errors of 0.1 pm on similar unseen cases



Janet, J.P. et al., *Inorg. Chem.*, 58(16):10592–10606, 2019.

Janet, J.P. et al., *Ind. Eng. Chem. Res.*, 56(17):4898–4910, 2017.

Janet, J.P. and Kulik, H.J., *Chem. Sci.*, 8:5137–5152, 2017.

Nandy, A. et al., *Ind. Eng. Chem. Res.*, 57(42):13973–13986, 2018.


Beyond prediction: live job management

In high-throughput DFT screening, job failure is a frequent issue

Beyond prediction: live job management


In high-throughput DFT screening, job failure is a frequent issue

new complex




Beyond prediction: live job management

In high-throughput DFT screening, job failure is a frequent issue




Beyond prediction: live job management

In high-throughput DFT screening, job failure is a frequent issue



Beyond prediction: live job management


In high-throughput DFT screening, job failure is a frequent issue




Beyond prediction: live job management

new complex


Beyond prediction: live job management




Beyond prediction: live job management




Beyond prediction: live job management




Beyond prediction: live job management




Beyond prediction: live job management



Beyond prediction: live job management




Beyond prediction: live job management



Beyond prediction: live job management

This approach leads to **30% time savings** and can abort almost all failures.



How far can we extrapolate?


Out-of-distribution test:

Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.

How far can we extrapolate?

Out-of-distribution test:


Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.



How far can we extrapolate?

Out-of-distribution test:


Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.



How far can we extrapolate?

Out-of-distribution test:


Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.



How far can we extrapolate?

Out-of-distribution test:

Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.




How far can we extrapolate?

Out-of-distribution test:

Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.

Test generalization performance of MLP trained on 1900 DFT results for complexes with common small ligands.




How far can we extrapolate?

Out-of-distribution test:

Computed spin-splitting energies of 116 diverse structures from the CSD, from training-like to very different.

Test generalization performance of MLP trained on 1900 DFT results for complexes with common small ligands.



Introduction
oooo

Prediction and uncertainty
oooo●ooo


Case study
oooooooo

Drug discovery
oooo


Final thoughts
oo

Latent distance similarity


Latent distance similarity




Latent distance similarity




Latent distance similarity




Latent distance similarity




Latent distance similarity




Latent distance similarity




Latent distance similarity




Latent distance similarity



Latent distance similarity



Latent distance similarity



How to measure chemical extrapolation?


How to measure chemical extrapolation?

How to measure chemical extrapolation?

latent distances are superior to feature space distances



How to measure chemical extrapolation?

latent distances are superior to feature space distances



How to measure chemical extrapolation?

latent distances are superior to feature space distances





competitive with ensembles and dropout

Janet, J.P., et al., *Chem. Sci.*, 10:7913–7922, 2019.

How to measure chemical extrapolation?

latent distances are superior to feature space distances





competitive with ensembles and dropout

Janet, J.P., et al., *Chem. Sci.*, 10:7913–7922, 2019.

How to measure chemical extrapolation?

latent distances are superior to feature space distances



Wang, D., et al., *J. Cheminform.*, 13:69, 2021

Hirschfeld, L., et al., *J. Chem. Inf. Model.*, 60(8):3770–3780, 2020

competitive with ensembles and dropout and works well for active learning

Janet, J.P., et al., *Chem. Sci.*, 10:7913–7922, 2019.


From distances to error distributions

We need to provide confidence estimates in appropriate units:

From distances to error distributions

We need to provide confidence estimates in appropriate units:


$$\varepsilon(d) \sim \mathcal{N}(0, \sigma_1^2 + d\sigma_2^2)$$



From distances to error distributions


We need to provide confidence estimates in appropriate units:

$$\varepsilon(d) \sim \mathcal{N}(0, \sigma_1^2 + d\sigma_2^2)$$



Redox flow batteries

Redox flow batteries (RFBs) are a promising option for scalable energy storage:




Perry, M.L. and Adam, Z., *J. Electrochem. Soc.*, 163(1):A5064–A5067, 2018.

Redox flow batteries

Redox flow batteries (RFBs) are a promising option for scalable energy storage:

Transition metal complexes make attractive redox couples for RFBs




Perry, M.L. and Adam, Z., *J. Electrochem. Soc.*, 163(1):A5064–A5067, 2018.

Redox flow batteries

Redox flow batteries (RFBs) are a promising option for scalable energy storage:

Transition metal complexes make attractive redox couples for RFBs


- good ion stability (compared to organics)



Perry, M.L. and Adam, Z., *J. Electrochem. Soc.*, 163(1):A5064–A5067, 2018.

Redox flow batteries

Redox flow batteries (RFBs) are a promising option for scalable energy storage:




Transition metal complexes make attractive redox couples for RFBs

- good ion stability (compared to organics)
- good range of redox potentials available

Perry, M.L. and Adam, Z., *J. Electrochem. Soc.*, 163(1):A5064–A5067, 2018.

Redox flow batteries

Redox flow batteries (RFBs) are a promising option for scalable energy storage:



Transition metal complexes make attractive redox couples for RFBs


- good ion stability (compared to organics)
- good range of redox potentials available
- **solubility is an issue!**

$$E_{\text{cell}} = 0.5 \times \Delta G_{\text{solv}} \times C \times n \times F$$

Perry, M.L. and Adam, Z., *J. Electrochem. Soc.*, 163(1):A5064–A5067, 2018.

Redox flow batteries

Redox flow batteries (RFBs) are a promising option for scalable energy storage:



Transition metal complexes make attractive redox couples for RFBs

- good ion stability (compared to organics)
- good range of redox potentials available
- **solubility is an issue!**

$$E_{\text{cell}} = 0.5 \times \Delta G_{\text{solv}} \times C \times n \times F$$

Perry, M.L. and Adam, Z., *J. Electrochem. Soc.*, 163(1):A5064–A5067, 2018.


We need complexes that have high redox potential **and** good solubility

Multiobjective optimization

We can construct a large design space of feasible candidates and screen them with NNs

Multiobjective optimization

We can construct a large design space of feasible candidates and screen them with NNs






Combinatorial library of 3M RFBs

Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Multiobjective optimization

We can construct a large design space of feasible candidates and screen them with NNs





Combinatorial library of $3M$ RFBs screened in < 4 mins on a regular workstation, c.f. 50 GPU-years with DFT

Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.



Multiobjective optimization

We can construct a large design space of feasible candidates and screen them with NNs



Multiobjective optimization

We can construct a large design space of feasible candidates and screen them with NNs




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

$$\Delta G_{\text{solv}} = \begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \hat{\mu}_1 \\ \hat{\mu}_2 \end{bmatrix}, \begin{bmatrix} \hat{\sigma}_1^2 & 0 \\ 0 & \hat{\sigma}_2^2 \end{bmatrix} \right)$$

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.


2D EGO Illustration

We will use a multiobjective expected improvement framework:



2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration


We will use a multiobjective expected improvement framework:



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration

We will use a multiobjective expected improvement framework:




$$P(I) = \int_{-\infty}^{y_1^1} \int_{-\infty}^{\infty} p(\hat{y}_1(x^*), \hat{y}_2(x^*)) dy_1 dy_2$$
$$+ \int_{y_1^1}^{y_2^1} \int_{-\infty}^{y_2^1} p(\hat{y}_1(x^*), \hat{y}_2(x^*)) dy_1 dy_2$$

Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration

We will use a multiobjective expected improvement framework:




Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration

We will use a multiobjective expected improvement framework:

$$\mathbb{E}[I] = \int \int d(\hat{y}, \mathcal{F}) P[I(x^*)] d\hat{y}_1 d\hat{y}_2$$




Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

2D EGO Illustration

We will use a multiobjective expected improvement framework:


$$\begin{aligned}\mathbb{E}[I] &= \int \int d(\hat{y}, \mathcal{F}) P[I(x^*)] d\hat{y}_1 d\hat{y}_2 \\ &= f(\mu_1, \mu_2, \sigma_1, \sigma_2, \mathcal{F})\end{aligned}$$



Keane, A. J., *AIAA Journal*, 44(4):879–891, 2006.

Design space and clustering


Jump start the design with diversity-oriented cluster:



Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Design space and clustering


Jump start the design with diversity-oriented cluster:



Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Design space and clustering



Jump start the design with diversity-oriented cluster:



Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.



Design space and clustering

Jump start the design with diversity-oriented cluster:





Design space and clustering

Jump start the design with diversity-oriented cluster:





Design space and clustering

Jump start the design with diversity-oriented cluster:



Design space and clustering



Jump start the design with diversity-oriented cluster:



Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Design space and clustering


Jump start the design with diversity-oriented cluster:




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




DFT Data




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$E[I]$




DFT Data




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$E[I]$




DFT Data




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$E[I]$



DFT Data




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI

 $P[I]$ $\mathbb{E}[I]$


DFT Data




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




k-medoids points (gen 1)




Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




pareto front (gen 1)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




EI points (generation 2)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




pareto front (gen 2)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




EI points (gen 3)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




pareto front (gen 3)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




EI points (gen 4)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




pareto front (gen 4)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




EI points (gen 5)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$




pareto front (gen 5)




Janet, J.P., et al., ACS Cent. Sci., 6(4):513–524, 2020.

Evolution of PI and EI


$P[I]$



$\mathbb{E}[I]$



convex hulls



Introduction
oooo

Prediction and uncertainty
oooooooo

Case study
oooooo●

Drug discovery
oooo

Final thoughts
oo

Case study conclusions

Case study conclusions

- EI framework provides high resolution in the region of interest, converges quickly
- Able to identify fruitful regions from a large chemical space based few DFT evaluations
- Multiobjective DFT optimization guided by ML efficiency generates lead complexes, $> 100\times$ faster than random search

Case study conclusions

- EI framework provides high resolution in the region of interest, converges quickly
- Able to identify fruitful regions from a large chemical space based few DFT evaluations
- Multiobjective DFT optimization guided by ML efficiency generates lead complexes, $> 100\times$ faster than random search

Thanks to Professor Heather Kulik and her lab at MIT



Case study conclusions

- EI framework provides high resolution in the region of interest, converges quickly
- Able to identify fruitful regions from a large chemical space based few DFT evaluations
- Multiobjective DFT optimization guided by ML efficiency generates lead complexes, $> 100\times$ faster than random search

Thanks to Professor Heather Kulik and her lab at MIT



github.com/hjkgrp/molSimplify
github.com/hjkgrp/AutomaticDesign

Drug discovery

And now for something completely different

Drug discovery


And now for something completely **moderately** different

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.

Drug discovery


Bringing a new drug costs upwards of \$1 billion and can take decades.



protein


Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.



Drug discovery


Bringing a new drug costs upwards of \$1 billion and can take decades.



This is very difficult because of the many requirements for drug-like molecules:

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard)
- Specificity, tolerability, toxicity (very hard)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard)
- Specificity, tolerability, toxicity (very hard)
- Pharmacokinetics and pharmacodynamics (hard)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard)
- Specificity, tolerability, toxicity (very hard)
- Pharmacokinetics and pharmacodynamics (hard)
- Synthetic accessibility (moderate)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard) – can be predicted with physics (FEP), sometimes
- Specificity, tolerability, toxicity (very hard)
- Pharmacokinetics and pharmacodynamics (hard)
- Synthetic accessibility (moderate)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard) – can be predicted with physics (FEP), sometimes
- Specificity, tolerability, toxicity (very hard) – mostly predicted from data
- Pharmacokinetics and pharmacodynamics (hard)
- Synthetic accessibility (moderate)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.




This is very difficult because of the many requirements for drug-like molecules:

- Affinity/potency (hard) – can be predicted with physics (FEP), sometimes
- Specificity, tolerability, toxicity (very hard) – mostly predicted from data
- Pharmacokinetics and pharmacodynamics (hard) – mostly predicted from data
- Synthetic accessibility (moderate)

Drug discovery

Bringing a new drug costs upwards of \$1 billion and can take decades.



This is very difficult because of the many requirements for drug-like molecules:


- Affinity/potency (hard) – can be predicted with physics (FEP), sometimes
- Specificity, tolerability, toxicity (very hard) – mostly predicted from data
- Pharmacokinetics and pharmacodynamics (hard) – mostly predicted from data
- Synthetic accessibility (moderate) – ML synthesis planning very advanced

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning

The REINVENT ecosystem


We tackle these challenges with generative reinforcement learning




text-based molecular representation generated by
an RNN

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




text-based molecular representation generated by
an RNN





Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.
Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




text-based molecular representation generated by an RNN



- Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.
Olivcrona, M., et al., *J. Cheminform.* 9(48), 2017.
Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.

Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.


Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.

Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.




The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




- Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.
Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.
Olivcrona, M., et al., *J. Cheminform.* 9(48), 2017.
Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.




The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning



$$R(a) = \sigma S(a) - \log \pi_\infty(a)$$



Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.


Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.

Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.


Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning



$$R(a) = \underbrace{\sigma S(a)}_{\text{actor}} - \underbrace{\log \pi_\infty(a)}_{\text{critic}}$$



Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.


Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.

Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.


Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




$$R(a) = \sigma S(a) - \log \pi_\infty(a)$$





- Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.
Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.
Olivcrona, M., et al., *J. Cheminform.* 9(48), 2017.
Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning



$$R(a) = \sigma S(a) - \log \pi_\infty(a)$$



Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.


Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.

Olivcrona, M., et al., *J. Cheminform.* 9(48), 2017.


Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning



$$R(a) = \sigma S(a) - \log \pi_\infty(a)$$





Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.

Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.


Olivcrona, M., et al., *J. Cheminform.* 9(48), 2017.

Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.




The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




$$R(a) = \sigma S(a) M(a) - \log \pi_\infty(a)$$



- Blaschke, T., et al., *J. Cheminform.* 12(68), 2020.
Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.
Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.
Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.
Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning




Blaschke, T., et al., *J. Cheminform.* 12(68), 2020.

Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.

Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.

Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.


Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.



Janet, J.P., et al., *in preparation*.

The REINVENT ecosystem

We tackle these challenges with generative reinforcement learning



Blaschke, T., et al., *J. Cheminform.* 12(68), 2020.


Arús-Pous, J., et al., *J. Cheminform.* 11(20), 2019.

Arús-Pous, J., et al., *J. Cheminform.* 11(71), 2019.

Olivecrona, M., et al., *J. Cheminform.* 9(48), 2017.

Blaschke, T., et al., *J. Chem. Inf. Model.* 60(12):5918–5922, 2020.


$$R(a) = \sigma S(a) M(a) - \log \pi_\infty(a)$$



Janet, J.P., et al., *in preparation*.


Docking and curriculum learning

Developed engine for conditioning on protein structure.






Docking and curriculum learning

Developed engine for conditioning on protein structure. Nice, but slow! ~ 8 minutes/epoch.







Docking and curriculum learning

Developed engine for conditioning on protein structure. Nice, but slow! ~ 8 minutes/epoch.



Docking and curriculum learning

Developed engine for conditioning on protein structure. Nice, but slow! ~ 8 minutes/epoch.







Implemented curriculum learning to speed up the process and learn difficult objectives.
Add physics-based models last!

Guo, J., et al., *ChemRxiv*:chemrxiv-2021-37m7w-v2
Guo, J., et al., *J. Cheminform.* 13(89), 2021.

Docking and curriculum learning

Developed engine for conditioning on protein structure. Nice, but slow! ~ 8 minutes/epoch.






Implemented curriculum learning to speed up the process and learn difficult objectives.
Add physics-based models last!



Guo, J., et al., *ChemRxiv*:chemrxiv-2021-37m7w-v2
Guo, J., et al., *J. Cheminform.* 13(89), 2021.

Docking and curriculum learning

Developed engine for conditioning on protein structure. Nice, but slow! ~ 8 minutes/epoch.




Implemented curriculum learning to speed up the process and learn difficult objectives.
Add physics-based models last!



Guo, J., et al., *ChemRxiv*:chemrxiv-2021-37m7w-v2
Guo, J., et al., *J. Cheminform.* 13(89), 2021.


Other Projects

Transfer learning to exploit 'big' affinity data:



Other Projects



Transfer learning to exploit 'big' affinity data:



Buterez, D., et al., *in preparation*.


Other Projects

Transfer learning to exploit 'big' affinity data:




Other Projects

Transfer learning to exploit 'big' affinity data:




Protein pocket analysis with e3nns:




Navarro, A.C., Ghabi, Y., et al., *in preparation*.
Buterez, D., et al., *in preparation*.

Other Projects

Transfer learning to exploit 'big' affinity data:




Protein pocket analysis with e3nns:



Navarro, A.C., Ghabi, Y., et al., *in preparation*.
Buterez, D., et al., *in preparation*.

More than just ML (or simulation)


physics-driven




slow, accurate (?)

More than just ML (or simulation)

physics-driven



data-driven




slow, accurate (?)

fast, uncertainty-aware


More than just ML (or simulation)

physics-driven



slow, accurate (?)


data-driven



fast, uncertainty-aware


More than just ML (or simulation)

physics-driven



slow, accurate (?)



data-driven



fast, uncertainty-aware



More than just ML (or simulation)

physics-driven



slow, accurate (?)


data-driven



fast, uncertainty-aware



More than just ML (or simulation)

physics-driven




slow, accurate (?)

data-driven



fast, uncertainty-aware

More than just ML (or simulation)



What's next?

Data-driven methods are changing how solve chemical design problems. What's next?

What's next?

Data-driven methods are changing how solve chemical design problems. What's next?

- Neural network potentials - quantum accuracy, force field cost!


Name	Molecule	MAE	RMSE	Scan (Left:ANI Right:DFT)
Cysteine-Dipeptide (25 atoms)		1.75	2.55	
DDT (28 atoms)		0.53	0.70	
Hexafluoroacetone (10 atoms)		0.08	0.11	
Bendamustine (44 atoms)		0.50	0.66	

Devereux, C., et al., *J. Chem. Theory Comput.*, 16(7):4192–4202, 2020

What's next?

Data-driven methods are changing how solve chemical design problems. What's next?

- Neural network potentials - quantum accuracy, force field cost!
- Synthesis planning and optimization. Fully automated chemistry!




Caramelli, D., et al., ACS Cent. Sci., 7(11):1821–1830, 2021

What's next?

Data-driven methods are changing how solve chemical design problems. What's next?

- Neural network potentials - quantum accuracy, force field cost!
- Synthesis planning and optimization. Fully automated chemistry!
- Generative models. Designing new drugs directly into the pocket, *de novo*!



Ragoza, M., et al., arXiv:2010.08687v3, 2020

Arcidiacono, M. & Koes, D.R., et al., <https://arxiv.org/abs/2109.15308>, 2021