

Chapter 1

Ancestral Reconstruction of Bat Echolocation Calls

J.P. Meagher*

*Department of Statistics,
University of Warwick,
J.Meagher@Warwick.ac.uk[†]*

1. Introduction

Advances in technology allowing the precise quantification and storage of information about the world around us continues to drive the emergence of Data Science as a discipline distinct from both Statistics and Computer Science.

Bioacoustics is one area of research generating vast quantities of data which also captures the imagination of the public, as evidenced by successful citizen science initiatives.¹² Bioacoustic techniques for biodiversity monitoring³⁴ have the potential to make real policy impacts, particularly with regard to sustainable economic development and nature conservation.

In the acoustic monitoring of biodiversity, bats (order *Chiroptera*) are of particular interest. Bats have been identified as ideal bioindicators for monitoring climate change and habitat quality,⁵ largely because bats broadcast information about themselves into their environment in the form of echolocation calls.⁶ The development of automatic acoustic monitoring algorithms for classifying species of bats³⁷ means that large scale, non-invasive monitoring is becoming possible.

While monitoring bat populations provides useful information, understanding the root causes and effects of what is observed requires that the natural history of extant bat species is also well understood. Traits, such as call structure or body size, exhibited by particular bat species are linked

* Author footnote.

[†] Affiliation footnote.

to the bats interactions with its environment.⁸ Existing fossil records are of limited use in inferring the traits exhibited by ancestral bats, particularly with respect to echolocation calls for example. The reconstruction of ancestral traits relies heavily on the comparative analysis⁹ of extant bat species. Thus, statistical data science techniques may be particularly useful for inferring the evolutionary dynamics and reconstructing ancestral states of echolocation in bats.

Previous studies of bat echolocation calls for both classification⁷ and ancestral reconstruction¹⁰ examined features of the call extracted from the spectrogram of the call. These, somewhat arbitrary, call features relied on significant domain knowledge to ensure they were sensibly selected and used. More recently however, general techniques for the classification of acoustic signals have been developed.^{11,4} These methods do not require, but can be augmented by, domain knowledge. General techniques for ancestral reconstruction of function-valued traits, such as speech sounds or echolocation calls, have been proposed.¹² The study of bat echolocation calls offers an opportunity to examine the efficacy of these techniques.

A function-valued trait is measured along some continuous scale, usually time, and can then be modelled as a continuous mathematical function using techniques for functional data analysis.¹³ Jones & Moriarty¹⁴ develop a method which extends Gaussian Process Regression¹⁵ to model the evolution of function-valued traits.

References

1. P. E. Allen and C. B. Cooper. Citizen science as a tool for biodiversity monitoring. (2006).
2. N. Pettorelli, J. E. Baillie, and S. M. Durant, Indicator bats program: a system for the global acoustic monitoring of bats (2013).
3. V. Stathopoulos, V. Zamora-Gutierrez, K. E. Jones, and M. Girolami, Bat echolocation call identification for biodiversity monitoring: a probabilistic approach, *Journal of the Royal Statistical Society: Series C (Applied Statistics)* (2017).
4. T. Damoulas, S. Henry, A. Farnsworth, M. Lanzone, and C. Gomes. Bayesian classification of flight calls with a novel dynamic time warping kernel. In *Machine Learning and Applications (ICMLA), 2010 Ninth International Conference on*, pp. 424–429 (2010).
5. G. Jones, D. S. Jacobs, T. H. Kunz, M. R. Willig, and P. A. Racey, Carpe noctem: the importance of bats as bioindicators, *Endangered species research*. **8**(1-2), 93–115 (2009).
6. G. Jones, Echolocation, *Current Biology*. **15**(13), R484–R488 (2005).

7. C. L. Walters, R. Freeman, A. Collen, C. Dietz, M. Brock Fenton, G. Jones, M. K. Obrist, S. J. Puechmaille, T. Sattler, B. M. Siemers, et al., A continental-scale tool for acoustic identification of european bats, *Journal of Applied Ecology*. **49**(5), 1064–1074 (2012).
8. H. Aldridge and I. Rautenbach, Morphology, echolocation and resource partitioning in insectivorous bats, *The Journal of Animal Ecology*. pp. 763–778 (1987).
9. J. Felsenstein and J. Felsenstein, *Inferring phylogenies*. vol. 2, Sinauer associates Sunderland (2004).
10. A. Collen. *The evolution of echolocation in bats: a comparative approach*. PhD thesis, UCL (University College London) (2012).
11. V. Stathopoulos, V. Zamora-Gutierrez, K. Jones, and M. Girolami. Bat call identification with gaussian process multinomial probit regression and a dynamic time warping kernel. In *Artificial Intelligence and Statistics*, pp. 913–921 (2014).
12. T. F. P. Group, Phylogenetic inference for function-valued traits: speech sound evolution, *Trends in ecology & evolution*. **27**(3), 160–166 (2012).
13. J. O. Ramsay, *Functional data analysis*. Wiley Online Library (2006).
14. N. S. Jones and J. Moriarty, Evolutionary inference for function-valued traits: Gaussian process regression on phylogenies, *Journal of The Royal Society Interface*. **10**(78), 20120616 (2013).
15. C. E. Rasmussen, Gaussian processes for machine learning (2006).