

# Shaping the chameleon

JONATHAN A. PEARSON\*

*School of Physics & Astronomy  
University of Nottingham  
Nottingham, NG7 2RD*

October 15, 2014

## **Abstract**

In conservation with Clare Burrage, Ed Copeland, James...

---

\*E-mail: [j.pearson@nottingham.ac.uk](mailto:j.pearson@nottingham.ac.uk)

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>The model</b>	<b>2</b>
<b>3</b>	<b>Numerical methods</b>	<b>2</b>
	<b>Acknowledgements</b>	<b>2</b>
<b>A</b>	<b>Schemes</b>	<b>2</b>
	A.1 Fourth order finite difference derivatives . . . . .	2
	<b>References</b>	<b>3</b>

## 1 Introduction

## 2 The model

The equations are

$$\nabla^2 \phi = -\frac{\Lambda^5}{\phi^2} + \frac{\rho}{M}, \quad (2.1a)$$

$$\nabla^2 \Phi = -\rho \quad (2.1b)$$

We compute the forces due to the chameleon and gravitational scalars by taking their gradients:

$$\mathbf{F}_{(\phi)} = \nabla \phi, \quad \mathbf{F}_{(\Phi)} = \nabla \Phi \quad (2.2)$$

## 3 Numerical methods

- Solve (2.1a) via gradient flow; finite difference derivatives discretized to fourth order accuracy.
- Solve (2.1b) via SoR; discretized to second order accuracy.
- The forces (2.2) are computed with finite difference derivatives discretized to fourth order

## Acknowledgements

## A Schemes

### *SoR*

$$Q_{i,j}^{n+1} = (1 - \omega)Q_{i,j}^n + \frac{\omega}{4} [Q_{i+1,j}^n + Q_{i-1,j}^{n+1} + Q_{i,j+1}^n + Q_{i,j-1}^{n+1} + h^2 \rho] \quad (A.1)$$

*Fourth order finite difference derivatives*

$$\frac{\partial Q}{\partial x} \approx \frac{-Q_{i+2} + 8Q_{i+1} - 8Q_{i-1} + Q_{i-2}}{12h} \quad (\text{A.2})$$

$$\frac{\partial^2 Q}{\partial x^2} \approx \frac{-Q_{i+2} + 16Q_{i+1} - 30Q_i + 16Q_{i-1} - Q_{i-2}}{12h^2} \quad (\text{A.3})$$