

ALL YOUR QUESTIONS ANSWERED

*The Comprehensive FAQ for Blood Pressure,
Hypertension, Natural Remedies, and Getting Off Meds*

**40 Real Questions From Real People.
Honest Answers From a 20-Year RN.**

40

questions answered from real
people

7

categories covered

20

years of clinical nursing
experience

0

BS or sugarcoating

Joel Polley, RN | @braveworksrn
BraveWorksRN.com

DISCLAIMER: This guide is for educational purposes only. Always consult your healthcare provider before making changes to medications or treatment plans. This is not medical advice.

HOW TO USE THIS FAQ

These are the 40 most common questions I get asked — in my TikTok comments, in my Skool community, in DMs, and in the questions I hear nurses and patients struggling with every single day. They are organized into 7 categories. I wrote every answer the way I would answer you if we were sitting across from each other at my kitchen table. Straight talk. Real science. No sugarcoating.

If your specific question is not in here, chances are the answer is woven into one of the other guides in this kit — the Pharma vs. Farmacy comparison, the White Coat Syndrome Guide, the Top 10 Herbs Deep Dive, or the 10-Day BP Reset Challenge PDFs. Between all of them, you have the most comprehensive natural blood pressure resource available anywhere.

CATEGORY	QUESTIONS	# OF Qs
1. General Blood Pressure	What is normal, is it genetic, can it be cured, am I too young for this	7
2. Medication Fears	Stopping meds, side effects, weaning off, lifetime dependency	7
3. Herbs & Supplements	Dosing, interactions, which ones, stacking, how long to work	8
4. Diet & Nutrition	Salt, coffee, alcohol, sugar, water, DASH diet, processed food	6
5. Lifestyle & Exercise	Stress, sleep, breathing, exercise type, weight loss	5
6. Monitoring & Readings	White coat, home monitors, which arm, fluctuations	4
7. Specific Situations	Anxiety-driven BP, pregnancy, young people, hormones	3

CATEGORY 1: GENERAL BLOOD PRESSURE QUESTIONS

Q: "What is actually considered normal blood pressure? I keep seeing different numbers."

Normal is less than 120/80. Elevated is 120-129 over less than 80. Stage 1 hypertension is 130-139 over 80-89. Stage 2 is 140+ over 90+. Anything over 180/120 is a hypertensive crisis — go to the ER. The confusion comes from guidelines changing in 2017 when the AHA lowered the threshold from 140/90 to 130/80. Overnight, 30 million more Americans became "hypertensive." Same blood, same vessels — different number on a chart.

Q: "My whole family has high blood pressure. Am I genetically screwed?"

Genetics writes the recipe. Lifestyle bakes the cake. Yes, there is a genetic component to blood pressure. But here is what they do not tell you: what gets passed down in families is not just DNA — it is habits. If your mother cooked with excessive sodium, you learned to cook with excessive sodium. If your father sat on the couch every night, you learned to sit on the couch every night. If your family handled stress with anger and alcohol, you learned to handle stress with anger and alcohol. Genetics may load the gun. But lifestyle pulls the trigger. The NEWSTART lifestyle program demonstrated that systolic blood pressure dropped an average of 19 points in just 14 days, and 93% of participants reduced or stopped BP medications — many of whom had "genetic" hypertension.

Nurse JP's Note: I have watched patients come into the hospital with 'genetic' hypertension and leave with normal readings after two weeks of dietary changes. The genes did not change. The inputs did.

Q: "I'm only in my 20s/30s — how is it possible I have hypertension?"

It is not only possible, it is increasingly common. The Standard American Diet is higher in sodium, sugar, and processed food than at any point in human history. Stress levels among young adults are at record highs. Screen time is replacing sleep. Caffeine intake starts younger and gets heavier. Energy drinks did not exist 30 years ago — now a single can contains 200-300mg of caffeine plus taurine and other stimulants. Your age is not protection. Your inputs are either protecting you or harming you, regardless of the year on your birth certificate.

Q: "Can high blood pressure actually be reversed, or is this a life sentence?"

If I believed it was a life sentence, I would not have built this entire program. High blood pressure is an **effect**. It has causes. Remove the causes, add the right support, and the effect changes. That is not opinion — it is the law of cause and effect. Clinical trials prove it. The PREMIER trial showed comprehensive lifestyle modifications reduced hypertension prevalence by 77% — matching medication. The question is not "can it be reversed?" The question is "are you willing to change the inputs?" If yes, your body will do what it was designed to do.

Q: "Is high blood pressure really dangerous if I feel completely fine?"

This is why it is called the silent killer. You can walk around with a reading of 160/100 and feel perfectly normal. No headaches. No dizziness. Meanwhile, that pressure is silently damaging your arterial lining, thickening your heart muscle, scarring your kidneys, and weakening blood vessels in your brain. By the time you "feel" something, the damage is often severe — a stroke, a heart attack, kidney failure, or vision loss. Feeling fine is not the same as being fine. The numbers do not lie. Trust the numbers.

Q: "What do the top and bottom numbers actually mean?"

Systolic (top number) is the pressure when your heart BEATS — the force of blood pushing against your artery walls during a contraction. Diastolic (bottom number) is the pressure when your heart RESTS between beats. Both matter, but systolic tends to rise with age and is the stronger predictor of cardiovascular events in people over 50. In younger adults, a high diastolic can signal early-stage hypertension. Do not ignore either number.

Q: "I had one really high reading but the rest were fine — should I be worried?"

One reading is a snapshot, not a diagnosis. Blood pressure fluctuates constantly — it changes with every heartbeat, every breath, every emotion, every sip of coffee, every trip to the bathroom. A single high reading could mean you were stressed, dehydrated, had a full bladder, just exercised, or drank caffeine. What matters is the **pattern** over days and weeks. Monitor at home, 2-3 times per day for 1-2 weeks, and look at the trend. If the trend is consistently elevated, act. If it was one spike, note it and move on.

CATEGORY 2: MEDICATION FEARS

Q: "Will I be on blood pressure pills forever?"

Not necessarily — and that is the entire premise of everything I teach. Blood pressure medication manages the **effect** while leaving the **cause** untouched. If you change the cause — the diet, the stress, the sleep, the movement, the inputs — your body can begin regulating pressure on its own again. I have seen it happen hundreds of times. The NEWSTART program documented 93% of participants reducing or stopping medications in 14 days. But — and this is critical — you must change the inputs FIRST. The pills come off LAST, and only under your doctor's supervision. Not the other way around.

Q: "My doctor wants to put me on medication but I really don't want to take it."

Here is my honest answer: if your blood pressure is dangerously high RIGHT NOW (Stage 2, above 140/90 consistently, or crisis level above 180/120), medication may be necessary to protect you while you make lifestyle changes. A stroke does not wait for your elimination diet to kick in. But here is what you CAN say to your doctor: "I would like to try intensive lifestyle modifications for 30-60 days and bring you my home readings. If my numbers have not improved, I am open to medication." Most doctors will respect that — especially if you show up with a plan and data. This kit IS that plan.

Q: "The side effects are worse than the condition — anyone else feel this way?"

You are not alone. ACE inhibitors (lisinopril) cause a persistent dry cough in up to 20% of users. Amlodipine causes ankle swelling. Beta-blockers cause fatigue, weight gain, and can suppress exercise tolerance. Diuretics flush potassium, magnesium, and CoQ10 from your body — the very nutrients your vessels need to relax. These are not rare side effects. They are common. And they are one of the top reasons 54% of patients either stop taking BP meds or do not take them as directed. This is why I teach the cause-and-effect approach: address the root cause so the medication becomes less necessary over time.

Q: "My BP is normal now — does that mean the meds are working or that I don't need them?"

If you are on medication and your BP is normal, it usually means the medication is doing its job. That does NOT mean you can stop cold turkey. Abruptly stopping blood pressure medication can cause a dangerous rebound spike — your body has adapted to the drug, and removing it suddenly can send your pressure skyrocketing. The safe path: continue medication, make lifestyle changes, monitor your numbers, and when readings are consistently LOW (not just normal), bring your data to your doctor. Let THEM reduce the dose gradually. Your results do the talking.

Q: "Can I wean off my meds if I've lost weight and changed my diet?"

Potentially — and this is the goal. Research shows that every 2.2 pounds of weight lost reduces systolic BP by approximately 1 mmHg. If you have lost significant weight, cleaned up your diet, added exercise, and your home readings are consistently lower than they were when you started medication, you have a strong case. Bring your tracking data to your doctor. The conversation is: "My numbers are improving. Can we talk about reducing my dosage?" Your doctor will typically taper one medication at a time while monitoring your response.

Q: "Do blood pressure meds cause weight gain, fatigue, or depression?"

Beta-blockers (metoprolol, atenolol, propranolol) are the most common culprits. They slow your heart rate and reduce your body's ability to burn calories during exercise. Many patients describe feeling like a "zombie." Weight gain of 5-10 pounds is documented. They can also worsen depression and fatigue. Diuretics can cause muscle weakness from mineral depletion. ACE inhibitors and ARBs tend to have fewer of these effects. If side effects are impacting your quality of life, tell your doctor. There are multiple drug classes — and a lifestyle-first approach may eventually reduce or eliminate the need for any of them.

Q: "Can over-the-counter pain meds or cold medicine raise my blood pressure?"

Yes. NSAIDs like ibuprofen (Advil, Motrin) and naproxen (Aleve) can raise blood pressure by 3-6 mmHg and counteract the effects of BP medication. Decongestants like pseudoephedrine (Sudafed) are even worse — they directly constrict blood vessels. Many cold medicines contain both. If you have hypertension, use acetaminophen (Tylenol) for pain and avoid decongestants. Read every label. The ingredients that raise BP are hiding in products you would never suspect.

CATEGORY 3: HERBS & SUPPLEMENTS

Q: "Can I take herbs/supplements with my blood pressure medication? Will they interact?"

This is the single most asked question on every blood pressure video on TikTok — and the answer requires nuance. Many herbs work through the SAME mechanism as your medication (see the Pharma vs. Farmacy guide in this kit). That means combining them can potentially **double the effect** and drop your pressure too low. Hibiscus is a natural ACE inhibitor — taking it with lisinopril is like taking a double dose. Magnesium is a natural calcium channel blocker — combining it with amlodipine needs monitoring. This does not mean you cannot do it. It means you **START LOW, MONITOR DAILY, and TELL YOUR DOCTOR** what you are taking. Add one supplement at a time, watch your readings for a week, then consider adding the next.

***Nurse JP's Note:** The goal is not to replace your medication overnight. It is to add natural support, watch the numbers improve, and then have YOUR DOCTOR reduce the pharmaceutical dose based on your data.*

Q: "Does hibiscus tea actually lower blood pressure? How many cups a day?"

Yes — and the research is solid. A meta-analysis of multiple clinical trials found hibiscus tea reduced systolic BP by an average of 7.5 mmHg and diastolic by 3.5 mmHg. It works as a natural ACE inhibitor AND a mild diuretic — two mechanisms in one cup. Dose: 2-3 cups per day of strong hibiscus tea (steep 5-10 minutes with a lid on to keep the volatile oils in). Use dried hibiscus flowers or quality tea bags. It can be consumed hot or iced. The deep red color comes from anthocyanins — the same antioxidants that protect your blood vessel walls.

Q: "Which type of magnesium is best for blood pressure?"

Magnesium glycinate is the gold standard — best absorbed, least GI upset, and has a calming effect that supports sleep. Magnesium taurate is excellent specifically for cardiovascular health — taurine itself supports heart function. Magnesium citrate is well-absorbed but has a mild laxative effect. **AVOID** magnesium oxide — it is the cheapest form on the shelf but has only 4% bioavailability. You are essentially paying to flush it down the toilet. Dose: 300-400mg daily, taken with dinner or before bed for best results.

Q: "Does garlic really work? Raw cloves vs. supplements vs. aged garlic — which one?"

Garlic is one of the most powerful multi-mechanism plants for blood pressure. It inhibits ACE, stimulates nitric oxide and hydrogen sulfide, blocks calcium channels, and blocks alpha-adrenergic receptors. That is **FOUR** drug classes in one clove. Raw garlic is the most potent — **CRUSH** the clove and let it sit for 10 minutes before eating. This activates the enzyme alliinase that converts alliin into allicin. Without those 10 minutes, you are just eating a vegetable. Aged garlic extract (Kyolic brand, 600-1200mg/day) is the most studied supplement form and is gentler on the stomach. Both work. The key is consistency — garlic's effects build over 2-4 weeks.

Q: "How long does it take for supplements to start working?"

It depends on the supplement. Some effects are immediate: beet juice can lower BP within hours (nitric oxide pathway). Cayenne pepper causes vasodilation within minutes of consumption. Others take time to build: ashwagandha takes 2-4 weeks for full cortisol reduction. Garlic takes 2-4 weeks for ACE inhibition to peak. Hawthorn berry can take 4-8 weeks for full effect. Magnesium begins working within days but

reaches full impact over 2-4 weeks. The common mistake is trying something for 3 days, seeing no change, and quitting. Give each supplement a minimum of 4 weeks before judging. And monitor your BP daily so you can SEE the trend.

Q: "Has anyone actually tried hawthorn berry? Tea or capsules?"

Hawthorn has been used for cardiovascular support for over 2,000 years, and a 2020 systematic review confirmed its ability to lower blood pressure and reduce heart rate. It works similarly to beta-blockers — slowing the heart and reducing cardiac workload — but without the fatigue and weight gain. Both tea and capsules work. For tea: steep 1-2 teaspoons of dried hawthorn berries for 15-20 minutes (longer steep = stronger extraction). For capsules: 250-500mg standardized extract, 2-3 times daily. Hawthorn takes longer to show effects than some herbs — give it 4-8 weeks of consistent use.

Q: "Can I take all of these together — hibiscus + garlic + magnesium + hawthorn?"

Yes — with caution and with monitoring. These herbs work through DIFFERENT mechanisms (ACE inhibition, calcium channel blocking, beta-blocking, diuretic), so stacking them is similar to how a doctor prescribes multiple medications targeting different pathways. The key is to add ONE at a time, take daily readings, and give each one a week before adding the next. If you are also on pharmaceutical medication, the combined effect can drop your pressure significantly. This is actually the goal — but you need to monitor so it does not drop TOO low. Signs of BP that is too low: dizziness when standing, lightheadedness, fatigue, blurred vision. If you experience these, reduce your supplementation and talk to your doctor about reducing your medication.

Q: "Does apple cider vinegar do anything for blood pressure?"

The honest answer: the evidence is thin. There are some small animal studies suggesting acetic acid may have a mild effect on blood pressure through renin-angiotensin modulation, but the human clinical trial data is not strong enough for me to recommend it as a primary intervention. If you enjoy it, it will not hurt you — 1-2 tablespoons diluted in water before meals. But I would not rely on it over hibiscus, garlic, magnesium, or any of the other herbs with significantly stronger research behind them. Do not fall for the TikTok hype on this one. Focus on what is proven.

CATEGORY 4: DIET & NUTRITION

Q: "Do I really have to give up salt completely? Everything tastes like cardboard."

No — you do not have to eliminate salt entirely. You need to eliminate EXCESS sodium. The AHA recommends no more than 1,500mg/day for people with hypertension. The average American consumes 3,400mg. Here is the secret: 75% of that sodium is coming from PROCESSED and RESTAURANT food — not your salt shaker. Bread, deli meat, canned soup, frozen meals, condiments, cheese — these are the real culprits. Cook from whole ingredients, season with herbs, garlic, lemon, and spices, and your taste buds will recalibrate within 5-7 days. Food will actually taste MORE flavorful, not less, once your palate resets.

Q: "Is coffee really that bad? Do I have to give it up?"

Caffeine can spike blood pressure by 5-10 mmHg within 30 minutes of consumption. It triggers adrenaline release, increases heart rate, and constricts blood vessels. For some people — especially those who are caffeine-sensitive or whose BP is already borderline — this is the difference between normal and hypertensive. My honest recommendation: eliminate caffeine for 10-14 days and monitor your numbers. If your BP drops, you have your answer. If it does not change, moderate coffee (1 cup in the morning, not 3-4 throughout the day) may be tolerable for YOU. But you will never know until you test it. Swap to chamomile or hibiscus tea — both actively LOWER BP while you sip.

Q: "Is Celtic salt / Himalayan pink salt better for blood pressure than regular salt?"

I need to be straight with you: sodium is sodium. Celtic salt and Himalayan pink salt contain trace minerals — tiny amounts of magnesium, potassium, and calcium — but the primary ingredient is still sodium chloride. The trace mineral content is so small that you would need to consume dangerously high amounts of salt to get any meaningful mineral benefit. One gram of Himalayan salt contains approximately 368mg of sodium — virtually identical to table salt. If you prefer the taste, use it. But do not fool yourself into thinking it is "healthy" salt that does not affect your blood pressure. It does.

Nurse JP's Note: This is one of the biggest myths circulating on TikTok right now. Do not fall for it. Reduce your sodium intake regardless of the color of the salt.

Q: "Can I drink alcohol with high blood pressure?"

Alcohol activates the sympathetic nervous system, increases cortisol, raises heart rate, and disrupts sleep quality — all of which directly raise blood pressure. Even "moderate" drinking (1-2 drinks per day) has been shown to raise BP by approximately 4 mmHg. The idea that red wine is "heart healthy" has been largely debunked — the resveratrol content is too low to provide meaningful benefit, and the alcohol itself does more harm than the antioxidants do good. My recommendation: eliminate alcohol completely during any blood pressure reset period. After your numbers normalize, if you choose to drink, limit it to rare occasions — not daily.

Q: "Does sugar affect blood pressure too, or is it just salt?"

Sugar is a MAJOR blood pressure driver that gets far less attention than salt. High sugar intake spikes insulin, which causes your kidneys to retain sodium — so sugar raises your BP through the SAME mechanism as salt. Additionally, high fructose corn syrup (found in sodas, juice, processed foods) increases uric acid, which directly reduces nitric oxide — your body's master vasodilator. Less nitric oxide means stiffer vessels and higher pressure. A study in the American Journal of Cardiology found that

reducing sugar intake lowered systolic BP by 8 mmHg in just 2 weeks. Sugar may be a bigger BP driver than salt for many people.

Q: "Can drinking more water lower my blood pressure?"

Yes — through multiple mechanisms. When you are dehydrated, your blood gets thicker (higher viscosity), which makes your heart work harder to pump it. Your kidneys go into conservation mode and retain sodium. Your body releases vasopressin, a hormone that directly constricts blood vessels. Proper hydration reverses all three of these. The formula: drink your body weight in pounds as ounces of water daily. A 180-pound person drinks 180 ounces. Water ONLY — not juice, not sports drinks, not soda. Start with 16 ounces first thing in the morning before anything else. Your kidneys will thank you.

CATEGORY 5: LIFESTYLE & EXERCISE

Q: "How much does stress REALLY affect blood pressure?"

More than almost anything else. Chronic stress keeps cortisol elevated 24/7. Cortisol constricts blood vessels, causes sodium retention, increases heart rate, spikes insulin, promotes inflammation, and suppresses vascular repair. A single episode of intense anger increases heart attack risk 8.5x in the following 2 hours. But it is not just "feeling stressed" — anger, unforgiveness, worry, anxiety, grief, guilt, and shame ALL produce the same cortisol response. Your body does not distinguish between a real threat and a replayed argument. Every grudge you carry, every worry you replay, every guilt you hold onto is producing real cortisol in your real bloodstream right now.

Nurse JP's Note: I have seen patients whose BP was resistant to THREE medications — and it came down when they forgave someone. That is not a metaphor. That is cortisol physiology.

Q: "Can losing weight actually fix my blood pressure? How much do I need to lose?"

Weight loss is one of the single most effective interventions for blood pressure. Research shows that for every 2.2 pounds (1 kg) lost, systolic BP drops approximately 1 mmHg. So a 20-pound weight loss could reduce your systolic reading by about 9 mmHg — that is the effect of many medications. You do not need to reach your "ideal" weight. Even losing 5-10% of your body weight produces meaningful results. The mechanism: excess body fat — especially visceral belly fat — acts as an endocrine organ, producing inflammatory compounds that stiffen arteries and raise pressure. Remove the fat, remove the inflammation, the pressure drops.

Q: "What is the best exercise for blood pressure — cardio or weights?"

Both help, but the research may surprise you. A 2023 systematic review of 270 randomized trials found that **isometric exercises** — like wall sits, planks, and isometric leg extensions — produced the LARGEST blood pressure reduction of any exercise type. Wall sits alone reduced systolic BP by approximately 10 mmHg. Aerobic exercise (walking, cycling, swimming) comes next at about 5-8 mmHg reduction. Resistance training also helps. But the simplest, most accessible option remains a 20-minute walk in the sunshine. It produces nitric oxide through shear stress, lowers cortisol, generates vitamin D, and improves arterial flexibility. You do not need a gym membership. You need shoes and a door.

Q: "I just started exercising and my BP is actually HIGHER — is that normal?"

Yes — temporarily. During exercise, blood pressure rises because your heart is pumping harder and faster to deliver oxygen to working muscles. Systolic can spike to 180+ during intense exertion — this is a normal physiological response. What matters is your **RESTING** blood pressure 30-60 minutes AFTER exercise. That number should be lower than your pre-exercise reading. Over time (weeks), your resting BP will decrease as your cardiovascular fitness improves, your arteries become more elastic, and your heart becomes more efficient. Do NOT take your blood pressure during or immediately after exercise. Wait at least 30 minutes.

Q: "Do breathing exercises actually work, or is it placebo?"

It is not placebo — it is vagus nerve activation, and it is measurable. Box breathing (4 counts in, 4 hold, 4 out, 4 hold) activates the vagus nerve, which is the master switch between your sympathetic (fight-or-flight) and parasympathetic (rest-and-digest) nervous systems. When the vagus nerve fires, heart rate drops, vessels dilate, and cortisol production slows. A study in the journal Hypertension Research

found that slow breathing at 6 breaths per minute reduced systolic BP by 3.5 mmHg in a single session. Practiced 2-3 times daily for weeks, the cumulative effect compounds significantly.

CATEGORY 6: MONITORING & READINGS

Q: "Why is my BP always high at the doctor but normal at home?"

This is called white coat syndrome, and it affects 15-30% of all people with elevated office readings. The medical environment triggers a fight-or-flight response — cortisol spikes, adrenaline surges, vessels constrict, and your reading captures the PEAK of anxiety, not your resting state. The solution: monitor at home with a validated upper-arm cuff, bring 2 weeks of readings to your doctor, and tell them you experience white coat hypertension. We have an entire guide dedicated to this in the BP Reset Kit — including 10 strategies to beat it, doctor conversation scripts, and a printable 2-week tracking log.

Q: "My readings are different every single time I check — which one is the real number?"

ALL of them are real — because blood pressure fluctuates constantly. It changes with every heartbeat, every breath, every emotion, every degree of temperature, every ounce of hydration. This is normal. The mistake people make is obsessing over individual readings. What matters is the AVERAGE over time. Take 2-3 readings one minute apart, discard the first (usually the highest), and average the remaining readings. Do this at the same time daily for 1-2 weeks. The TREND is your truth. Not any single number.

Q: "Which home blood pressure monitor should I buy? Are wrist monitors accurate?"

Upper-arm cuff monitors are the clinical standard. Omron is the most recommended brand in both medical settings and by Reddit communities that have tested extensively. Avoid wrist monitors — they are highly position-sensitive and consistently less accurate than upper-arm models. Spend \$30-60 on a validated upper-arm monitor. Pro tip: take your home monitor to your next doctor visit and compare its reading to the office cuff at the same time. If they match within 5 mmHg, you know your home device is accurate.

***Nurse JP's Note:** Less than 15% of home blood pressure monitors sold internationally meet clinical accuracy standards. Buy a validated one. It is the best \$40 health investment you will ever make.*

Q: "Does it matter which arm I use? My left is always higher than my right."

Yes, it matters. A difference of up to 10 mmHg between arms is considered normal. A difference greater than 10 mmHg consistently may indicate peripheral artery disease and should be reported to your doctor. The general recommendation: use the arm that reads HIGHER as your primary arm, and use that SAME arm every time. Most clinical guidelines suggest the left arm. Consistency is more important than which arm — pick one and stick with it.

CATEGORY 7: SPECIFIC SITUATIONS

Q: "Does anxiety cause REAL hypertension, or just temporary spikes?"

Both. Anxiety causes immediate, measurable blood pressure spikes through the cortisol and adrenaline cascade. But here is what most people do not realize: if anxiety is chronic — if you live in a state of constant worry, hypervigilance, or health anxiety — those "temporary" spikes become your new baseline. Your nervous system forgets what "calm" feels like. Your vessels stay constricted. Your cortisol stays elevated. Over months and years, this IS hypertension. Many people — especially in their 20s and 30s — are being diagnosed with hypertension that is primarily anxiety-driven. The treatment is not a blood pressure pill. The treatment is addressing the anxiety. Box breathing, gratitude practice, prayer, walking, sleep optimization, and professional counseling if needed. Address the cause, and the effect changes.

***Nurse JP's Note:** If you have developed a fear of checking your own blood pressure — where the act of taking a reading spikes your reading — limit checks to 3-5 times per week at a set time. Do not obsessively recheck.*

Q: "I developed high blood pressure during pregnancy — will it go away?"

Gestational hypertension and preeclampsia typically resolve within 6-12 weeks after delivery. However — and this is important — women who experience hypertension during pregnancy are at significantly higher risk of developing chronic hypertension later in life. Consider it an early warning signal from your body. Use it as motivation to build the lifestyle habits now that will protect you for decades. Everything in this kit applies to postpartum recovery: the dietary changes, the walking, the sleep optimization, the stress management. Your body just told you it is vulnerable. Listen to it.

***Nurse JP's Note:** If you are currently pregnant and experiencing high blood pressure, please work closely with your OB-GYN. Do NOT take herbal supplements during pregnancy without your provider's explicit approval.*

Q: "I'm going through menopause / my testosterone is low — is that why my BP is climbing?"

Very likely. This is one of the most underdiagnosed causes of blood pressure elevation. For women: estrogen is a powerful cardiovascular protector — it stimulates nitric oxide production, reduces inflammation, maintains vascular elasticity, and regulates the renin-angiotensin system. When estrogen declines during perimenopause and menopause, blood pressure often rises for the first time. For men: testosterone supports nitric oxide production, reduces visceral fat, and counterbalances cortisol. As testosterone drops 1-2% per year after age 30, cortisol becomes dominant, belly fat increases, and blood pressure climbs. Everything in this kit — walking, sleep, whole foods, stress reduction, magnesium, ashwagandha — directly supports hormone balance. See Day 7 of the 10-Day Challenge for the full hormone-BP deep dive.

STILL HAVE QUESTIONS?

This FAQ covers the 40 most common questions, but I know there are more. Here is where to find me:

TikTok: @braveworksrn — I answer comments daily

Skool Community: Free to join. Post your question and I will answer it personally.

Website: BraveWorksRN.com

The fact that you are reading this means you are not accepting the default. You are not just taking the pill and going home. You are asking questions. You are learning. You are becoming the most informed person in the room — including, in many cases, the exam room.

That is what it means to be your own doctor.

FINAL REMINDER

Nothing in this FAQ replaces the advice of your personal healthcare provider. I am an RN with 20 years of clinical experience, and I built this to educate and empower you — not to replace the relationship between you and your doctor.

NEVER stop blood pressure medication without your doctor's supervision.

Use the knowledge in this kit to become an informed, active partner in your own care. Bring data. Ask questions. Let your results speak. And let your doctor make medication decisions based on the evidence YOUR body is producing.

Joel Polley, RN | @braveworksrn | BraveWorksRN.com

Your body is a temple. Start treating it like one.

This guide is part of the BP Reset Kit from BraveWorks RN.