

# Homogeneous Systems

## Definition

**Homogeneous:** A system of equations is **homogeneous** if it can be written in the form  $A\mathbf{x} = \mathbf{0}$ .

## Example

$$\begin{aligned}x_1 + x_2 &= 1 \\x_1 - 4x_2 &= 0\end{aligned}$$

$$\begin{aligned}x_1 + x_2 &= 0 \\x_1 - 4x_2 &= 0\end{aligned}$$

# Homogeneous Systems

## Definition

**Homogeneous:** A system of equations is **homogeneous** if it can be written in the form  $A\mathbf{x} = \mathbf{0}$ .

## Example

$$\begin{aligned}x_1 + x_2 &= 1 \\x_1 - 4x_2 &= 0\end{aligned}$$

Not homogeneous!

$$\begin{aligned}x_1 + x_2 &= 0 \\x_1 - 4x_2 &= 0\end{aligned}$$

Yes it is homogeneous!

# Homogeneous Systems

## Definition

**Homogeneous:** A system of equations is **homogeneous** if it can be written in the form  $A\mathbf{x} = \mathbf{0}$ .

## Example

$$\begin{aligned}x_1 + x_2 &= 1 \\x_1 - 4x_2 &= 0\end{aligned}$$

Not homogeneous!

$$\begin{aligned}x_1 + x_2 &= 0 \\x_1 - 4x_2 &= 0\end{aligned}$$

Yes it is homogeneous!

Every homogeneous system has a **trivial solution** where  $\mathbf{x} = \mathbf{0}$ .

## Nontrivial Solutions Existence

Recall that the system  $A\mathbf{x} = \mathbf{b}$  for all  $\mathbf{b} \in \mathbb{R}^m$  has:

## Nontrivial Solutions Existence

Recall that the system  $A\mathbf{x} = \mathbf{b}$  for all  $\mathbf{b} \in \mathbb{R}^m$  has:

- ▶ A unique solution if  $A$  has a pivot in every row

# Nontrivial Solutions Existence

Recall that the system  $A\mathbf{x} = \mathbf{b}$  for all  $\mathbf{b} \in \mathbb{R}^m$  has:

- ▶ A unique solution if  $A$  has a pivot in every row
- ▶ Infinite number of solutions if  $A$  has a solution and at least one free variable

# Nontrivial Solutions Existence

Recall that the system  $A\mathbf{x} = \mathbf{b}$  for all  $\mathbf{b} \in \mathbb{R}^m$  has:

- ▶ A unique solution if  $A$  has a pivot in every row
- ▶ Infinite number of solutions if  $A$  has a solution and at least one free variable

## Theorem

*The homogeneous equation  $A\mathbf{x} = \mathbf{0}$  has a nontrivial solution if and only if the solution set has at least one free variable.*

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$x_1 + 3x_2 - 2x_3 = 0$$

$$-2x_1 - 6x_2 + 4x_3 = 0$$



## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$x_1 + 3x_2 - 2x_3 = 0$$

$$-2x_1 - 6x_2 + 4x_3 = 0$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right]$$

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$x_1 + 3x_2 - 2x_3 = 0$$

$$-2x_1 - 6x_2 + 4x_3 = 0$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$\begin{aligned}x_1 + 3x_2 - 2x_3 &= 0 \\ -2x_1 - 6x_2 + 4x_3 &= 0\end{aligned}$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This gives us:  $x_1 = -3x_2 + 2x_3$  or as a vector:

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$\begin{aligned}x_1 + 3x_2 - 2x_3 &= 0 \\ -2x_1 - 6x_2 + 4x_3 &= 0\end{aligned}$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This gives us:  $x_1 = -3x_2 + 2x_3$  or as a vector:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$\begin{aligned}x_1 + 3x_2 - 2x_3 &= 0 \\ -2x_1 - 6x_2 + 4x_3 &= 0\end{aligned}$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This gives us:  $x_1 = -3x_2 + 2x_3$  or as a vector:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -3x_2 + 2x_3 \\ x_2 \\ x_3 \end{bmatrix}$$

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$\begin{aligned}x_1 + 3x_2 - 2x_3 &= 0 \\ -2x_1 - 6x_2 + 4x_3 &= 0\end{aligned}$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This gives us:  $x_1 = -3x_2 + 2x_3$  or as a vector:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -3x_2 + 2x_3 \\ x_2 \\ x_3 \end{bmatrix} = x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$$

## Solution Sets of Consistent Systems

Find the solution set to the homogeneous system write in vector form (from lecture 4):

$$\begin{aligned}x_1 + 3x_2 - 2x_3 &= 0 \\ -2x_1 - 6x_2 + 4x_3 &= 0\end{aligned}$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ -2 & -6 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

This gives us:  $x_1 = -3x_2 + 2x_3$  or as a vector:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -3x_2 + 2x_3 \\ x_2 \\ x_3 \end{bmatrix} = x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} = s \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$



## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right]$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 2 & 6 & 10 \end{array} \right]$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 2 & 6 & 10 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 2 & 6 & 10 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & -2 & -11 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Which can be written as

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 2 & 6 & 10 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & -2 & -11 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Which can be written as

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -11 + 2x_3 \\ 5 - 3x_3 \\ x_3 \end{bmatrix}$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 2 & 6 & 10 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & -2 & -11 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Which can be written as

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -11 + 2x_3 \\ 5 - 3x_3 \\ x_3 \end{bmatrix} = \begin{bmatrix} -11 \\ 5 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix}$$

## More Practice!

Find the solution set to the **nonhomogeneous** system

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ -2 & -4 & -8 & 2 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 2 & 6 & 10 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 7 & 4 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & -2 & -11 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Which can be written as

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -11 + 2x_3 \\ 5 - 3x_3 \\ x_3 \end{bmatrix} = \begin{bmatrix} -11 \\ 5 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} = \mathbf{p} + t\mathbf{v} (t \in \mathbb{R})$$

## Comparing Homogeneous and Nonhomogeneous Solutions

$$x_1 + 3x_2 + 7x_3 = 0$$

$$x_2 + 3x_3 = 0$$

$$-2x_1 - 4x_2 - 8x_3 = 0$$

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$



# Comparing Homogeneous and Nonhomogeneous Solutions

$$x_1 + 3x_2 + 7x_3 = 0$$

$$x_2 + 3x_3 = 0$$

$$-2x_1 - 4x_2 - 8x_3 = 0$$

$$\mathbf{x} = x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} = t\mathbf{v} (t \in \mathbb{R})$$

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\mathbf{x} = \begin{bmatrix} -11 \\ 5 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} = \mathbf{p} + t\mathbf{v} (t \in \mathbb{R})$$

# Comparing Homogeneous and Nonhomogeneous Solutions

$$x_1 + 3x_2 + 7x_3 = 0$$

$$x_2 + 3x_3 = 0$$

$$-2x_1 - 4x_2 - 8x_3 = 0$$


$$\mathbf{x} = x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} = t\mathbf{v} \quad (t \in \mathbb{R})$$

$$x_1 + 3x_2 + 7x_3 = 4$$

$$x_2 + 3x_3 = 5$$

$$-2x_1 - 4x_2 - 8x_3 = 2$$

$$\mathbf{x} = \begin{bmatrix} -11 \\ 5 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} = \mathbf{p} + t\mathbf{v} \quad (t \in \mathbb{R})$$



# Infinite Solutions from Homogeneous System

## Theorem

Let  $A\mathbf{x} = \mathbf{b}$  be a consistent system with solution  $\mathbf{x} = \mathbf{p}$ . Then there exists a solution set of vectors of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}.$$

Where  $t\mathbf{v}$  is a solution to the *homogeneous* system  $A\mathbf{x} = \mathbf{0}$ .

# Infinite Solutions from Homogeneous System

## Theorem

Let  $A\mathbf{x} = \mathbf{b}$  be a consistent system with solution  $\mathbf{x} = \mathbf{p}$ . Then there exists a solution set of vectors of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}.$$

Where  $t\mathbf{v}$  is a solution to the *homogeneous* system  $A\mathbf{x} = \mathbf{0}$ .

## Proof.

We will show this by showing  $A\mathbf{x} = A(\mathbf{p} + t\mathbf{v}) = \mathbf{b}$ .

# Infinite Solutions from Homogeneous System

## Theorem

Let  $A\mathbf{x} = \mathbf{b}$  be a consistent system with solution  $\mathbf{x} = \mathbf{p}$ . Then there exists a solution set of vectors of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}.$$

Where  $t\mathbf{v}$  is a solution to the *homogeneous* system  $A\mathbf{x} = \mathbf{0}$ .

## Proof.

We will show this by showing  $A\mathbf{x} = A(\mathbf{p} + t\mathbf{v}) = \mathbf{b}$ .

$$A\mathbf{x} = A(\mathbf{p} + t\mathbf{v})$$

# Infinite Solutions from Homogeneous System

## Theorem

Let  $A\mathbf{x} = \mathbf{b}$  be a consistent system with solution  $\mathbf{x} = \mathbf{p}$ . Then there exists a solution set of vectors of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}.$$

Where  $t\mathbf{v}$  is a solution to the *homogeneous* system  $A\mathbf{x} = \mathbf{0}$ .

## Proof.

We will show this by showing  $A\mathbf{x} = A(\mathbf{p} + t\mathbf{v}) = \mathbf{b}$ .

$$\begin{aligned} A\mathbf{x} &= A(\mathbf{p} + t\mathbf{v}) \\ &= A\mathbf{p} + A(t\mathbf{v}) \end{aligned}$$

# Infinite Solutions from Homogeneous System

## Theorem

Let  $A\mathbf{x} = \mathbf{b}$  be a consistent system with solution  $\mathbf{x} = \mathbf{p}$ . Then there exists a solution set of vectors of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}.$$

Where  $t\mathbf{v}$  is a solution to the *homogeneous* system  $A\mathbf{x} = \mathbf{0}$ .

## Proof.

We will show this by showing  $A\mathbf{x} = A(\mathbf{p} + t\mathbf{v}) = \mathbf{b}$ .

$$\begin{aligned} A\mathbf{x} &= A(\mathbf{p} + t\mathbf{v}) \\ &= A\mathbf{p} + A(t\mathbf{v}) \\ &= \mathbf{b} + \mathbf{0} \end{aligned}$$

# Infinite Solutions from Homogeneous System

## Theorem

Let  $A\mathbf{x} = \mathbf{b}$  be a consistent system with solution  $\mathbf{x} = \mathbf{p}$ . Then there exists a solution set of vectors of the form

$$\mathbf{x} = \mathbf{p} + t\mathbf{v}.$$

Where  $t\mathbf{v}$  is a solution to the *homogeneous* system  $A\mathbf{x} = \mathbf{0}$ .

## Proof.

We will show this by showing  $A\mathbf{x} = A(\mathbf{p} + t\mathbf{v}) = \mathbf{b}$ .

$$\begin{aligned} A\mathbf{x} &= A(\mathbf{p} + t\mathbf{v}) \\ &= A\mathbf{p} + A(t\mathbf{v}) \\ &= \mathbf{b} + \mathbf{0} \\ &= \mathbf{b} \end{aligned}$$

