

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible

# Theorem Statement

## Theorem

*Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .*

- 1.  $A$  is invertible*
- 2.  $A$  has a pivot in every row*
- 3.  $A$  has a pivot in every column*

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
6. The columns of  $A$  are linearly independent

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
6. The columns of  $A$  are linearly independent
7. The columns of  $A$  span all of  $\mathbb{R}^n$ .

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
6. The columns of  $A$  are linearly independent
7. The columns of  $A$  span all of  $\mathbb{R}^n$ .
8.  $T$  is invertible

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
6. The columns of  $A$  are linearly independent
7. The columns of  $A$  span all of  $\mathbb{R}^n$ .
8.  $T$  is invertible
9.  $T$  is injective



# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
6. The columns of  $A$  are linearly independent
7. The columns of  $A$  span all of  $\mathbb{R}^n$ .
8.  $T$  is invertible
9.  $T$  is injective
10.  $T$  is surjective

# Theorem Statement

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a linear transformation with associated matrix  $A \in \mathbb{R}^{n \times n}$ .

1.  $A$  is invertible
2.  $A$  has a pivot in every row
3.  $A$  has a pivot in every column
4. The system  $A\mathbf{x} = \mathbf{b}$  has a unique solution for all  $\mathbf{b} \in \mathbb{R}^n$
5.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution
6. The columns of  $A$  are linearly independent
7. The columns of  $A$  span all of  $\mathbb{R}^n$ .
8.  $T$  is invertible
9.  $T$  is injective
10.  $T$  is surjective

We have the tools to prove all these, but really, it's just a summary of all the linear transformation and matrix algebra we've done so far!

# One-sided inverse Implies the Other!

## Theorem

*Let  $A \in \mathbb{R}^{n \times n}$ . If there is some  $B \in \mathbb{R}^{n \times n}$  such that*

$$BA = I_n \text{ or } AB = I_n$$

*Then,  $A$  is invertible and  $A^{-1} = B$ .*

$AB = I_n$  case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

$AB = I_n$  case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

$AB = I_n$  case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b}$$

## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b}$$



## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b} = A(B\mathbf{b})$$

## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b} = A(B\mathbf{b}) = T(B\mathbf{b})$$

$AB = I_n$  case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b} = A(B\mathbf{b}) = T(B\mathbf{b})$$

Since we have an input,  $\mathbf{x} = B\mathbf{b}$  such that  $T(\mathbf{x}) = \mathbf{b}$ ,

## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b} = A(B\mathbf{b}) = T(B\mathbf{b})$$

Since we have an input,  $\mathbf{x} = B\mathbf{b}$  such that  $T(\mathbf{x}) = \mathbf{b}$ ,

We know that  $T$  is surjective, so by above,  $A$  is invertible.

## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b} = A(B\mathbf{b}) = T(B\mathbf{b})$$

Since we have an input,  $\mathbf{x} = B\mathbf{b}$  such that  $T(\mathbf{x}) = \mathbf{b}$ ,

We know that  $T$  is surjective, so by above,  $A$  is invertible.

Also, see that

## $AB = I_n$ case

Proof.

First, let  $B \in \mathbb{R}^{n \times n}$  such that  $AB = I_n$ .

We will show that  $T$  is surjective by showing how we can get any  $\mathbf{b} \in \mathbb{R}^n$ .

Let  $\mathbf{b} \in \mathbb{R}^n$ . See that

$$\mathbf{b} = I_n \mathbf{b} = AB\mathbf{b} = A(B\mathbf{b}) = T(B\mathbf{b})$$

Since we have an input,  $\mathbf{x} = B\mathbf{b}$  such that  $T(\mathbf{x}) = \mathbf{b}$ ,

We know that  $T$  is surjective, so by above,  $A$  is invertible.

Also, see that



$BA = I_n$  case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.



## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ .

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$A\mathbf{x} = \mathbf{0}$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \end{aligned}$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$A\mathbf{x} = \mathbf{0}$$

$$BA\mathbf{x} = B\mathbf{0}$$

$$I_n\mathbf{x} = \mathbf{0}$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$A\mathbf{x} = \mathbf{0}$$

$$BA\mathbf{x} = B\mathbf{0}$$

$$I_n\mathbf{x} = \mathbf{0}$$

$$\mathbf{x} = \mathbf{0}$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !, which means  $A$  is invertible.



## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !, which means  $A$  is invertible.

Next, we show  $B = A^{-1}$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !, which means  $A$  is invertible.

Next, we show  $B = A^{-1}$

$$A^{-1} = I_n A^{-1}$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !, which means  $A$  is invertible.

Next, we show  $B = A^{-1}$

$$A^{-1} = I_n A^{-1} = BAA^{-1}$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !, which means  $A$  is invertible.

Next, we show  $B = A^{-1}$

$$A^{-1} = I_n A^{-1} = BAA^{-1} = BI_n$$

## $BA = I_n$ case

Proof.

Let  $B \in \mathbb{R}^{n \times n}$  such that  $BA = I_n$ .

We will show that the system  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.

Assume that  $\mathbf{x} \in \mathbb{R}^n$  such that  $A\mathbf{x} = \mathbf{0}$ . See that

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ BA\mathbf{x} &= B\mathbf{0} \\ I_n\mathbf{x} &= \mathbf{0} \\ \mathbf{x} &= \mathbf{0} \end{aligned}$$

So, if  $A\mathbf{x} = \mathbf{0}$ , then  $\mathbf{x}$  must be  $\mathbf{0}$ !, which means  $A$  is invertible.

Next, we show  $B = A^{-1}$

$$A^{-1} = I_n A^{-1} = BAA^{-1} = BI_n = B$$

