

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:

$$\mathbf{u} + \mathbf{v} \in V$$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$
3. Vector addition is associative:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:



# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:

$$\mathbf{u} + \mathbf{v} \in V$$

2. Vector addition is commutative:

$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$$

3. Vector addition is associative:

$$(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$$

4. Additive Identity:

There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:

$$\mathbf{u} + \mathbf{v} \in V$$

2. Vector addition is commutative:

$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$$

3. Vector addition is associative:

$$(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$$

4. Additive Identity:

There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$

5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some

$$-\mathbf{u} \in V \text{ such that } \mathbf{u} + (-\mathbf{u}) = \mathbf{0}$$

6.  $V$  is closed under scalar multiplication:

$$c\mathbf{v} \in V.$$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8. Scalar multiplication distributes over scalar addition:



# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8. Scalar multiplication distributes over scalar addition:  
 $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8. Scalar multiplication distributes over scalar addition:  
 $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$
9. Scalar multiplication is associative:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8. Scalar multiplication distributes over scalar addition:  
 $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$
9. Scalar multiplication is associative:  
 $c(d\mathbf{v}) = (cd)\mathbf{v}$

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8. Scalar multiplication distributes over scalar addition:  
 $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$
9. Scalar multiplication is associative:  
 $c(d\mathbf{v}) = (cd)\mathbf{v}$
10. Multiplicative Identity:

# What is a Vector Space?

A **Vector Space** is a set  $V$  that contains our *vectors*, a set  $F$  that contains our scalars with a **vector addition** operation and **scalar multiplication** operation where the following properties are true for every  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$  and  $c, d \in F$ .

1.  $V$  is closed under addition:  
 $\mathbf{u} + \mathbf{v} \in V$
2. Vector addition is commutative:  
 $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
3. Vector addition is associative:  
 $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. Additive Identity:  
There exists some  $\mathbf{0} \in V$  where  $\mathbf{v} + \mathbf{0} = \mathbf{v}$
5. Additive Inverse: for each  $\mathbf{u} \in V$ , there is some  $-\mathbf{u} \in V$  such that  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6.  $V$  is closed under scalar multiplication:  
 $c\mathbf{v} \in V$ .
7. Scalar multiplication distributes over vector addition:  
 $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ .
8. Scalar multiplication distributes over scalar addition:  
 $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$
9. Scalar multiplication is associative:  
 $c(d\mathbf{v}) = (cd)\mathbf{v}$
10. Multiplicative Identity:  
There exists some  $1 \in F$  such that  $1\mathbf{u} = \mathbf{u}$ .

# What are some examples?

$$V = \mathbb{R}^n, F = \mathbb{R}.$$

## What are some examples?

$V = \mathbb{R}^n, F = \mathbb{R}$ . See Slide 8 of Lecture slide 3 for properties 2-5 and 7-10.

For properties 1 and 2, we have the definitions of vector addition and scalar multiplication that guarantees this!

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.



## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

$$\begin{aligned} 1. (a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = \\ (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0) \end{aligned}$$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$
3.  $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$
3.  $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$
4.  $\mathbf{0} = 0x^2 + 0x + 0$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$
3.  $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$
4.  $\mathbf{0} = 0x^2 + 0x + 0$
5.  $-f(x) = -a_2x^2 - a_1x - a_0$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

- $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
- $f(x) + g(x) = g(x) + f(x)$
- $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$
- $\mathbf{0} = 0x^2 + 0x + 0$
- $-f(x) = -a_2x^2 - a_1x - a_0$
- $cf(x) = ca_2x^2 + ca_1x + ca_0$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

$$1. (a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$$

$$2. f(x) + g(x) = g(x) + f(x)$$

$$3. (f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$$

$$4. \mathbf{0} = 0x^2 + 0x + 0$$

$$5. -f(x) = -a_2x^2 - a_1x - a_0$$

$$6. cf(x) = ca_2x^2 + ca_1x + ca_0$$

$$7. c(f(x) + g(x)) = cf(x) + cg(x)$$



## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$
3.  $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$
4.  $\mathbf{0} = 0x^2 + 0x + 0$
5.  $-f(x) = -a_2x^2 - a_1x - a_0$
6.  $cf(x) = ca_2x^2 + ca_1x + ca_0$
7.  $c(f(x) + g(x)) = cf(x) + cg(x)$
8.  $(c + d)f(x) = cf(x) + df(x)$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$
3.  $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$
4.  $\mathbf{0} = 0x^2 + 0x + 0$
5.  $-f(x) = -a_2x^2 - a_1x - a_0$
6.  $cf(x) = ca_2x^2 + ca_1x + ca_0$
7.  $c(f(x) + g(x)) = cf(x) + cg(x)$
8.  $(c + d)f(x) = cf(x) + df(x)$
9.  $c(df(x)) = (cd)f(x)$

## More Examples

$V = \mathcal{P}_2$  is the set of all polynomials with real coefficients of degree 2 or less.

$F = \mathbb{R}$ , with the operations we'd expect.

1.  $(a_2x^2 + a_1x + a_0) + (b_2x^2 + b_1x + b_0) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0)$
2.  $f(x) + g(x) = g(x) + f(x)$
3.  $(f(x) + g(x)) + h(x) = f(x) + (g(x) + h(x))$
4.  $\mathbf{0} = 0x^2 + 0x + 0$
5.  $-f(x) = -a_2x^2 - a_1x - a_0$
6.  $cf(x) = ca_2x^2 + ca_1x + ca_0$
7.  $c(f(x) + g(x)) = cf(x) + cg(x)$
8.  $(c + d)f(x) = cf(x) + df(x)$
9.  $c(df(x)) = (cd)f(x)$
10.  $1f(x) = f(x)$

## Is this a Vector Space?

$V = \mathbb{R}^3, F = \mathbb{R}$  using standard scalar multiplication but  $\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_2 \\ u_2 + v_2 \\ u_3 + v_3 \end{bmatrix}$

# Uniqueness of $\mathbf{0}$

## Theorem

*If  $V$  is a vector space, then the  $\mathbf{0}$  element is unique*

# Uniqueness of $\mathbf{0}$

## Theorem

*If  $V$  is a vector space, then the  $\mathbf{0}$  element is unique*

## Proof.

Let  $\mathbf{w} \in V$  such that for every  $\mathbf{u} \in V$  we have

$$\mathbf{w} + \mathbf{u} = \mathbf{u} + \mathbf{w} = \mathbf{u}$$

# Uniqueness of $\mathbf{0}$

## Theorem

*If  $V$  is a vector space, then the  $\mathbf{0}$  element is unique*

## Proof.

Let  $\mathbf{w} \in V$  such that for every  $\mathbf{u} \in V$  we have

$$\mathbf{w} + \mathbf{u} = \mathbf{u} + \mathbf{w} = \mathbf{u}$$

By taking  $\mathbf{u} = \mathbf{0}$ , we have:

# Uniqueness of $\mathbf{0}$

## Theorem

*If  $V$  is a vector space, then the  $\mathbf{0}$  element is unique*

## Proof.

Let  $\mathbf{w} \in V$  such that for every  $\mathbf{u} \in V$  we have

$$\mathbf{w} + \mathbf{u} = \mathbf{u} + \mathbf{w} = \mathbf{u}$$

By taking  $\mathbf{u} = \mathbf{0}$ , we have:

$$\mathbf{0} + \mathbf{w} = \mathbf{0}$$



# Uniqueness of $\mathbf{0}$

## Theorem

*If  $V$  is a vector space, then the  $\mathbf{0}$  element is unique*

## Proof.

Let  $\mathbf{w} \in V$  such that for every  $\mathbf{u} \in V$  we have

$$\mathbf{w} + \mathbf{u} = \mathbf{u} + \mathbf{w} = \mathbf{u}$$

By taking  $\mathbf{u} = \mathbf{0}$ , we have:

$$\mathbf{0} + \mathbf{w} = \mathbf{0}$$

$$\mathbf{w} + \mathbf{0} = \mathbf{w}$$

# Uniqueness of $\mathbf{0}$

## Theorem

*If  $V$  is a vector space, then the  $\mathbf{0}$  element is unique*

## Proof.

Let  $\mathbf{w} \in V$  such that for every  $\mathbf{u} \in V$  we have

$$\mathbf{w} + \mathbf{u} = \mathbf{u} + \mathbf{w} = \mathbf{u}$$

By taking  $\mathbf{u} = \mathbf{0}$ , we have:

$$\mathbf{0} + \mathbf{w} = \mathbf{0}$$

$$\mathbf{w} + \mathbf{0} = \mathbf{w}$$

Thus, we see that  $\mathbf{w} = \mathbf{0}$



# Uniqueness of Additive Inverse

## Theorem

*If  $V$  is a vector space, then for every  $\mathbf{u} \in V$ , we have that  $-\mathbf{u}$  is unique.*

# Uniqueness of Additive Inverse

## Theorem

*If  $V$  is a vector space, then for every  $\mathbf{u} \in V$ , we have that  $-\mathbf{u}$  is unique.*

## Proof.

Let  $\mathbf{u} \in V$ , and suppose there are two additive identities  $\mathbf{1E}$  that  $-\mathbf{u}_1, -\mathbf{u}_2 \in V$  such that  $\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} = \mathbf{u} + (-\mathbf{u}_2)$  See that

# Uniqueness of Additive Inverse

## Theorem

*If  $V$  is a vector space, then for every  $\mathbf{u} \in V$ , we have that  $-\mathbf{u}$  is unique.*

## Proof.

Let  $\mathbf{u} \in V$ , and suppose there are two additive identities  $\mathbf{0}$  that  $-\mathbf{u}_1, -\mathbf{u}_2 \in V$  such that  $\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} = \mathbf{u} + (-\mathbf{u}_2)$ . See that

$$\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} \rightarrow -\mathbf{u}_2 + (\mathbf{u} + (-\mathbf{u}_1)) = -\mathbf{u}_2$$

# Uniqueness of Additive Inverse

## Theorem

*If  $V$  is a vector space, then for every  $\mathbf{u} \in V$ , we have that  $-\mathbf{u}$  is unique.*

## Proof.

Let  $\mathbf{u} \in V$ , and suppose there are two additive identities  $\mathbf{0}$  that  $-\mathbf{u}_1, -\mathbf{u}_2 \in V$  such that  $\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} = \mathbf{u} + (-\mathbf{u}_2)$ . See that

$$\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} \rightarrow -\mathbf{u}_2 + (\mathbf{u} + (-\mathbf{u}_1)) = -\mathbf{u}_2 \rightarrow (-\mathbf{u}_2 + \mathbf{u}) + -\mathbf{u}_1 = -\mathbf{u}_2$$

# Uniqueness of Additive Inverse

## Theorem

*If  $V$  is a vector space, then for every  $\mathbf{u} \in V$ , we have that  $-\mathbf{u}$  is unique.*

## Proof.

Let  $\mathbf{u} \in V$ , and suppose there are two additive identities  $\mathbf{0}$  that  $-\mathbf{u}_1, -\mathbf{u}_2 \in V$  such that  $\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} = \mathbf{u} + (-\mathbf{u}_2)$ . See that

$$\mathbf{u} + (-\mathbf{u}_1) = \mathbf{0} \rightarrow -\mathbf{u}_2 + (\mathbf{u} + (-\mathbf{u}_1)) = -\mathbf{u}_2 \rightarrow (-\mathbf{u}_2 + \mathbf{u}) + -\mathbf{u}_1 = -\mathbf{u}_2$$

$$\rightarrow -\mathbf{u}_1 = -\mathbf{u}_2$$



## Vector Space Practice

Work with your neighbors to determine if the following spaces are vector spaces

$V = \mathbb{R}^3$ ,  $F = \mathbb{R}$  with the usual vector addition

and  $c\mathbf{u} = \begin{bmatrix} -cu_1 \\ -cu_2 \\ -cu_3 \end{bmatrix}$ .

$V = \mathbb{R}^{3 \times 3}$ ,  $F = \mathbb{R}$  with the standard operations.



# Vector Subspaces

## Definition

A subspace of a vector space  $V$  is a subset  $H$  of  $V$  ( $H \subseteq V$ ) that has the following properties (using the same vector addition, scalar multiplication, and  $F$ )

# Vector Subspaces

## Definition

A subspace of a vector space  $V$  is a subset  $H$  of  $V$  ( $H \subseteq V$ ) that has the following properties (using the same vector addition, scalar multiplication, and  $F$ )

1. The  $\mathbf{0}$  from  $V$  is in  $H$ .

# Vector Subspaces

## Definition

A subspace of a vector space  $V$  is a subset  $H$  of  $V$  ( $H \subseteq V$ ) that has the following properties (using the same vector addition, scalar multiplication, and  $F$ )

1. The  $\mathbf{0}$  from  $V$  is in  $H$ .
2.  $H$  is closed under vector addition: for each  $\mathbf{u}, \mathbf{v} \in H$ , we have  $\mathbf{u} + \mathbf{v} \in H$ .

# Vector Subspaces

## Definition

A subspace of a vector space  $V$  is a subset  $H$  of  $V$  ( $H \subseteq V$ ) that has the following properties (using the same vector addition, scalar multiplication, and  $F$ )

1. The  $\mathbf{0}$  from  $V$  is in  $H$ .
2.  $H$  is closed under vector addition: for each  $\mathbf{u}, \mathbf{v} \in H$ , we have  $\mathbf{u} + \mathbf{v} \in H$ .
3.  $H$  is closed under scalar multiplication: for each  $c \in F$  and  $\mathbf{v} \in H$ , we have  $c\mathbf{v} \in H$ .

## Is it a Subspace?

Determine with your neighbors if each of the following sets are subspaces of  $V = \mathbb{R}^3$ .

$$H = \left\{ \mathbf{v} \in \mathbb{R}^3 \mid \mathbf{v} = \begin{bmatrix} 3a + b \\ a + 5 \\ 2a - 5b \end{bmatrix} \text{ for } a, b \in \mathbb{R} \right\} \qquad H = \left\{ \mathbf{v} \in \mathbb{R}^3 \mid \mathbf{v} = \begin{bmatrix} a \\ 0 \\ b \end{bmatrix} \text{ for } a, b \in \mathbb{R} \right\}$$

# Spanning Sets and Subspaces

Let  $\mathbf{v}_1, \dots, \mathbf{v}_p$  denote a set of  $p$  vectors in  $V$ . Then  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p)$  is a subspace of  $V$ .

# Spanning Sets and Subspaces

Let  $\mathbf{v}_1, \dots, \mathbf{v}_p$  denote a set of  $p$  vectors in  $V$ . Then  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p)$  is a subspace of  $V$ .

1.  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p)$  is the subspace spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_p$

# Spanning Sets and Subspaces

Let  $\mathbf{v}_1, \dots, \mathbf{v}_p$  denote a set of  $p$  vectors in  $V$ . Then  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p)$  is a subspace of  $V$ .

1.  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p)$  is the subspace spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_p$
2. Given any subspace  $H$  of  $V$ , a spanning set for  $H$  is a set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  of vectors in  $H$  such that  $H = \text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_p)$



## Example

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if so, give a spanning set for  $H$ .

## Example

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if

so, give a spanning set for  $H$ . Show that  $\mathbf{0} \in H$ , Show that we are closed under “vector” addition, and show we are closed under scalar multiplication.

## Example

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if

so, give a spanning set for  $H$ . Show that  $\mathbf{0} \in H$ , Show that we are closed under “vector” addition, and show we are closed under scalar multiplication.

1. Set  $a = b = c = 0$ , clearly  $a + b + c = 0$  and then we have the 0 matrix!

## Example

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if

so, give a spanning set for  $H$ . Show that  $\mathbf{0} \in H$ , Show that we are closed under “vector” addition, and show we are closed under scalar multiplication.

1. Set  $a = b = c = 0$ , clearly  $a + b + c = 0$  and then we have the 0 matrix!
2. Let  $A, B \in H$ . See that

$$A + B = \begin{bmatrix} a_1 + a_2 & 0 & 0 \\ 0 & b_1 + b_2 & 0 \\ 0 & 0 & c_1 + c_2 \end{bmatrix}$$

## Example

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if

so, give a spanning set for  $H$ . Show that  $\mathbf{0} \in H$ , Show that we are closed under “vector” addition, and show we are closed under scalar multiplication.

1. Set  $a = b = c = 0$ , clearly  $a + b + c = 0$  and then we have the 0 matrix!
2. Let  $A, B \in H$ . See that

$$A + B = \begin{bmatrix} a_1 + a_2 & 0 & 0 \\ 0 & b_1 + b_2 & 0 \\ 0 & 0 & c_1 + c_2 \end{bmatrix}$$

$$\text{and } (a_1 + a_2) + (b_1 + b_2) + (c_1 + c_2) = a_1 + b_1 + c_1 + a_2 + b_2 + c_2$$

## Example

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if

so, give a spanning set for  $H$ . Show that  $\mathbf{0} \in H$ , Show that we are closed under “vector” addition, and show we are closed under scalar multiplication.

1. Set  $a = b = c = 0$ , clearly  $a + b + c = 0$  and then we have the 0 matrix!
2. Let  $A, B \in H$ . See that

$$A + B = \begin{bmatrix} a_1 + a_2 & 0 & 0 \\ 0 & b_1 + b_2 & 0 \\ 0 & 0 & c_1 + c_2 \end{bmatrix}$$

and  $(a_1 + a_2) + (b_1 + b_2) + (c_1 + c_2) = a_1 + b_1 + c_1 + a_2 + b_2 + c_2 = 0 + 0 = 0$ , so  $A + B \in H$ .

## Example continued

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if so, give a spanning set for  $H$ .

## Example continued

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if so, give a spanning set for  $H$ .

3. let  $x \in \mathbb{R}$ . See that

$$xA = \begin{bmatrix} xa & 0 & 0 \\ 0 & xb & 0 \\ 0 & 0 & xc \end{bmatrix}$$



## Example continued

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if so, give a spanning set for  $H$ .

3. let  $x \in \mathbb{R}$ . See that

$$xA = \begin{bmatrix} xa & 0 & 0 \\ 0 & xb & 0 \\ 0 & 0 & xc \end{bmatrix}$$

And

$$xa + xb + xc = x(a + b + c)$$

## Example continued

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if so, give a spanning set for  $H$ .

3. let  $x \in \mathbb{R}$ . See that

$$xA = \begin{bmatrix} xa & 0 & 0 \\ 0 & xb & 0 \\ 0 & 0 & xc \end{bmatrix}$$

And

$$xa + xb + xc = x(a + b + c) = x \cdot 0 = 0$$

## Example continued

Determine if  $H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$  is a subspace of  $\mathbb{R}^{3 \times 3}$  and if so, give a spanning set for  $H$ .

3. let  $x \in \mathbb{R}$ . See that

$$xA = \begin{bmatrix} xa & 0 & 0 \\ 0 & xb & 0 \\ 0 & 0 & xc \end{bmatrix}$$

And

$$xa + xb + xc = x(a + b + c) = x \cdot 0 = 0$$

So,  $H$  is a subspace of  $\mathbb{R}^3$ !

## Example continued pt. 2

See that our “vectors” are  $3 \times 3$  matrices, so our spanning set will have these kinds of matrices!

## Example continued pt. 2

See that our “vectors” are  $3 \times 3$  matrices, so our spanning set will have these kinds of matrices! Define

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

## Example continued pt. 2

See that our “vectors” are  $3 \times 3$  matrices, so our spanning set will have these kinds of matrices! Define

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

See that

$$\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) = \{A \in \mathbb{R}^{3 \times 3} \mid A = a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3, \text{ for } a, b, c \in \mathbb{R}\}$$

## Example continued pt. 2

See that our “vectors” are  $3 \times 3$  matrices, so our spanning set will have these kinds of matrices! Define

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

See that

$$\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) = \{A \in \mathbb{R}^{3 \times 3} \mid A = a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3, \text{ for } a, b, c \in \mathbb{R}\}$$

And

## Example continued pt. 3

$$a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3 = a \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} + b \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix} + c \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$



## Example continued pt. 3

$$\begin{aligned} a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3 &= a \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} + b \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix} + c \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2a - b - c & 0 & 0 \\ 0 & -a + 2b - c & 0 \\ 0 & 0 & -a - b + 2c \end{bmatrix} \end{aligned}$$

Where

$$2a - b - c + (-a + 2b - c) + (-a - b + 2c)$$

## Example continued pt. 3

$$\begin{aligned} a\mathbf{v}_1 + b\mathbf{v}_2 + c\mathbf{v}_3 &= a \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} + b \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix} + c \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2a - b - c & 0 & 0 \\ 0 & -a + 2b - c & 0 \\ 0 & 0 & -a - b + 2c \end{bmatrix} \end{aligned}$$

Where

$$2a - b - c + (-a + 2b - c) + (-a - b + 2c) = 0$$

# Basis of a Vector Space

## Definition

**Basis:** A **basis** of a vector space  $V$  is a set of  $v_1, \dots, v_p \in V$  such that

# Basis of a Vector Space

## Definition

**Basis:** A **basis** of a vector space  $V$  is a set of  $v_1, \dots, v_p \in V$  such that

1.  $\text{Span}(v_1, \dots, v_p) = V$

# Basis of a Vector Space

## Definition

**Basis:** A **basis** of a vector space  $V$  is a set of  $v_1, \dots, v_p \in V$  such that

1.  $\text{Span}(v_1, \dots, v_p) = V$
2.  $v_1, \dots, v_p$  are linearly independent

# Basis of a Vector Space

## Definition

**Basis:** A **basis** of a vector space  $V$  is a set of  $v_1, \dots, v_p \in V$  such that

1.  $\text{Span}(v_1, \dots, v_p) = V$
2.  $v_1, \dots, v_p$  are linearly independent

## Example

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

# Basis of a Vector Space

## Definition

**Basis:** A **basis** of a vector space  $V$  is a set of  $v_1, \dots, v_p \in V$  such that

1.  $\text{Span}(v_1, \dots, v_p) = V$
2.  $v_1, \dots, v_p$  are linearly independent

## Example

$$\mathbf{v}_1 = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

is a basis of

$$H = \left\{ A \in \mathbb{R}^{3 \times 3} \mid A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, a + b + c = 0 \right\}$$

# Length of Basis and Dimension of Vector Space

## Theorem

*All bases of a vector space  $V$  have the same number of elements*



# Length of Basis and Dimension of Vector Space

## Theorem

*All bases of a vector space  $V$  have the same number of elements*

## Definition

The **dimension** of a vector space, denoted  $\dim(V)$  is the length of a basis of  $V$ .

# Spanning and Independent List of Correct Size is a Basis

## Theorem

*Let  $V$  be a vector space with  $n = \dim(V)$ . Then, any linearly independent list of  $n$  vectors,  $\mathbf{v}_1, \dots, \mathbf{v}_n \in V$  forms a basis of  $V$ .*

# Spanning and Independent List of Correct Size is a Basis

## Theorem

*Let  $V$  be a vector space with  $n = \dim(V)$ . Then, any linearly independent list of  $n$  vectors,  $\mathbf{v}_1, \dots, \mathbf{v}_n \in V$  forms a basis of  $V$ .*

## Theorem

*Let  $V$  be a vector space with  $n = \dim(V)$ . Then, any spanning of  $n$  vectors,  $\mathbf{v}_1, \dots, \mathbf{v}_n \in V$  such that  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_n) = V$  is also a basis of  $V$ .*