

# Diagonal Matrices

Recall that a diagonal matrix looks something like

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

# Diagonal Matrices

Recall that a diagonal matrix looks something like

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

And we can see that computing  $D\mathbf{x}$  is pretty easy!

$$D\mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

# Diagonal Matrices

Recall that a diagonal matrix looks something like

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

And we can see that computing  $D\mathbf{x}$  is pretty easy!

$$D\mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ 3x_2 \\ 2x_3 \end{bmatrix}$$

# Diagonal Matrices

Recall that a diagonal matrix looks something like

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

And we can see that computing  $D\mathbf{x}$  is pretty easy!

$$D\mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ 3x_2 \\ 2x_3 \end{bmatrix}$$

So, our lives could be easier if we can find a diagonal matrix that a given matrix behaves like!

# Similarity

## Definition

Let  $A, B \in \mathbb{R}^{n \times n}$ . Then we say that  $A$  and  $B$  are **similar** if there is some  $C \in \mathbb{R}^{n \times n}$  such that  $C$  is invertible and

$$A = CBC^{-1}$$

# Similarity

## Definition

Let  $A, B \in \mathbb{R}^{n \times n}$ . Then we say that  $A$  and  $B$  are **similar** if there is some  $C \in \mathbb{R}^{n \times n}$  such that  $C$  is invertible and

$$A = CBC^{-1}$$

Or equivalently without using an inverse

$$AC = CB$$

# Similarity

## Definition

Let  $A, B \in \mathbb{R}^{n \times n}$ . Then we say that  $A$  and  $B$  are **similar** if there is some  $C \in \mathbb{R}^{n \times n}$  such that  $C$  is invertible and

$$A = CBC^{-1}$$

Or equivalently without using an inverse

$$AC = CB$$

## Example

If

$$A = \begin{bmatrix} -12 & 15 \\ -10 & 13 \end{bmatrix}, B = \begin{bmatrix} 3 & 0 \\ 0 & -2 \end{bmatrix}, C = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}$$

Then  $A$  and  $B$  are similar as  $AC = CB$ .

# Computing Large Matrix Powers

One application of similar matrices is that we can compute matrix powers really easy!



# Computing Large Matrix Powers

One application of similar matrices is that we can compute matrix powers really easy! Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}$$

# Computing Large Matrix Powers

One application of similar matrices is that we can compute matrix powers really easy! Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}$$

Then for any  $k \geq 1$  we have that

$$A^k = \underbrace{A \cdots A}_{k \text{ times}}$$

# Computing Large Matrix Powers

One application of similar matrices is that we can compute matrix powers really easy! Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}$$

Then for any  $k \geq 1$  we have that

$$A^k = \underbrace{A \cdots A}_{k \text{ times}} = CB^kC^{-1}$$

# Computing Large Matrix Powers

One application of similar matrices is that we can compute matrix powers really easy! Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}$$

Then for any  $k \geq 1$  we have that

$$A^k = \underbrace{A \cdots A}_{k \text{ times}} = CB^kC^{-1}$$

Which is really easy to compute if  $B$  is diagonal or some other nice structure!

# Similarity Transformation

Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}.$$

# Similarity Transformation

Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}.$$

We sometimes call this a **similarity transformation from  $A$  to  $B$** .

# Similarity Transformation

Let  $A, B, C \in \mathbb{R}^{n \times n}$  such that

$$A = CBC^{-1}.$$

We sometimes call this a **similarity transformation from  $A$  to  $B$** . We will see why this is important in the next few slides!

# Similarity Transformation as a Change of Basis

Let's consider an invertible  $C \in \mathbb{R}^{n \times n}$  with columns denoted  $\mathbf{v}_1, \dots, \mathbf{v}_n$  as follows

$$C = [\mathbf{v}_1 \quad \dots \quad \mathbf{v}_n]$$



# Similarity Transformation as a Change of Basis

Let's consider an invertible  $C \in \mathbb{R}^{n \times n}$  with columns denoted  $\mathbf{v}_1, \dots, \mathbf{v}_n$  as follows

$$C = [\mathbf{v}_1 \quad \dots \quad \mathbf{v}_n]$$

Since  $C$  is invertible,  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly independent!

# Similarity Transformation as a Change of Basis

Let's consider an invertible  $C \in \mathbb{R}^{n \times n}$  with columns denoted  $\mathbf{v}_1, \dots, \mathbf{v}_n$  as follows

$$C = [\mathbf{v}_1 \quad \dots \quad \mathbf{v}_n]$$

Since  $C$  is invertible,  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly independent! So, this means

$$\mathcal{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$$

forms a basis of  $\mathbb{R}^n$ , so we can talk about  $\mathcal{B}$ -Coordinates!

## $\mathcal{B}$ -Coordinates of $\mathbf{x}$

## $\mathcal{B}$ -Coordinates of $\mathbf{x}$

So, if we want to find the  $\mathcal{B}$ -Coordinates of some vector  $\mathbf{x}$  we would get

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \text{ where } \mathbf{x} = c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n$$

## $\mathcal{B}$ -Coordinates of $\mathbf{x}$

So, if we want to find the  $\mathcal{B}$ -Coordinates of some vector  $\mathbf{x}$  we would get

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \text{ where } \mathbf{x} = c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n$$

This means that

$$C [\mathbf{x}]_{\mathcal{B}} = \mathbf{x} \leftrightarrow C^{-1} \mathbf{x} = [\mathbf{x}]_{\mathcal{B}}$$

## $\mathcal{B}$ -Coordinates of $\mathbf{x}$

So, if we want to find the  $\mathcal{B}$ -Coordinates of some vector  $\mathbf{x}$  we would get

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \text{ where } \mathbf{x} = c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n$$

This means that

$$C [\mathbf{x}]_{\mathcal{B}} = \mathbf{x} \leftrightarrow C^{-1} \mathbf{x} = [\mathbf{x}]_{\mathcal{B}}$$

So  $C^{-1}$  takes a vector in the standard basis and converts it to coordinates in the  $\mathcal{B}$  basis.

## $\mathcal{B}$ -Coordinates of $\mathbf{x}$

So, if we want to find the  $\mathcal{B}$ -Coordinates of some vector  $\mathbf{x}$  we would get

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \text{ where } \mathbf{x} = c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n$$

This means that

$$C [\mathbf{x}]_{\mathcal{B}} = \mathbf{x} \leftrightarrow C^{-1} \mathbf{x} = [\mathbf{x}]_{\mathcal{B}}$$

So  $C^{-1}$  takes a vector in the standard basis and converts it to coordinates in the  $\mathcal{B}$  basis. Or, in otherwords, we're finding a basis under which the matrix  $A$  behaves “like”  $B$  does!

## Putting it All Together for Similarity Transformations

Since  $C^{-1}$  takes a vector  $\mathbf{x}$  and computes the  $\mathcal{B}$ -Coordinates of that vector and  $C$  returns it to the standard coordinates, we see that:



## Putting it All Together for Similarity Transformations

Since  $C^{-1}$  takes a vector  $\mathbf{x}$  and computes the  $\mathcal{B}$ -Coordinates of that vector and  $C$  returns it to the standard coordinates, we see that:

$$A\mathbf{x} = CBC^{-1}\mathbf{x}$$

## Putting it All Together for Similarity Transformations

Since  $C^{-1}$  takes a vector  $\mathbf{x}$  and computes the  $\mathcal{B}$ -Coordinates of that vector and  $C$  returns it to the standard coordinates, we see that:

$$A\mathbf{x} = CBC^{-1}\mathbf{x} = C(B(C^{-1}\mathbf{x}))$$

performs the following actions

## Putting it All Together for Similarity Transformations

Since  $C^{-1}$  takes a vector  $\mathbf{x}$  and computes the  $\mathcal{B}$ -Coordinates of that vector and  $C$  returns it to the standard coordinates, we see that:

$$A\mathbf{x} = CBC^{-1}\mathbf{x} = C(B(C^{-1}\mathbf{x}))$$

performs the following actions

1. Computes the  $\mathcal{B}$ -Coordinates of  $\mathbf{x}$

## Putting it All Together for Similarity Transformations

Since  $C^{-1}$  takes a vector  $\mathbf{x}$  and computes the  $\mathcal{B}$ -Coordinates of that vector and  $C$  returns it to the standard coordinates, we see that:

$$A\mathbf{x} = CBC^{-1}\mathbf{x} = C(B(C^{-1}\mathbf{x}))$$

performs the following actions

1. Computes the  $\mathcal{B}$ -Coordinates of  $\mathbf{x}$
2. Transforms  $[\mathbf{x}]_{\mathcal{B}}$  via  $B$ . IE  $[\mathbf{y}]_{\mathcal{B}} = B[\mathbf{x}]_{\mathcal{B}}$

## Putting it All Together for Similarity Transformations

Since  $C^{-1}$  takes a vector  $\mathbf{x}$  and computes the  $\mathcal{B}$ -Coordinates of that vector and  $C$  returns it to the standard coordinates, we see that:

$$A\mathbf{x} = CBC^{-1}\mathbf{x} = C(B(C^{-1}\mathbf{x}))$$

performs the following actions

1. Computes the  $\mathcal{B}$ -Coordinates of  $\mathbf{x}$
2. Transforms  $[\mathbf{x}]_{\mathcal{B}}$  via  $B$ . IE  $[\mathbf{y}]_{\mathcal{B}} = B[\mathbf{x}]_{\mathcal{B}}$
3. Returns  $[\mathbf{y}]_{\mathcal{B}}$  to the standard coordinates

# Similarity Transformations Preserve Eigenvalues!

The final properties that we will discuss are how eigenpairs behave under similarity transformations

# Similarity Transformations Preserve Eigenvalues!

The final properties that we will discuss are how eigenpairs behave under similarity transformations. We first claim that the eigenvalues are preserved. See that

$$\det(A - \lambda I) = \det(CBC^{-1} - \lambda CC^{-1})$$

# Similarity Transformations Preserve Eigenvalues!

The final properties that we will discuss are how eigenpairs behave under similarity transformations. We first claim that the eigenvalues are preserved. See that

$$\det(A - \lambda I) = \det(CBC^{-1} - \lambda CC^{-1}) = \det(C(B - \lambda I)C^{-1})$$



# Similarity Transformations Preserve Eigenvalues!

The final properties that we will discuss are how eigenpairs behave under similarity transformations. We first claim that the eigenvalues are preserved. See that

$$\begin{aligned}\det(A - \lambda I) &= \det(CBC^{-1} - \lambda CC^{-1}) = \det(C(B - \lambda I)C^{-1}) \\ &= \det(C) \det(B - \lambda I) \det(C^{-1})\end{aligned}$$

# Similarity Transformations Preserve Eigenvalues!

The final properties that we will discuss are how eigenpairs behave under similarity transformations. We first claim that the eigenvalues are preserved. See that

$$\begin{aligned}\det(A - \lambda I) &= \det(CBC^{-1} - \lambda CC^{-1}) = \det(C(B - \lambda I)C^{-1}) \\ &= \det(C) \det(B - \lambda I) \det(C^{-1}) = \det(B - \lambda I)\end{aligned}$$

# Similarity Transformations Preserve Eigenvalues!

The final properties that we will discuss are how eigenpairs behave under similarity transformations. We first claim that the eigenvalues are preserved. See that

$$\begin{aligned}\det(A - \lambda I) &= \det(CBC^{-1} - \lambda CC^{-1}) = \det(C(B - \lambda I)C^{-1}) \\ &= \det(C) \det(B - \lambda I) \det(C^{-1}) = \det(B - \lambda I)\end{aligned}$$

So, any value  $\lambda$  that makes  $\det(A - \lambda I) = 0$  will necessarily make  $\det(B - \lambda I) = 0$ , so the eigenvalues must be the same!

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .



## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

$$AC\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

$$AC\mathbf{v} = (CBC^{-1})C\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

$$AC\mathbf{v} = (CBC^{-1})C\mathbf{v} = CB\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

$$AC\mathbf{v} = (CBC^{-1})C\mathbf{v} = CB\mathbf{v} = \lambda C\mathbf{v}$$

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

$$AC\mathbf{v} = (CBC^{-1})C\mathbf{v} = CB\mathbf{v} = \lambda C\mathbf{v}$$

So  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ !

## Similarity Transformations Also Transform Eigenvectors

We claim that if  $(\lambda, \mathbf{v})$  is an eigenpair of  $A$ , then  $(\lambda, C^{-1}\mathbf{v})$  is an eigenvector of  $B$ . See that

$$BC^{-1}\mathbf{v} = (C^{-1}C)BC^{-1}\mathbf{v} = C^{-1}A\mathbf{v} = \lambda C^{-1}\mathbf{v}$$

So  $(\lambda, C^{-1}\mathbf{v})$  is an eigenpair of  $B$ .

Similarly if  $(\lambda, \mathbf{v})$  is an eigenpair of  $B$  then  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ . See that

$$AC\mathbf{v} = (CBC^{-1})C\mathbf{v} = CB\mathbf{v} = \lambda C\mathbf{v}$$

So  $(\lambda, C\mathbf{v})$  is an eigenpair of  $A$ !

This means we can think of eigenvectors of  $A$  and  $B$  as the same objects just with different coordinates!

# Geometry of Similarity Transformations

See Section 5.3 of textbook. The images there are much better than what I will come up with



## Some More Statements About Similarity

Some more properties that are nice to know are as follows

## Some More Statements About Similarity

Some more properties that are nice to know are as follows

1. The only matrix similar to  $I_n$  is  $I_n$  itself

## Some More Statements About Similarity

Some more properties that are nice to know are as follows

1. The only matrix similar to  $I_n$  is  $I_n$  itself
2. The only matrix similar to  $0_{n \times n}$  is  $0_{n \times n}$

## Some More Statements About Similarity

Some more properties that are nice to know are as follows

1. The only matrix similar to  $I_n$  is  $I_n$  itself
2. The only matrix similar to  $0_{n \times n}$  is  $0_{n \times n}$
3. Similarity has nothing to do with row equivalence