

Advanced Statistical Methodology and Psychological Flourishing

Joshua N. Pritikin

Virginia Institute for Psychiatric and Behavioral Genetics
Virginia Commonwealth University

Nov 2019



Acknowledgment

Some mentors and collaborators


- ▶ Mike Neale
 - ▶ Tim Brick (Penn State Univ)
 - ▶ Steven Boker (Univ of Virginia)
 - ▶ Karen Schmidt (Univ of Virginia)
 - ▶ OpenMx development team



A detailed illustration of a guinea pig from a side-on perspective. The guinea pig has a light grey body with a dark grey patch on its back and a dark grey mask-like patch around its eyes. It has pink ears, a pink nose, and pink paws. Its long whiskers are clearly visible.


Funded in part by NIDA R25 DA-26119

Number of estimated parameters




Agenda

- ▶ OpenMx
 - ▶ Heritability of cortical thickness
 - ▶ Confidential analyses
 - ▶ Psychological wellbeing
 - ▶ Flow & physical activities
 - ▶ Understanding human wellbeing (*if time permits*)



Agenda

- ▶ OpenMx
 - ▶ Heritability of cortical thickness
 - ▶ Confidential analyses
 - ▶ Psychological wellbeing
 - ▶ Flow & physical activities
 - ▶ Understanding human wellbeing (*if time*)



OpenMx

Extended structural equation modelling¹



Written in R & C++

Open source

11 authors over 12 years

Team meeting Friday 9am-11 (subject to change)

Comparable to lavaan, Mplus, EQS, LISREL, SPSS, etc...

¹Neale et al. (2016)



Some of my contributions



Modern test theory/Item response theory²

Many-level multilevel structural equation modeling³

Multivariate normal maximum likelihood with both ordinal and continuous variables, and data missing at random⁴

Likelihood-based confidence intervals for a parameter with an upper or lower bound⁵

Main source of publications

²Pritikin (2017a); Pritikin and Schmidt (2016); Pritikin, Hunter, and Boker (2015); Pritikin (2016)

³Pritikin, Hunter, von Oertzen, Brick, and Boker (2017)

⁴Pritikin, Brick, and Neale (2018)


⁵Pritikin, Rappaport, and Neale (2017)

5 years, 500k lines of code

Apr 21, 2013 – Nov 20, 2019


Contributions: **Commits** ▾

Contributions to master, excluding merge commits






Latent growth curve + autoregressive model



Factor model, F scales loadings



Potential student projects


Some ideas:

- ▶ Higher moments to fit product models
 - ▶ Automatic conversion to Bayesian
 - ▶ Copula for diverse link functions
 - ▶ Analogue of quantile regression



Agenda

- ▶ OpenMx
 - ▶ Heritability of cortical thickness
 - ▶ Confidential analyses
 - ▶ Psychological wellbeing
 - ▶ Flow & physical activities
 - ▶ Understanding human wellbeing (*if time permits*)



Cortical thickness⁶

Application of OpenMx


Recently evolved phenotypes exhibits variability

- ▶ 410 families
 - ▶ 792 normally developing children, adolescents, and young adults
 - ▶ Up to 8 MRI scans performed per individual
 - ▶ Mean interval between scans was 2.4 yrs
 - ▶ 1,748 MRI datasets acquired


⁶Schmitt et al. (2014)




Genetically informative latent growth curve model




Univariate twin model




Latent growth curve



Genetically informative latent growth curve model



Results



Cloud computing

Original 2014 analyses⁷

- ▶ 3.7 GHz Quad Core Mac Pro Desktop computer
 - ▶ 4 months of processing time



Reanalysis⁸

- ▶ Rented 1024 CPUs
 - ▶ Kubernetes
 - ▶ Done in 2 hr
 - ▶ Identical results
 - ▶ Cost: \$200



Recently applied to new data⁹


⁷Schmitt et al. (2014)

⁸Pritikin, Schmitt, and Neale (2019)

⁹Schmitt, Neale, et al. (2019); Schmitt, Raznahan, et al. (2019)

Agenda

- ▶ OpenMx
 - ▶ Heritability of cortical thickness
 - ▶ Confidential analyses
 - ▶ Psychological wellbeing
 - ▶ Flow & physical activities
 - ▶ Understanding human wellbeing (*if time permits*)





Confidential analyses

What if we could keep participant data confidential,
never revealed to researchers,
and still fit statistical models to data
and test hypotheses?



Statistical models

Given

x_i row x of data from person i (1)

θ parameter vector (2)

Full-information likelihood often has the form

$$\sum_{i=1}^I \log L(x_i | \theta) \quad (3)$$

and rows are assumed independent and identically distributed.

Distributed likelihood evaluation (DLE)

$$\begin{aligned} \sum_{i=1}^I \log L(x_i | \theta) &= \log L(x_1 | \theta) \\ &\quad + \log L(x_2 | \theta) + \\ &\quad + \log L(x_3 | \theta) + \\ &\quad + \log L(x_4 | \theta) + \\ &\quad \cdots + \\ &\quad + \log L(x_I | \theta) \end{aligned} \tag{4}$$




Maintained individual data (MID)



Your personal $L(x_i|\theta)$ runs on your smartphone

- ▶ Data remain confidential
 - ▶ Automatic data sharing across experiments
 - ▶ Larger participant pool with more generalizable estimates

Next steps



Proof-of-concept stage¹⁰

Apply to federal health agencies for funding

¹⁰Boker et al. (2015)

Potential student projects


Some ideas:

- ▶ Delineate security and confidentiality trade-offs
 - ▶ Investigate power and inference in the context of a continuously changing sample
 - ▶ Recommend changes to informed consent and IRB processes



Agenda

- ▶ OpenMx
- ▶ Heritability of cortical thickness
- ▶ Confidential analyses
- ▶ Psychological wellbeing
- ▶ Flow & physical activities
- ▶ Understanding human wellbeing (*if time permits*)




History of sophisticated thinking

Not a study, but a research program (to be)

Recent evolution

- ▶ 1.8-1 MYO – Cooking¹¹
 - ▶ 200k-100k BCE – Homo sapiens
 - ▶ 3400-3100 BCE – Written language
 - ▶ 1500-200 BCE – Sophisticated thinking



¹¹Wrangham (2009)



Ouch



Conscious thinking has costs¹²


- ▶ maladaptive rumination
 - ▶ jealousy, guilt
 - ▶ anxiety, excessive planning



12 Leary (2007)

Solutions

Think better (e.g., cognitive behavioral therapy¹³)




In part due to hypofrontality¹⁴

¹³Butler, Chapman, Forman, and Beck (2006)

¹⁴Dietrich (2003)

Solutions

Think better (e.g., cognitive behavioral therapy¹³)



Temporarily cease thinking (to some extent)

- ▶ Selflessness (i.e., dissolution of the self non-self boundary)
- ▶ Timelessness
- ▶ Effortlessness (i.e., spontaneous or involuntary)

In part due to hypofrontality¹⁴

¹³Butler et al. (2006)

¹⁴Dietrich (2003)

Substantive research focus

Non-ordinary states of consciousness
that support

- ▶ psychological well-being
- ▶ optimal functioning
- ▶ mental and physical health
- ▶ fulfillment



Includes: flow, meditation, mindfulness, psychedelic drugs¹⁵

Excludes: dreaming, daydreaming, hypnosis

¹⁵Pollan (2018)



Big challenges

Tricky to induce in the lab

Ephemeral

Prior work has often
underestimated the difficulty¹⁶



¹⁶E.g., Goyal et al. (2014)




Big challenges

... and it's gone



Agenda

- ▶ OpenMx
- ▶ Heritability of cortical thickness
- ▶ Confidential analyses
- ▶ Psychological wellbeing
- ▶ Flow & physical activities
- ▶ Understanding human wellbeing (*if time permits*)



Exploratory survey

Personal factors

- ▶ Psychological¹⁷
- ▶ Physiological¹⁸
- ▶ Neurological¹⁹

What about situational factors?



¹⁷E.g., Jackson, Ford, Kimiecik, and Marsh (1998)

¹⁸E.g., de Manzano, Theorell, Harmat, and Ullén (2010)

¹⁹E.g., Limb and Braun (2008)

Optimal performance

Flow involves²⁰

- ▶ Complete immersion without reflective self-consciousness but with a deep sense of control
- ▶ Preconditions: skills-demands compatibility, clear goals, and immediate feedback

Contexts:

- ▶ Running²¹
- ▶ Jazz improvisation²²
- ▶ Etc...



²⁰Keller and Landhäußer (2012)

²¹Csikszentmihalyi, Latter, and Duranso (2017)

²²Limb and Braun (2008)



Sample item (template)

Participant picks: A, B

How predictable is the action?

- ▶ B is much more predictable than A.
- ▶ B is somewhat more predictable than A.
- ▶ Both offer roughly equal predictability.
- ▶ A is somewhat more predictable than B.
- ▶ A is much more predictable than B.



Sample item

Participant picks: running, golf

How predictable is the action?

- ▶ golf is much more predictable than running.
- ▶ golf is somewhat more predictable than running.
- ▶ Both offer roughly equal predictability.
- ▶ running is somewhat more predictable than golf.
- ▶ running is much more predictable than golf.

20 items inspired by prior research on flow²³

²³Kotler (2014); Sawyer (2007); Wegner (2002)

Birnbaum's dichotomous response model²⁴

$$\pi(1|\alpha, c, \theta) = \frac{1}{1 + \exp(-(αθ + c))} \quad (5)$$

$$\pi(0|\alpha, c, \theta) = 1 - \pi(1|\alpha, c, \theta) \quad (6)$$

where

- ▶ α is the discrimination parameter
- ▶ c is the intercept
- ▶ θ is the latent ability

²⁴Birnbaum (1968)

Build your understanding

Open your laptop and run R

- ▶ `install.packages(c('ifaTools', 'pcFactorStan'))`
- ▶ `library(ifaTools)`
- ▶ `ifaTools::itemModelExplorer()`
- ▶ <https://www.youtube.com/watch?v=KlnSiarko0g>

Adjust for paired comparisons

$$\pi(1|\alpha, c, \theta) = \frac{1}{1 + \exp(-(\alpha\theta + c))} \quad (7)$$

$$\pi(0|\alpha, c, \theta) = 1 - \pi(1|\alpha, c, \theta) \quad (8)$$

where

- ▶ α is the discrimination parameter
- ▶ c is the intercept
- ▶ θ is the latent ability



Adjusted for paired comparisons

$$\pi(1|\alpha, c, \theta_i, \theta_j) = \frac{1}{1 + \exp(-(\alpha(\theta_j - \theta_i) + c))} \quad (9)$$

$$\pi(0|\alpha, c, \theta_i, \theta_j) = 1 - \pi(1|\alpha, c, \theta_i, \theta_j) \quad (10)$$

where

- ▶ α is the discrimination parameter
- ▶ c is the intercept
- ▶ $\theta_{i,j}$ are two latent worths



What about symmetry?

Given,

$$\pi(1|\alpha, \textcolor{blue}{c}, \theta_i, \theta_j) = \frac{1}{1 + \exp(-(\alpha(\theta_j - \theta_i) + \textcolor{blue}{c}))} \quad (11)$$

If

$$\pi(1|\alpha, c, \theta_1, \theta_2) = 1 - \pi(1|\alpha, c, \theta_2, \theta_1) \quad (12)$$

then

$$c = 0. \quad (13)$$



Paired comparison response function²⁵

$$\pi(1|\alpha, \theta_i, \theta_j) = \frac{1}{1 + \exp(-\alpha(\theta_j - \theta_i))} \quad (14)$$

$$\pi(0|\alpha, \theta_i, \theta_j) = 1 - \pi(1|\alpha, \theta_i, \theta_j) \quad (15)$$

where

- ▶ α is the discrimination parameter
 - ▶ $\theta_{i,j}$ are two latent worths

²⁵Bradley and Terry (1952)

Build your understanding

In R,

- ```
► library(pcFactorStan)
► pcFactorStan::itemModelExplorer()
► https://www.youtube.com/watch?v=6Q31lPByM4Q
```

# Some differences

Dichotomous response model

- ▶ Sum score
- ▶ Judge differences (latent ability)

Paired comparison model

- ▶ No sum score
- ▶ Judge differences regarded as measurement error

## Item 1/10: Predictability

Participant picks: running , golf

How predictable is the action?

- ▶ golf is much more predictable than running.
  - ▶ golf is somewhat more predictable than running.
  - ▶ Both offer roughly equal predictability.
  - ▶ running is somewhat more predictable than golf.
  - ▶ running is much more predictable than golf.



## Item 2/10: Skill

Participant picks: running , golf

How much skill is required?

- ▶ running requires much more skill than golf.
  - ▶ running requires somewhat more skill than golf.
  - ▶ Both require roughly equal skill.
  - ▶ golf requires somewhat more skill than running.
  - ▶ golf requires much more skill than running.



## Item 3/10: Creativity

Participant picks: running, golf

To what extent are there opportunities to express creativity during the activity?

- ▶ golf admits much more creativity than running.
- ▶ golf admits somewhat more creativity than running.
- ▶ Both admit roughly the same amount of creativity.
- ▶ running admits somewhat more creativity than golf.
- ▶ running admits much more creativity than golf.



# Factor model



# Manuscript under review

jpritikin / pcFactorStan

Code Issues 0 Pull requests 0 Actions Projects 0 Wiki Security Insights Settings

Stan Models for the Paired Comparison Factor Model Edit

stan bayesian-inference factor-analysis paired-comparisons r-package Manage topics

104 commits 1 branch 0 packages 0 releases 1 contributor

Branch: master New pull request Create new file Upload files Find file Clone or download

jpritikin Doc psiScalePrior Latest commit 3e7432d on Oct 7

|                        |                                                                   |              |
|------------------------|-------------------------------------------------------------------|--------------|
| R                      | Doc psiScalePrior                                                 | last month   |
| data                   | Sort factor levels alphabetically (as in genCloud script)         | 6 months ago |
| inst/itemModelExplorer | Reparameterize                                                    | 4 months ago |
| man-roxygen            | Bump version                                                      | 2 months ago |
| src                    | Remove extra prior on factor score; add factor correlation matrix | last month   |
| tests                  | Remove extra prior on factor score; add factor correlation matrix | last month   |
| tools                  | Remove extra prior on factor score; add factor correlation matrix | last month   |
| vignettes              | Remove extra prior on factor score; add factor correlation matrix | last month   |
| .Rbuildignore          | Doc tweaks                                                        | 2 months ago |
| .gitignore             | Doc tweaks                                                        | 2 months ago |
| .travis.yml            | Try to cov faster                                                 | 2 months ago |




# Demographics, sex & country


|                | source            |                  |
|----------------|-------------------|------------------|
|                | public<br>n = 230 | MTurk<br>n = 757 |
| sex            |                   |                  |
| female         | 150 (65%)         | 299 (39%)        |
| male           | 75 (33%)          | 456 (60%)        |
| <i>missing</i> | 5 (2%)            | 2 (0%)           |
| country        |                   |                  |
| Austria        | 4 (2%)            | 0 (0%)           |
| Canada         | 6 (3%)            | 9 (1%)           |
| Germany        | 16 (7%)           | 0 (0%)           |
| India          | 1 (0%)            | 79 (10%)         |
| United Kingdom | 30 (13%)          | 0 (0%)           |
| USA            | 147 (64%)         | 641 (85%)        |
| other          | 16 (7%)           | 13 (2%)          |
| <i>missing</i> | 10 (4%)           | 15 (2%)          |




# Incomplete design



## Sample size distribution




# Posterior item loadings



## Posterior flow propensity scores

For activities with a sample size  $\geq 11$



## Possible student projects

## Some ideas:

- ▶ Confirmatory factor analysis
  - ▶ Structural equation modeling
  - ▶ Get more data on high flow propensity areas (e.g., martial arts)
  - ▶ Investigate measurement invariance in a new sample




## Questions?



## Agenda

- ▶ OpenMx
- ▶ Heritability of cortical thickness
- ▶ Confidential analyses
- ▶ Psychological wellbeing
- ▶ Flow & physical activities
- ▶ Understanding human wellbeing (*if time permits*)




# 2017 fNIRS Workshop at Kingston, RI




#### Possible future research direction

# Working principles of NIRS



Absorption spectra of Hb



## NIRS compared to MRI

Both measure the blood oxygen level-dependent (BOLD) response<sup>26</sup>

| method | resolution |                     | depth<br>pervasion | mobility | cost         |
|--------|------------|---------------------|--------------------|----------|--------------|
|        | temporal   | spatial             |                    |          |              |
| MRI    | 1-2 s      | 64 mm <sup>3</sup>  | good               | poor     | >\$1 million |
| NIRS   | 100-400 ms | 100 mm <sup>3</sup> | 2-3 cm             | good     | ~\$20-100k   |

(Table from Min, Marzelli, & Yoo, 2010)



<sup>26</sup>Cui, Bray, Bryant, Glover, and Reiss (2011)

<sup>27</sup><https://www.openwater.cc>

## NIRS compared to MRI

Both measure the blood oxygen level-dependent (BOLD) response<sup>26</sup>

| method | resolution |                     | depth<br>pervasion | mobility | cost         |
|--------|------------|---------------------|--------------------|----------|--------------|
|        | temporal   | spatial             |                    |          |              |
| MRI    | 1-2 s      | 64 mm <sup>3</sup>  | good               | poor     | >\$1 million |
| NIRS   | 100-400 ms | 100 mm <sup>3</sup> | 2-3 cm             | good     | ~\$20-100k   |

(Table from Min et al., 2010)


New tech on the horizon ( $2 \mu\text{m}^3$  spatial resolution)<sup>27</sup>




<sup>26</sup>Cui et al. (2011)

<sup>27</sup> <https://www.openwater.cc>

## Triangulate



# Flow inductions



Tetris, pong<sup>28</sup>

---

<sup>28</sup>Yoshida et al. (2014); de Sampaio Barros, Araújo-Moreira, Trevelin, and Radel (2018)

# Wim Hof method

Not mere relaxation

Clear physiological correlates<sup>29</sup>

Simple method:

- ▶ Cold exposure
- ▶ Breathing
- ▶ Commitment



---

<sup>29</sup>Kox et al. (2014); Muzik, Reilly, and Diwadkar (2018)




# Data analysis challenge

Multivariate, multilevel time series data

Possible approaches:

- ▶ Windowed cross correlation
- ▶ Autoregressive models
- ▶ State space models
- ▶ Dynamic connectivity<sup>30</sup>
- ▶ Novel methods<sup>31</sup>



<sup>30</sup>Santosa, Aarabi, Perlman, and Huppert (2017)

<sup>31</sup>Pritikin, Hunter, et al. (2017); Pritikin (2017b)



## Next steps

# Speculative

With Co-PI Tim Brick,  
assistant professor at Penn State Univ

Apply for a grant from

- ▶ John Templeton Foundation
  - ▶ The Mind and Life Institute
  - ▶ Nat'l Institute of Mental Health?




# Questions?



# Summary

Think of

- ▶ Applied Statistician
- ▶ Data Scientist
- ▶ Software Engineer
- ▶ Your friendly collaborator
- ▶ Psychological Flourishing



## Contact

jpritikin@pobox.com

<http://exuberant-island.surge.sh/>



Birnbaum, A. (1968). Some latent trait models and their use in inferring an examinee's ability. In F. M. Lord & M. R. Novick (Eds.), *Statistical theories of mental test scores* (pp. 397–479). Reading, MA: Addison-Wesley.

Boker, S. M., Brick, T. R., Pritikin, J. N., Wang, Y., von Oertzen, T., Brown, D., ... Neale, M. C. (2015). Maintained individual data distributed likelihood estimation. *Multivariate Behavioral Research*, 50(6), 706–720. doi: 10.1080/00273171.2015.1094387

Bradley, R. A., & Terry, M. E. (1952). Rank analysis of incomplete block designs: I. The method of paired comparisons. *Biometrika*, 39(3/4), 324–345. doi: 10.2307/2334029

Butler, A. C., Chapman, J. E., Forman, E. M., & Beck, A. T. (2006). The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*, 26(1), 17–31.

Csikszentmihalyi, M., Latter, P., & Duranso, C. W. (2017). *Running Flow*. Champaign, IL: Human Kinetics.

Cui, X., Bray, S., Bryant, D. M., Glover, G. H., & Reiss, A. L. (2011). A quantitative comparison of nirs and fmri across multiple cognitive tasks. *Neuroimage*, 54(4), 2808–2821.


- de Manzano, Ö., Theorell, T., Harmat, L., & Ullén, F. (2010). The psychophysiology of flow during piano playing. *Emotion, 10*(3), 301. doi: 10.1037/a0018432

de Sampaio Barros, M. F., Araújo-Moreira, F. M., Trevelin, L. C., & Radel, R. (2018). Flow experience and the mobilization of attentional resources. *Cognitive, Affective, & Behavioral Neuroscience*. doi: 10.3758/s13415-018-0606-4

Dietrich, A. (2003). Functional neuroanatomy of altered states of consciousness: The transient hypofrontality hypothesis. *Consciousness and Cognition, 12*(2), 231–256.

Goyal, M., Singh, S., Sibinga, E. M., Gould, N. F., Rowland-Seymour, A., Sharma, R., ... others (2014). Meditation programs for psychological stress and well-being: A systematic review and meta-analysis. *JAMA Internal Medicine, 174*(3), 357–368.

Jackson, S. A., Ford, S. K., Kimiecik, J. C., & Marsh, H. W. (1998). Psychological correlates of flow in sport. *Journal of Sport and Exercise Psychology, 20*(4), 358–378. doi: 10.1123/jsep.20.4.358

Keller, J., & Landhäußer, A. (2012). The flow model revisited. In S. Engeser (Ed.), *Advances in flow research* (pp. 51–64). 

Springer. doi: 10.1007/978-1-4614-2359-1

- Kotler, S. (2014). *The rise of superman: Decoding the science of ultimate human performance*. Houghton Mifflin Harcourt.
- Kox, M., van Eijk, L. T., Zwaag, J., van den Wildenberg, J., Sweep, F. C., van der Hoeven, J. G., & Pickkers, P. (2014). Voluntary activation of the sympathetic nervous system and attenuation of the innate immune response in humans. *Proceedings of the National Academy of Sciences*, 201322174.
- Leary, M. R. (2007). *The curse of the self: Self-awareness, egotism, and the quality of human life*. Oxford University Press.
- Limb, C. J., & Braun, A. R. (2008). Neural substrates of spontaneous musical performance: An fMRI study of jazz improvisation. *PLoS ONE*, 3(2), 1–9. doi: 10.1371/journal.pone.0001679
- Min, B.-K., Marzelli, M. J., & Yoo, S.-S. (2010). Neuroimaging-based approaches in the brain–computer interface. *Trends in Biotechnology*, 28(11), 552–560.
- Muzik, O., Reilly, K. T., & Diwadkar, V. A. (2018). “brain over body”—a study on the willful regulation of autonomic function during cold exposure. *NeuroImage*, 172, 632 - 641. doi: 10.1016/j.neuroimage.2018.01.030



10.1016/j.neuroimage.2018.01.067

- Neale, M. C., Hunter, M. D., Pritikin, J. N., Zahery, M., Brick, T. R., Kirkpatrick, R., ... Boker, S. M. (2016). OpenMx 2.0: Extended structural equation and statistical modeling. *Psychometrika*, 81(2), 535–549. doi: 10.1007/s11336-014-9435-8

Pollan, M. (2018). *How to change your mind: The new science of psychedelics*. London, England: Penguin Books Limited.

Pritikin, J. N. (2016). A computational note on the application of the Supplemented EM algorithm to item response models. *arXiv preprint arXiv:1605.00860*.

Pritikin, J. N. (2017a). A comparison of parameter covariance estimation methods for item response models in an expectation-maximization framework. *Cogent Psychology*, 4(1), 1279435. doi: 10.1080/23311908.2017.1279435

Pritikin, J. N. (2017b, May). *Toward multilevel variance decomposition of interactions in non-linear structural equation models*. (Talk presented at Modern Modeling Methods)

Pritikin, J. N., Brick, T. R., & Neale, M. C. (2018). Multivariate



normal maximum likelihood with both ordinal and continuous variables, and data missing at random. *Behavior Research Methods*, 50(2), 395–401. doi: 10.3758/s13428-017-1011-6

Pritikin, J. N., Hunter, M. D., & Boker, S. M. (2015). Modular open-source software for Item Factor Analysis. *Educational and Psychological Measurement*, 75(3), 458-474. doi: 10.1177/0013164414554615

Pritikin, J. N., Hunter, M. D., von Oertzen, T., Brick, T. R., & Boker, S. M. (2017). Many-level multilevel structural equation modeling: An efficient evaluation strategy. *Structural Equation Modeling: A Multidisciplinary Journal*, 24(5), 684-698. doi: 10.1080/10705511.2017.1293542

Pritikin, J. N., Rappaport, L. M., & Neale, M. C. (2017). Likelihood-based confidence intervals for a parameter with an upper or lower bound. *Structural Equation Modeling: A Multidisciplinary Journal*, 24(3), 395-401. doi: 10.1080/10705511.2016.1275969

Pritikin, J. N., & Schmidt, K. M. (2016). Model builder for Item Factor Analysis with OpenMx. *R Journal*, 8(1), 182-203.



Pritikin, J. N., Schmitt, J. E., & Neale, M. C. (2019). Cloud computing for voxel-wise SEM analysis of MRI data. *Structural Equation Modeling: A Multidisciplinary Journal*, 26(3), 470-480. doi: 10.1080/10705511.2018.1521285

Santosa, H., Aarabi, A., Perlman, S. B., & Huppert, T. J. (2017). Characterization and correction of the false-discovery rates in resting state connectivity using functional near-infrared spectroscopy. *Journal of Biomedical Optics*, 22(5), 055002–055002. doi: 10.1117/1.JBO.22.5.055002

Sawyer, K. (2007). *Group genius: The creative power of collaboration*. Basic Books.

Schmitt, J. E., Neale, M. C., Clasen, L. S., Liu, S., Seidlitz, J., Pritikin, J. N., ... Raznahan, A. (2019). A comprehensive quantitative genetic analysis of cerebral surface area in youth. *Journal of Neuroscience*, 39(16), 3028–3040. doi: 10.1523/JNEUROSCI.2248-18.2019

Schmitt, J. E., Neale, M. C., Fassassi, B., Perez, J., Lenroot, R. K., Wells, E. M., & Giedd, J. N. (2014). The dynamic role of genetics on cortical patterning during childhood and adolescence. *Development and Psychopathology*, 26, 1371–1383.



adolescence. *Proceedings of the National Academy of Sciences*, 111(18), 6774–6779. Retrieved from <http://www.pnas.org/content/111/18/6774> doi: 10.1073/pnas.1311630111

Schmitt, J. E., Raznahan, A., Clasen, L. S., Wallace, G. L., Pritikin, J. N., Lee, N. R., ... Neale, M. C. (2019). The dynamic associations between cortical thickness and general intelligence are genetically-mediated. *Cerebral Cortex*, 1–10. doi: 10.1093/cercor/bhz007

Wegner, D. M. (2002). *The illusion of conscious will*. MIT press.

Wrangham, R. (2009). *Catching fire: How cooking made us human.* Basic Books.

Yoshida, K., Sawamura, D., Inagaki, Y., Ogawa, K., Ikoma, K., & Sakai, S. (2014). Brain activity during the flow experience: A functional near-infrared spectroscopy study. *Neuroscience Letters*, 573, 30–34. doi: 10.1016/j.neulet.2014.05.011

