

# A new implementation of Item Factor Analysis: Accuracy, flexibility, and speed

Joshua N. Pritikin

Department of Psychology  
University of Virginia

28 Mar 2013



# Why a new implementation?

- ▶ ConQuest, \$750
- ▶ IRTPRO, \$495
- ▶ flexMIRT,  $\approx$  \$100 per year

## Weaknesses:

- ▶ flexibility, customization
- ▶ Windows-centric
- ▶ non-zero \$ barrier to entry

Also, current open-source software is fragmented and uncompetitive.



# Goal

Meet or exceed capabilities of all commercial software

Where to start?

Until recently, marginal maximum likelihood (Bock & Aitkin, 1981) with adaptive Gauss-Hermite quadrature (Schilling & Bock, 2005) was the leading algorithm (Wirth & Edwards, 2007).

The IRTPRO manual recommended

- ▶ marginal maximum likelihood (Bock & Aitkin, 1981) with analytic dimension reduction (Cai, 2010b)
- ▶ Metropolis-Hastings Robbins-Monro (Cai, 2010a)



# A starting point

- ▶ marginal maximum likelihood (Bock & Aitkin, 1981) with analytic dimension reduction (Cai, 2010b)
- ▶ Metropolis-Hastings Robbins-Monro (Cai, 2010a)

Which? *Both*

- ▶ Radically different approaches
- ▶ Similar performance on a broad class of problems

An open source implementation of MH-RM exists (Chalmers, 2012).

Start with marginal maximum likelihood.



# Analytic dimension reduction

Necessary restriction of the factor covariance structure

$$\begin{pmatrix} \Sigma & \\ 0 & \text{diag}(\tau) \end{pmatrix}$$

where  $\Sigma$  is any covariance structure. For example,

$$\begin{pmatrix} 1 & & & \\ \rho & 1 & & \\ 0 & 0 & 1 & \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

All factors have unit variance. In a multigroup model, the primary factors can be correlated. The specific factors are mutually uncorrelated and uncorrelated with the primary factors (Cai, 2010b).



# Analytic dimension reduction




FIGURE 1.

A two-tier model for testlet-based assessments. The items measure 2 correlated primary dimensions and the test is based up of 4 testlets, creating 4 additional specific dimensions.

(Cai, 2010b, p. 583)



# Analytic dimension reduction




FIGURE 2.

A two-tier model for longitudinal item response data. The 9 items are administered to the same group of individuals at two time points, creating two time-specific primary dimensions and 9 item-specific doublets capturing residual correlations.

(Cai, 2010b, p. 584)



# Analytic dimension reduction




FIGURE 3.

Random intercept item bifactor model as a two-tier model. The inclusion of a random intercept that is orthogonal to the other factors is a distinguishing feature of this model. The general factor, on which all items load, and the random intercept are the primary dimensions.

(Cai, 2010b, p. 585)





# Status

- ▶ R,  $\approx$  1300 lines
- ▶ C,  $\approx$  2350 lines

Implements marginal maximum likelihood (Bock & Aitkin, 1981) with analytic dimension reduction (Cai, 2010b)

OpenMx plugin, not yet merged

EAP person scores (Bock & Mislevy, 1982)

Item models: M3PL, GPCM, or invent your own

$S - \chi^2$  with polytomous extension (Orlando & Thissen, 2000)

Infit & outfit (Wright & Masters, 1982)

Plots: item characteristic curve (1d), data vs model, and information



$$S - \chi^2$$

(Explain on blackboard)



# High-priority wish list

Merge into OpenMx (Boker et al., 2011)

Nominal model w/ analytic Newton-Raphson (Baker & Kim, 2004)  
(open source math exist; Chalmers, 2012)

Item parameter standard errors (Cai, 2008)

Multi-group (Cai, Yang, & Hansen, 2011)

Structural latent trait model

Hierarchical factor model



# Item parameter standard errors

What do they mean?

$$\mathcal{I}(\hat{\theta}|Y_o) = \mathcal{I}_c(\hat{\theta}) - \mathcal{I}_m(\hat{\theta})$$

where the information matrix  $\mathcal{I}$  at the item parameter vector estimate  $\hat{\theta}$  conditional on the observed data  $Y_o$  is the complete information matrix  $\mathcal{I}_c$  minus the information matrix of the missing data  $\mathcal{I}_m$  (Cai, 2008).



# Accuracy, performance

Let's take a look ...



# Standardize (rescale) the latent distribution or not?

- ▶ 20 2PL items
- ▶ 500 persons
- ▶ 17 point GH quadrature
- ▶ 0% missing
- ▶ 500 Monte Carlo replications  $M$
- ▶ Rescale or not (Liu, Rubin, & Wu, 1998)

Examine  $-2LL$ ,  $S-\chi^2$ , and bias.

$$\text{bias} = \hat{\theta} - \theta_{true} \text{ where } \hat{\theta} = \frac{1}{M} \sum_{m=1}^M \theta_m$$



# -2LL




$$S - \chi^2$$





# Bias



Bias ranged from -0.0695 to 0.1188 with 50% of the bias between -0.0033 to 0.0174 with the median at 0.0056. For comparison, Winstep obtained bias ranging from 0.01 to 0.13 (Wang & Chen, 2005).



# Which model? How many quadrature points?

- ▶ 20 2PL items
- ▶ 500 persons
- ▶ 0% missing
- ▶ GH quadrature
- ▶ This and subsequent studies are rescaled (Liu et al., 1998)

Traditional parameterization


$$\frac{1}{1 + \exp(-a(\theta - b))}$$

or slope-intercept form

$$\frac{1}{1 + \exp(-(a\theta + c))} \text{ where } b = \frac{c}{-a}$$



# Likelihood by item model and quadrature points




# Bias by % missing

- ▶ 20 2PL items
- ▶ 500 persons
- ▶ 17 point GH quadrature
- ▶ 500 Monte Carlo replications


For the missing at random condition, data was replaced by NA depending on the first 5 items.




# Bias by % missing




# Bias by % missing, MCAR 0%



# Bias by % missing, MCAR 2%




# Bias by % missing, MCAR 4%







# Bias by % missing, MCAR 6%




# Bias by % missing, MCAR 8%




# Bias by % missing, MCAR 10%




# Bias by % missing, MAR 0%




# Bias by % missing, MAR 2%




# Bias by % missing, MAR 4%



# Bias by % missing, MAR 6%




# Bias by % missing, MAR 8%





# Bias by % missing, MAR 10%



# Cai (2010b) parameter recovery simulation


- ▶ 20 M2PL items
- ▶ 2 primary dimensions
- ▶ 4 specific dimensions formed by 4 pairs of item doublets
- ▶  $N = 500$  per replication
- ▶ 13 point GH quadrature<sup>1</sup>
- ▶ 500 Monte Carlo replications

---

<sup>1</sup>IRTPRO uses a 21 point equally spaced quadrature by default.



# Cai (2010b) parameter recovery simulation



Note: slight pos bias; comparable to Cai (2010b) except for green; *very* slow




# Acknowledgment

- ▶ Karen
- ▶ OpenMx development team
- ▶ Steve
- ▶ Timo
- ▶ UVa grad students

& colleagues who I forgot to mention

## Questions?



- Baker, F. B., & Kim, S. H. (2004). *Item response theory: Parameter estimation techniques* (2nd ed.). CRC Press.
- Bock, R. D., & Aitkin, M. (1981). Marginal maximum likelihood estimation of item parameters: Application of an EM algorithm. *Psychometrika*, 46, 443-459.
- Bock, R. D., & Mislevy, R. J. (1982). Adaptive EAP estimation of ability in a microcomputer environment. *Applied Psychological Measurement*, 6(4), 431-444.
- Boker, S., Neale, M., Maes, H., Wilde, M., Spiegel, M., Brick, T., ... others (2011). OpenMx: An open source extended structural equation modeling framework. *Psychometrika*, 76(2), 306-317.
- Cai, L. (2008). SEM of another flavour: Two new applications of the supplemented EM algorithm. *British Journal of Mathematical and Statistical Psychology*, 61, 309-329.
- Cai, L. (2010a). High-dimensional exploratory item factor analysis by a Metropolis-Hastings Robbins-Monro algorithm. *Psychometrika*, 75, 33-57.
- Cai, L. (2010b). A two-tier full-information item factor analysis model with applications. *Psychometrika*, 75, 581-612. Retrieved from <http://dx.doi.org/10.1007/s11336-010-9178-0> doi: 



10.1007/s11336-010-9178-0

- Cai, L., Yang, J. S., & Hansen, M. (2011). Generalized Full-Information Item Bifactor Analysis. *Psychological Methods*, 16(3), 221–248.
- Chalmers, R. P. (2012). mirt: A multidimensional item response theory package for the R environment. *Journal of Statistical Software*, 48(6), 1–29. Retrieved from <http://www.jstatsoft.org/v48/i06/>
- Liu, C., Rubin, D. B., & Wu, Y. N. (1998). Parameter expansion to accelerate EM: the PX-EM algorithm. *Biometrika*, 85(4), 755–770.
- Orlando, M., & Thissen, D. (2000). Likelihood-Based Item-Fit Indices for Dichotomous Item Response Theory Models. *Applied Psychological Measurement*, 24(1), 50–64.
- Schilling, S., & Bock, R. D. (2005). High-dimensional maximum marginal likelihood item factor analysis by adaptive quadrature. *Psychometrika*, 70(3), 533–555.
- Wang, W.-C., & Chen, C.-T. (2005). Item parameter recovery, standard error estimates, and fit statistics of the Winsteps program for the family of Rasch models. *Educational and Psychological Measurement*, 25(1), 1–20.



*Psychological Measurement*, 65(3), 376–404.

Wirth, R. J., & Edwards, M. C. (2007). Item factor analysis: Current approaches and future directions. *Psychological Methods*, 12(1), 58.

Wright, B. D., & Masters, G. N. (1982). *Rating scale analysis*. Chicago: MESA Press.

