

# Cooperative Game Theory: Solution concepts

Fall 2020

# References

- G. Chalkiadakis, E. Elkind, M. Wooldridge  
Computational Aspects of Cooperative Game Theory  
Morgan & Claypool, 2012 Wikipedia.
- G. Owen F  
Game Theory  
3rd edition, Academic Press, 1995

- 1 Definitions
- 2 Stability notions

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where
  - agents can benefit by cooperating, and
  - binding agreements are possible.

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where
  - agents can benefit by cooperating, and
  - binding agreements are possible.
- In cooperative games, actions are taken by groups of agents, **coalitions**, and payoffs are given to

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where
  - agents can benefit by cooperating, and
  - binding agreements are possible.
- In cooperative games, actions are taken by groups of agents, **coalitions**, and payoffs are given to
  - the group. Those have to be divided among its members:  
**Transferable utility games (TU)**.



# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where
  - agents can benefit by cooperating, and
  - binding agreements are possible.
- In cooperative games, actions are taken by groups of agents, **coalitions**, and payoffs are given to
  - the group. Those have to be divided among its members: **Transferable utility games (TU)**.
  - individuals. **Non-transferable utility games (NTU)**.

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where
  - agents can benefit by cooperating, and
  - binding agreements are possible.
- In cooperative games, actions are taken by groups of agents, **coalitions**, and payoffs are given to
  - the group. Those have to be divided among its members: **Transferable utility games (TU)**.
  - individuals. **Non-transferable utility games (NTU)**.
- For the moment we focus on **TU games**

# Non-Cooperative versus cooperative Games

- Non-cooperative game theory model scenarios where players cannot make binding agreements.
- Cooperative game theory model scenarios, where
  - agents can benefit by cooperating, and
  - binding agreements are possible.
- In cooperative games, actions are taken by groups of agents, **coalitions**, and payoffs are given to
  - the group. Those have to be divided among its members: **Transferable utility games (TU)**.
  - individuals. **Non-transferable utility games (NTU)**.
- For the moment we focus on **TU games**
- **Notation:**  $N$ , set of players,  $C, S, X \subseteq N$  are coalitions.

# Notation

# Notation

- For a set  $A$ :

# Notation

- For a set  $A$ :
  - $\mathcal{C}_A$  denotes the subsets of  $A$ , i.e.,  $C \subseteq A$ .
  - $\mathcal{P}_A$  denotes the partitions of  $A$ .

# Notation

- For a set  $A$ :
  - $\mathcal{C}_A$  denotes the subsets of  $A$ , i.e.,  $C \subseteq A$ .
  - $\mathcal{P}_A$  denotes the partitions of  $A$ .
- For a set of players  $N$ , a **coalition** is any subset of  $N$ .

# Notation

- For a set  $A$ :
  - $\mathcal{C}_A$  denotes the subsets of  $A$ , i.e.,  $C \subseteq A$ .
  - $\mathcal{P}_A$  denotes the partitions of  $A$ .
- For a set of players  $N$ , a **coalition** is any subset of  $N$ .  
 $N$  is the **grand coalition**.



# Notation

- For a set  $A$ :
  - $\mathcal{C}_A$  denotes the subsets of  $A$ , i.e.,  $C \subseteq A$ .
  - $\mathcal{P}_A$  denotes the partitions of  $A$ .
- For a set of players  $N$ , a **coalition** is any subset of  $N$ .  
 $N$  is the **grand coalition**.
- A **partition** of  $N$  is a splitting of all the players into disjoint coalitions.

# Characteristic Function Games

- A **characteristic function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \rightarrow \mathbb{R}$  is the **characteristic function**.

# Characteristic Function Games

- A **characteristic function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \rightarrow \mathbb{R}$  is the **characteristic function**.
  - for each coalition of players  $C \subseteq N$ ,  $v(C)$  is the amount that the members of  $C$  can earn by working together

# Characteristic Function Games

- A **characteristic function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \rightarrow \mathbb{R}$  is the **characteristic function**.
  - for each coalition of players  $C \subseteq N$ ,  $v(C)$  is the amount that the members of  $C$  can earn by working together
- usually it is assumed that  $v$  is

# Characteristic Function Games

- A **characteristic function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \rightarrow \mathbb{R}$  is the **characteristic function**.
  - for each coalition of players  $C \subseteq N$ ,  $v(C)$  is the amount that the members of  $C$  can earn by working together
- usually it is assumed that  $v$  is
  - normalized:  $v(\emptyset) = 0$ ,
  - non-negative:  $v(C) \geq 0$ , for any  $C \subseteq N$ , and
  - monotone:  $v(C) \leq v(D)$ , for any  $C, D$  such that  $C \subseteq D$
- Example:  $N = \{A, B, C\}$  and

$\mathcal{C}_N$	$\emptyset$	A	B	C	AB	AC	BC	ABC
$v$	0	12	0	0	18	18	18	24

# Partition Function Games

- A **partition function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \times \mathcal{P}_{N \setminus S} \rightarrow \mathbb{R}$  is the **partition function**.

# Partition Function Games

- A **partition function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \times \mathcal{P}_{N \setminus S} \rightarrow \mathbb{R}$  is the **partition function**.
  - for each subset of players  $C \subseteq N$  and partition  $P$ ,  $v(C; P)$  is the amount that the members of  $C$  can earn by working together assuming  $N$  is splitted according to  $P$ .

# Partition Function Games

- A **partition function** game is a pair  $(N, v)$ , where:
  - $N = \{1, \dots, n\}$  is the **set of players** and
  - $v : \mathcal{C}_N \times \mathcal{P}_{N \setminus S} \rightarrow \mathbb{R}$  is the **partition function**.
  - for each subset of players  $C \subseteq N$  and partition  $P$ ,  $v(C; P)$  is the amount that the members of  $C$  can earn by working together assuming  $N$  is splitted according to  $P$ .
- Example:  $N = \{A, B, C\}$  and

$\mathcal{P}_N$	$\emptyset, ABC$		$AB, C$		$AC, B$		$BC, A$		$A, B, C$		
$C$	$\emptyset$	ABC	AB	C	AC	B	BC	A	A	B	C
$v$	0	24	18	0	18	0	18	0	12	6	0



# Characteristic Function Games vs. Partition Function Games

# Characteristic Function Games vs. Partition Function Games

- In partition function games (PFG) the payoff obtained by a coalition depends on the **other coalitions**

# Characteristic Function Games vs. Partition Function Games

- In partition function games (PFG) the payoff obtained by a coalition depends on the **other coalitions**
- In characteristic function games (CFG) the payoff of each coalition only depends on **the action of that coalition**  
in such games, each coalition can be identified with the profit it obtains by choosing its best action

# Characteristic Function Games vs. Partition Function Games

- In partition function games (PFG) the payoff obtained by a coalition depends on the **other coalitions**
- In characteristic function games (CFG) the payoff of each coalition only depends on **the action of that coalition** in such games, each coalition can be identified with the profit it obtains by choosing its best action
- We restrict in this course to focus on **characteristic function games**, and use the term **coalition game** to refer to such a game.

# Buying Ice-Cream Game

# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.

# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.
- There are three types of ice-cream tubs for sale:

# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.
- There are three types of ice-cream tubs for sale:





# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.
- There are three types of ice-cream tubs for sale:

Type 1 costs \$7, contains 500g

Type 2 costs \$9, contains 750g

Type 3 costs \$11, contains 1kg



# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.
- There are three types of ice-cream tubs for sale:
  - Type 1 costs \$7, contains 500g
  - Type 2 costs \$9, contains 750g
  - Type 3 costs \$11, contains 1kg
- The children have utility for ice-cream but do not care about money.



# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.

- There are three types of ice-cream tubs for sale:

Type 1 costs \$7, contains 500g

Type 2 costs \$9, contains 750g

Type 3 costs \$11, contains 1kg



- The children have utility for ice-cream but do not care about money.
- The payoff of each group is the maximum quantity of ice-cream the members of the group can buy by pooling all their money.

# Buying Ice-Cream Game

- We have a group of  $n$  children, each has some amount of money the  $i$ -th child has  $b_i$  dollars.
- There are three types of ice-cream tubs for sale:
  - Type 1 costs \$7, contains 500g
  - Type 2 costs \$9, contains 750g
  - Type 3 costs \$11, contains 1kg
- The children have utility for ice-cream but do not care about money.
- The payoff of each group is the maximum quantity of ice-cream the members of the group can buy by pooling all their money.
- The ice-cream can be shared arbitrarily within the group.



# Ice-Cream Game: Characteristic Function

# Ice-Cream Game: Characteristic Function



Charlie: \$6



Marcie: \$4



Pattie: \$3

# Ice-Cream Game: Characteristic Function



Charlie: \$6



Marcie: \$4



Pattie: \$3



$w = 500$

$p = \$7$



$w = 750$

$p = \$9$



$w = 100$

$p = \$11$

# Ice-Cream Game: Characteristic Function



Charlie: \$6



Marcie: \$4



Pattie: \$3



$w = 500$

$p = \$7$



$w = 750$

$p = \$9$



$w = 100$

$p = \$11$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 750, v(\{C, P\}) = 750, v(\{M, P\}) = 500$
- $v(\{C, M, P\}) = 1000$



# Outcomes

An **outcome** of a game  $\Gamma = (N, v)$  is a pair  $(P, x)$ , where:

# Outcomes

An **outcome** of a game  $\Gamma = (N, v)$  is a pair  $(P, x)$ , where:

- $P = (C_1, \dots, C_k) \in \mathcal{P}_N$  is a **coalition structure**

# Outcomes

An **outcome** of a game  $\Gamma = (N, v)$  is a pair  $(P, x)$ , where:

- $P = (C_1, \dots, C_k) \in \mathcal{P}_N$  is a **coalition structure**
- $x = (x_1, \dots, x_n)$  is a **payoff vector**, which distributes the value of each coalition in  $P$ :
  - $x_i \geq 0$ , for all  $i \in N$
  - $\sum_{i \in C} x_i = v(C)$ , for each  $C \in P$ ,

# Outcomes

An **outcome** of a game  $\Gamma = (N, v)$  is a pair  $(P, x)$ , where:

- $P = (C_1, \dots, C_k) \in \mathcal{P}_N$  is a **coalition structure**
- $x = (x_1, \dots, x_n)$  is a **payoff vector**, which distributes the value of each coalition in  $P$ :
  - $x_i \geq 0$ , for all  $i \in N$
  - $\sum_{i \in C} x_i = v(C)$ , for each  $C \in P$ , **feasibility**

# Outcome:example

Suppose  $v(\{1, 2, 3\}) = 9$  and  $v(\{4, 5\}) = 4$

# Outcome:example

Suppose  $v(\{1, 2, 3\}) = 9$  and  $v(\{4, 5\}) = 4$

- $((\{1, 2, 3\}, \{4, 5\}), (3, 3, 3, 3, 1))$  is an outcome

# Outcome:example

Suppose  $v(\{1, 2, 3\}) = 9$  and  $v(\{4, 5\}) = 4$

- $((\{1, 2, 3\}, \{4, 5\}), (3, 3, 3, 3, 1))$  is an outcome
- $((\{1, 2, 3\}, \{4, 5\}), (2, 3, 2, 3, 3))$  is **NOT** an outcome as transfers between coalitions are not allowed

# Imputations



# Imputations

An outcome  $(P, x)$  is called an **imputation** if it satisfies **individual rationality**:

$$x_i \geq v(\{i\}),$$

for all  $i \in N$ .

# Imputations

An outcome  $(P, x)$  is called an **imputation** if it satisfies **individual rationality**:

$$x_i \geq v(\{i\}),$$

for all  $i \in N$ .

Notation: we denote  $\sum_{i \in A} x_i$  by  $x(A)$

## 1 Definitions

## 2 Stability notions

# What Is a Good Outcome?

# What Is a Good Outcome?

- The solutions of a game should provide **good outcomes**.

# What Is a Good Outcome?

- The solutions of a game should provide **good outcomes**.
- Let us present some stability notions related to outcomes or imputations.

# What Is a Good Outcome?

- The solutions of a game should provide **good outcomes**.
- Let us present some stability notions related to outcomes or imputations.
- To simplify the presentation we consider **superadditive** games.

# Superadditive Games

- A game  $G = (N, v)$  is called **superadditive** if

$$v(C \cup D) \geq v(C) + v(D),$$

for any two disjoint coalitions  $C$  and  $D$



# Superadditive Games

- A game  $G = (N, v)$  is called **superadditive** if

$$v(C \cup D) \geq v(C) + v(D),$$

for any two disjoint coalitions  $C$  and  $D$

- Example:  $v(C) = |C|^2$

$$v(C \cup D) = (|C| + |D|)^2 \geq |C|^2 + |D|^2 = v(C) + v(D)$$

# Superadditive Games

# Superadditive Games

- In superadditive games, two coalitions can always merge without losing money; hence, we can assume in a stable outcome  $P = (N, \emptyset)$ .

# Superadditive Games

- In superadditive games, two coalitions can always merge without losing money; hence, we can assume in a stable outcome  $P = (N, \emptyset)$ .  
Players must form the **grand coalition**

# Superadditive Games

- In superadditive games, two coalitions can always merge without losing money; hence, we can assume in a stable outcome  $P = (N, \emptyset)$ .  
Players must form the **grand coalition**
- In superadditive games, we identify outcomes with **payoff vectors** for the grand coalition

# Superadditive Games

- In superadditive games, two coalitions can always merge without losing money; hence, we can assume in a stable outcome  $P = (N, \emptyset)$ .

Players must form the **grand coalition**

- In superadditive games, we identify outcomes with **payoff vectors** for the grand coalition

i.e., an outcome is a vector  $x = (x_1, \dots, x_n)$  with  $x(N) = v(N)$

# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$



# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$

This is a superadditive game, so outcomes are payoff vectors!

# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$

This is a superadditive game, so outcomes are payoff vectors!

How should the players share the ice-cream?

# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$

This is a superadditive game, so outcomes are payoff vectors!

How should the players share the ice-cream?

$(200, 200, 350)$ ?

# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$

This is a superadditive game, so outcomes are payoff vectors!

How should the players share the ice-cream?

$(200, 200, 350)$ ?

Charlie and Marcie can get more ice-cream by buying a 500g tub on their own, and splitting it equally

# What Is a Good Outcome?



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$

This is a superadditive game, so outcomes are payoff vectors!

How should the players share the ice-cream?

$(200, 200, 350)$ ?

Charlie and Marcie can get more ice-cream by buying a 500g tub on their own, and splitting it equally

$(200, 200, 350)$  is not stable!

# The core

# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

$$\text{core}(\Gamma) = \{(P, x) \mid x(C) \geq v(C) \text{ for any } C \subseteq N\}$$



# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

$$\text{core}(\Gamma) = \{(P, x) \mid x(C) \geq v(C) \text{ for any } C \subseteq N\}$$

each coalition earns, **according to  $x$** , at least as much as it can make on its own.

# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

$$\text{core}(\Gamma) = \{(P, x) \mid x(C) \geq v(C) \text{ for any } C \subseteq N\}$$

each coalition earns, **according to  $x$** , at least as much as it can make on its own.

- Example:  $v(\{1, 2, 3\}) = 9$ ,  $v(\{4, 5\}) = 4$ ,  $v(\{2, 4\}) = 7$

# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

$$\text{core}(\Gamma) = \{(P, x) | x(C) \geq v(C) \text{ for any } C \subseteq N\}$$

each coalition earns, **according to  $x$** , at least as much as it can make on its own.

- Example:  $v(\{1, 2, 3\}) = 9$ ,  $v(\{4, 5\}) = 4$ ,  $v(\{2, 4\}) = 7$   
 $((\{1, 2, 3\}, \{4, 5\}), (3, 3, 3, 3, 1))$  is **NOT** in the core

# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

$$\text{core}(\Gamma) = \{(P, x) \mid x(C) \geq v(C) \text{ for any } C \subseteq N\}$$

each coalition earns, **according to  $x$** , at least as much as it can make on its own.

- Example:  $v(\{1, 2, 3\}) = 9$ ,  $v(\{4, 5\}) = 4$ ,  $v(\{2, 4\}) = 7$   
 $((\{1, 2, 3\}, \{4, 5\}), (3, 3, 3, 3, 1))$  is **NOT** in the core  
as  $x(\{2, 4\}) = 6$  and  $v(\{2, 4\}) = 7$

# The core

The **core** of a game  $\Gamma$  is the set of all **stable outcomes**, i.e., outcomes that no coalition wants to deviate from

$$\text{core}(\Gamma) = \{(P, x) \mid x(C) \geq v(C) \text{ for any } C \subseteq N\}$$

each coalition earns, **according to  $x$** , at least as much as it can make on its own.

- Example:  $v(\{1, 2, 3\}) = 9$ ,  $v(\{4, 5\}) = 4$ ,  $v(\{2, 4\}) = 7$   
 $((\{1, 2, 3\}, \{4, 5\}), (3, 3, 3, 3, 1))$  is **NOT** in the core  
as  $x(\{2, 4\}) = 6$  and  $v(\{2, 4\}) = 7$
- no subgroup of players can deviate so that each member of the subgroup gets more.

# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$

# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$
- $(200, 200, 350)$

# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$
- $(200, 200, 350)$  is not in the core:  $v(\{C, M\}) > x(\{C, M\})$



# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$
- $(200, 200, 350)$  is not in the core:  $v(\{C, M\}) > x(\{C, M\})$
- $(250, 250, 250)$

# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$
- $(200, 200, 350)$  is not in the core:  $v(\{C, M\}) > x(\{C, M\})$
- $(250, 250, 250)$  is in the core: alone or in pairs do not get more.
- $(750, 0, 0)$

# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$
- $(200, 200, 350)$  is not in the core:  $v(\{C, M\}) > x(\{C, M\})$
- $(250, 250, 250)$  is in the core: alone or in pairs do not get more.
- $(750, 0, 0)$  is also in the core:

# Ice-cream game: Core



Charlie: \$4



Marcie: \$3



Pattie: \$3

Ice-cream pots:  $w = (500, 750, 100)$  and  $p = (\$7, \$9, \$11)$

- $v(\emptyset) = v(\{C\}) = v(\{M\}) = v(\{P\}) = 0$
- $v(\{C, M\}) = 500, v(\{C, P\}) = 500, v(\{M, P\}) = 0$
- $v(\{C, M, P\}) = 750$
- $(200, 200, 350)$  is not in the core:  $v(\{C, M\}) > x(\{C, M\})$
- $(250, 250, 250)$  is in the core: alone or in pairs do not get more.
- $(750, 0, 0)$  is also in the core:  
Marcie and Pattie cannot get more on their own!

# Games with empty core?

# Games with empty core?

- Let  $\Gamma = (N, v)$ , where  $N = \{1, 2, 3\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise.

# Games with empty core?

- Let  $\Gamma = (N, v)$ , where  $N = \{1, 2, 3\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise.
- Consider an outcome  $(P, x)$ .

# Games with empty core?

- Let  $\Gamma = (N, v)$ , where  $N = \{1, 2, 3\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise.
- Consider an outcome  $(P, x)$ .
  - We have  $x_1, x_2, x_3 \geq 0$ ,  $x_1 + x_2 + x_3 = 1$ , and  $x_i + x_j = 1$ , for  $i \neq j$
  - As,  $x_1 + x_2 + x_3 \geq 1$ , for some  $i \in \{1, 2, 3\}$ ,  $x_i \geq 1/3$ .
  - Assume that  $i = 1$ , we have  $x_2 + x_3 = 1 - x_1 \leq 1 - 1/3 \leq 1$ !



# Games with empty core?

- Let  $\Gamma = (N, v)$ , where  $N = \{1, 2, 3\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise.
- Consider an outcome  $(P, x)$ .
  - We have  $x_1, x_2, x_3 \geq 0$ ,  $x_1 + x_2 + x_3 = 1$ , and  $x_i + x_j = 1$ , for  $i \neq j$
  - As,  $x_1 + x_2 + x_3 \geq 1$ , for some  $i \in \{1, 2, 3\}$ ,  $x_i \geq 1/3$ .
  - Assume that  $i = 1$ , we have  $x_2 + x_3 = 1 - x_1 \leq 1 - 1/3 \leq 1$ !
- Thus the core of  $\Gamma$  is empty.

# Core on payoff vectors

## Core on payoff vectors

- Suppose the game is not necessarily superadditive, but the outcomes are defined as payoff vectors for the grand coalition.

## Core on payoff vectors

- Suppose the game is not necessarily superadditive, but the outcomes are defined as payoff vectors for the grand coalition.
- Then the core may be empty, even if according to the standard definition it is not.

# Core on payoff vectors

- Suppose the game is not necessarily superadditive, but the outcomes are defined as payoff vectors for the grand coalition.
- Then the core may be empty, even if according to the standard definition it is not.
- $\Gamma = (N, v)$  with  $N = \{1, 2, 3, 4\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise

# Core on payoff vectors

- Suppose the game is not necessarily superadditive, but the outcomes are defined as payoff vectors for the grand coalition.
- Then the core may be empty, even if according to the standard definition it is not.
- $\Gamma = (N, v)$  with  $N = \{1, 2, 3, 4\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise
  - not superadditive:  $v(\{1, 2\}) + v(\{3, 4\}) = 2 > v(\{1, 2, 3, 4\})$

# Core on payoff vectors

- Suppose the game is not necessarily superadditive, but the outcomes are defined as payoff vectors for the grand coalition.
- Then the core may be empty, even if according to the standard definition it is not.
- $\Gamma = (N, v)$  with  $N = \{1, 2, 3, 4\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise
  - not superadditive:  $v(\{1, 2\}) + v(\{3, 4\}) = 2 > v(\{1, 2, 3, 4\})$
  - no payoff vector for the grand coalition is in the core: either  $\{1, 2\}$  or  $\{3, 4\}$  get less than 1, so can deviate

# Core on payoff vectors

- Suppose the game is not necessarily superadditive, but the outcomes are defined as payoff vectors for the grand coalition.
- Then the core may be empty, even if according to the standard definition it is not.
- $\Gamma = (N, v)$  with  $N = \{1, 2, 3, 4\}$  and  $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise
  - not superadditive:  $v(\{1, 2\}) + v(\{3, 4\}) = 2 > v(\{1, 2, 3, 4\})$
  - no payoff vector for the grand coalition is in the core:  
either  $\{1, 2\}$  or  $\{3, 4\}$  get less than 1, so can deviate
  - But  $((\{1, 2\}, \{3, 4\}), (1/2, 1/2, 1/2, 1/2))$  is in the core



# $\epsilon$ -Core

# $\epsilon$ -Core

- When the core is empty, we may want to find approximately stable outcomes.

# $\epsilon$ -Core

- When the core is empty, we may want to find approximately stable outcomes.
- We need to relax the notion of the core:

# $\epsilon$ -Core

- When the core is empty, we may want to find approximately stable outcomes.
- We need to relax the notion of the core:  
core:  $(P, x) : x(C) \geq v(C)$ , for all  $C \subseteq N$

# $\epsilon$ -Core

- When the core is empty, we may want to find approximately stable outcomes.
- We need to relax the notion of the core:  
core:  $(P, x) : x(C) \geq v(C), \text{ for all } C \subseteq N$   
 $\epsilon$ -core:  $\{(P, x) : x(C) \geq v(C) - \epsilon, \text{ for all } C \subseteq N\}$

# $\epsilon$ -Core

- When the core is empty, we may want to find approximately stable outcomes.
- We need to relax the notion of the core:  
core:  $(P, x) : x(C) \geq v(C)$ , for all  $C \subseteq N$   
 $\epsilon$ -core:  $\{(P, x) : x(C) \geq v(C) - \epsilon, \text{ for all } C \subseteq N\}$
- $\Gamma = (N, v)$ ,  $N = \{1, 2, 3\}$  and  
 $v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise

# $\epsilon$ -Core

- When the core is empty, we may want to find approximately stable outcomes.

- We need to relax the notion of the core:

core:  $(P, x) : x(C) \geq v(C), \text{ for all } C \subseteq N$

$\epsilon$ -core:  $\{(P, x) : x(C) \geq v(C) - \epsilon, \text{ for all } C \subseteq N\}$

- $\Gamma = (N, v), N = \{1, 2, 3\}$  and

$v(C) = 1$  if  $|C| > 1$  and  $v(C) = 0$  otherwise

- $1/3$ -core is non-empty:  $(1/3, 1/3, 1/3) \in 1/3\text{-core}$

- $\epsilon$ -core is empty for any  $\epsilon < 1/3$ :

$x_i \geq 1/3$ , for some  $i = 1, 2, 3$ , so  $x(N \setminus \{i\}) \leq 2/3$ ,  
 $v(N \setminus \{i\}) = 1$

# Least Core



# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$

# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$
- We are interested in outcomes that minimize the worst-case deficit

# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$
- We are interested in outcomes that minimize the worst-case deficit
- Let  $\epsilon^*(\Gamma) = \inf\{\epsilon \mid \epsilon\text{-core of } \Gamma \text{ is not empty}\}$ .

# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$
- We are interested in outcomes that minimize the worst-case deficit
- Let  $\epsilon^*(\Gamma) = \inf\{\epsilon \mid \epsilon\text{-core of } \Gamma \text{ is not empty}\}$ .
- It can be shown that, for all  $\Gamma$ , the  $\epsilon^*(\Gamma)$ -core is not empty.

# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$
- We are interested in outcomes that minimize the worst-case deficit
- Let  $\epsilon^*(\Gamma) = \inf\{\epsilon \mid \epsilon\text{-core of } \Gamma \text{ is not empty}\}$ .
- It can be shown that, for all  $\Gamma$ , the  $\epsilon^*(\Gamma)$ -core is not empty.
- The  $\epsilon^*(\Gamma)$ -core is called the **least core** of  $\Gamma$  and  $\epsilon^*(\Gamma)$  is called the **value of the least core**

# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$
- We are interested in outcomes that minimize the worst-case deficit
- Let  $\epsilon^*(\Gamma) = \inf\{\epsilon \mid \epsilon\text{-core of } \Gamma \text{ is not empty}\}$ .
- It can be shown that, for all  $\Gamma$ , the  $\epsilon^*(\Gamma)$ -core is not empty.
- The  $\epsilon^*(\Gamma)$ -core is called the **least core** of  $\Gamma$  and  $\epsilon^*(\Gamma)$  is called the **value of the least core**
- $\Gamma = (N, v)$ ,  $N = \{1, 2, 3\}$ ,  $v(C) = (|C| > 1)$ 
  - $1/3$ -core is non-empty:  $(1/3, 1/3, 1/3) \in 1/3\text{-core}$
  - $\epsilon$ -core is empty for any  $\epsilon < 1/3$

# Least Core

- If an outcome  $(P, x)$  is in the  $\epsilon$ -core, the deficit  $v(C) - x(C)$  of any coalition is at most  $\epsilon$
- We are interested in outcomes that minimize the worst-case deficit
- Let  $\epsilon^*(\Gamma) = \inf\{\epsilon \mid \epsilon\text{-core of } \Gamma \text{ is not empty}\}$ .
- It can be shown that, for all  $\Gamma$ , the  $\epsilon^*(\Gamma)$ -core is not empty.
- The  $\epsilon^*(\Gamma)$ -core is called the **least core** of  $\Gamma$  and  $\epsilon^*(\Gamma)$  is called the **value of the least core**
- $\Gamma = (N, v)$ ,  $N = \{1, 2, 3\}$ ,  $v(C) = (|C| > 1)$ 
  - $1/3$ -core is non-empty:  $(1/3, 1/3, 1/3) \in 1/3\text{-core}$
  - $\epsilon$ -core is empty for any  $\epsilon < 1/3$
  - The least core is the  $1/3$ -core.

# Stability vs. Fairness

- Outcomes in the core may be unfair.



# Stability vs. Fairness

- Outcomes in the core may be unfair.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0$ ,  $v(\{1\}) = v(\{2\}) = 5$ ,  $v(\{1, 2\}) = 20$ 
  - $(15, 5)$  is in the core: player 2 cannot benefit by deviating.

# Stability vs. Fairness

- Outcomes in the core may be unfair.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0, v(\{1\}) = v(\{2\}) = 5, v(\{1, 2\}) = 20$ 
  - $(15, 5)$  is in the core: player 2 cannot benefit by deviating.
  - However, this is unfair since 1 and 2 are symmetric

# Stability vs. Fairness

- Outcomes in the core may be unfair.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0, v(\{1\}) = v(\{2\}) = 5, v(\{1, 2\}) = 20$ 
  - $(15, 5)$  is in the core: player 2 cannot benefit by deviating.
  - However, this is unfair since 1 and 2 are symmetric
- How do we divide payoffs in a fair way?

# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.

# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.
- **Attempt:**

# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.
- **Attempt:** given a game  $\Gamma = (N, v)$ , set

$$x_i = v(\{1, \dots, i-1, i\}) - v(\{1, \dots, i-1\}).$$

# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.
- **Attempt:** given a game  $\Gamma = (N, v)$ , set

$$x_i = v(\{1, \dots, i-1, i\}) - v(\{1, \dots, i-1\}).$$

The payoff to each player is his marginal contribution to the coalition of his predecessors

# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.
- **Attempt:** given a game  $\Gamma = (N, v)$ , set

$$x_i = v(\{1, \dots, i-1, i\}) - v(\{1, \dots, i-1\}).$$

The payoff to each player is his marginal contribution to the coalition of his predecessors

- We have  $x_1 + \dots + x_n = v(N)$  thus  $x$  is a payoff vector



# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.
- **Attempt:** given a game  $\Gamma = (N, v)$ , set

$$x_i = v(\{1, \dots, i-1, i\}) - v(\{1, \dots, i-1\}).$$

The payoff to each player is his marginal contribution to the coalition of his predecessors

- We have  $x_1 + \dots + x_n = v(N)$  thus  $x$  is a payoff vector
- However, payoff to each player depends on the order

# Marginal Contribution

- A fair payment scheme rewards each agent according to his contribution.
- **Attempt:** given a game  $\Gamma = (N, v)$ , set

$$x_i = v(\{1, \dots, i-1, i\}) - v(\{1, \dots, i-1\}).$$

The payoff to each player is his marginal contribution to the coalition of his predecessors

- We have  $x_1 + \dots + x_n = v(N)$  thus  $x$  is a payoff vector
- However, payoff to each player depends on the order
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0$ ,  $v(\{1\}) = v(\{2\}) = 5$ ,  $v(\{1, 2\}) = 20$   
 $x_1 = v(\{1\}) - v(\emptyset) = 5$ ,  $x_2 = v(\{1, 2\}) - v(\{1\}) = 15$

# Average Marginal Contribution

- **Idea:** Remove the dependence on ordering taking the **average** over all possible orderings.

# Average Marginal Contribution

- **Idea:** Remove the dependence on ordering taking the **average over all possible orderings**.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0$ ,  $v(\{1\}) = v(\{2\}) = 5$ ,  $v(\{1, 2\}) = 20$

# Average Marginal Contribution

- **Idea:** Remove the dependence on ordering taking the **average over all possible orderings**.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0$ ,  $v(\{1\}) = v(\{2\}) = 5$ ,  $v(\{1, 2\}) = 20$ 
  - 1, 2:  $x_1 = v(\{1\}) - v(\emptyset) = 5$ ,  $x_2 = v(\{1, 2\}) - v(\{1\}) = 15$
  - 2, 1:  $y_2 = v(\{2\}) - v(\emptyset) = 5$ ,  $y_1 = v(\{1, 2\}) - v(\{2\}) = 15$

# Average Marginal Contribution

- **Idea:** Remove the dependence on ordering taking the **average over all possible orderings**.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0$ ,  $v(\{1\}) = v(\{2\}) = 5$ ,  $v(\{1, 2\}) = 20$ 
  - 1, 2:  $x_1 = v(\{1\}) - v(\emptyset) = 5$ ,  $x_2 = v(\{1, 2\}) - v(\{1\}) = 15$
  - 2, 1:  $y_2 = v(\{2\}) - v(\emptyset) = 5$ ,  $y_1 = v(\{1, 2\}) - v(\{2\}) = 15$
- $z_1 = (x_1 + y_1)/2 = 10$ ,  $z_2 = (x_2 + y_2)/2 = 10$   
the resulting outcome is fair!

# Average Marginal Contribution

- **Idea:** Remove the dependence on ordering taking the **average over all possible orderings**.
- $\Gamma = (\{1, 2\}, v)$  with  
 $v(\emptyset) = 0$ ,  $v(\{1\}) = v(\{2\}) = 5$ ,  $v(\{1, 2\}) = 20$ 
  - 1, 2:  $x_1 = v(\{1\}) - v(\emptyset) = 5$ ,  $x_2 = v(\{1, 2\}) - v(\{1\}) = 15$
  - 2, 1:  $y_2 = v(\{2\}) - v(\emptyset) = 5$ ,  $y_1 = v(\{1, 2\}) - v(\{2\}) = 15$
- $z_1 = (x_1 + y_1)/2 = 10$ ,  $z_2 = (x_2 + y_2)/2 = 10$   
the resulting outcome is fair!
- Can we generalize this idea?

# Shapley Value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself  
 $\Pi(N)$  denotes the set of all permutations of  $N$



# Shapley Value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself  
 $\Pi(N)$  denotes the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$

# Shapley Value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself  
 $\Pi(N)$  denotes the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$
- For  $C \subseteq N$ , let  $\delta_i(C) = v(C \cup \{i\}) - v(C)$

# Shapley Value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself  
 $\Pi(N)$  denotes the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$
- For  $C \subseteq N$ , let  $\delta_i(C) = v(C \cup \{i\}) - v(C)$
- The **Shapley value of player  $i$**  in a game  $\Gamma = (N, v)$  with  $n$  players is

$$\Phi_i(\Gamma) = \frac{1}{n!} \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$$

# Shapley Value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself  
 $\Pi(N)$  denotes the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$
- For  $C \subseteq N$ , let  $\delta_i(C) = v(C \cup \{i\}) - v(C)$
- The **Shapley value of player  $i$**  in a game  $\Gamma = (N, v)$  with  $n$  players is

$$\Phi_i(\Gamma) = \frac{1}{n!} \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$$

- In the previous slide we have  $\Phi_1 = \Phi_2 = 10$

# Shapley Value: Probabilistic Interpretation

# Shapley Value: Probabilistic Interpretation

- $\Phi_i$  is  $i$ 's **average marginal contribution** to the coalition of its predecessors, over all permutations

# Shapley Value: Probabilistic Interpretation

- $\Phi_i$  is  $i$ 's **average marginal contribution** to the coalition of its predecessors, over all permutations
- Suppose that we choose a permutation of players uniformly at random, then  $\Phi_i$  is the **expected marginal contribution of player  $i$**  to the coalition of his predecessors

# Player's properties



# Player's properties

Given a game  $\Gamma = (N, v)$

# Player's properties

Given a game  $\Gamma = (N, v)$

- A player  $i$  is a **dummy** in  $\Gamma$  if

$$v(C) = v(C \cup \{i\}), \text{ for any } C \subseteq N$$

# Player's properties

Given a game  $\Gamma = (N, v)$

- A player  $i$  is a **dummy** in  $\Gamma$  if

$$v(C) = v(C \cup \{i\}), \text{ for any } C \subseteq N$$

- Two players  $i$  and  $j$  are said to be **symmetric** in  $\Gamma$  if

$$v(C \cup \{i\}) = v(C \cup \{j\}), \text{ for any } C \subseteq N \setminus \{i, j\}$$

# Shapley value: Axiomatic Characterization

Properties of the Shapley value:

- Efficiency:  $\Phi_1 + \dots + \Phi_n = v(N)$
- Dummy: if  $i$  is a dummy,  $\Phi_i = 0$
- Symmetry: if  $i$  and  $j$  are symmetric,  $\Phi_i = \Phi_j$
- Additivity:  $\Phi_i(\Gamma_1 + \Gamma_2) = \Phi_i(\Gamma_1) + \Phi_i(\Gamma_2)$

# Shapley value: Axiomatic Characterization

Properties of the Shapley value:

- Efficiency:  $\Phi_1 + \dots + \Phi_n = v(N)$
- Dummy: if  $i$  is a dummy,  $\Phi_i = 0$
- Symmetry: if  $i$  and  $j$  are symmetric,  $\Phi_i = \Phi_j$
- Additivity:  $\Phi_i(\Gamma_1 + \Gamma_2) = \Phi_i(\Gamma_1) + \Phi_i(\Gamma_2)$

## Theorem

*The Shapley value is the only payoff distribution scheme that has properties (1) - (4)*

$\Gamma = \Gamma_1 + \Gamma_2$  is the game  $(N, v)$  with  $v(C) = v_1(C) + v_2(C)$

# Computational Issues

- We have defined some solution concepts  
can we compute them efficiently?

# Computational Issues

- We have defined some solution concepts  
can we compute them efficiently?
- We need to determine how to represent a coalitional game  
 $\Gamma = (N, v)$

# Computational Issues

- We have defined some solution concepts  
can we compute them efficiently?
- We need to determine how to represent a coalitional game  $\Gamma = (N, v)$ 
  - **Extensive** list values of all coalitions  
exponential in the number of players  $n$
  - **Succinct** a TM describing the function  $v$   
some undecidable questions might arise



# Computational Issues

- We have defined some solution concepts  
can we compute them efficiently?
- We need to determine how to represent a coalitional game  $\Gamma = (N, v)$ 
  - **Extensive** list values of all coalitions  
exponential in the number of players  $n$
  - **Succinct** a TM describing the function  $v$   
some undecidable questions might arise
- We are usually interested in algorithms whose running time is polynomial in  $n$
- So what can we do? subclasses?

# Checking Non-emptiness of the Core: Superadditive Games

- An outcome in the core of a superadditive game satisfies the following constraints:

$$x_i \geq 0 \text{ for all } i \in N$$

$$\sum_{i \in N} x_i = v(N)$$

$$\sum_{i \in C} x_i \geq v(C), \text{ for any } C \subseteq N$$

# Checking Non-emptiness of the Core: Superadditive Games

- An outcome in the core of a superadditive game satisfies the following constraints:

$$x_i \geq 0 \text{ for all } i \in N$$

$$\sum_{i \in N} x_i = v(N)$$

$$\sum_{i \in C} x_i \geq v(C), \text{ for any } C \subseteq N$$

- A linear feasibility program, with one constraint for each coalition:  $2^n + n + 1$  constraints

# Superadditive Games: Computing the Least Core

- Starting from the linear feasibility problem for the core

$$\min \epsilon$$

$$x_i \geq 0 \text{ for all } i \in N$$

$$\sum_{i \in N} x_i = v(N)$$

$$\sum_{i \in C} x_i \geq v(C) - \epsilon, \text{ for any } C \subseteq N$$

# Superadditive Games: Computing the Least Core

- Starting from the linear feasibility problem for the core

$$\begin{aligned} \min \quad & \epsilon \\ \text{s.t.} \quad & x_i \geq 0 \text{ for all } i \in N \\ & \sum_{i \in N} x_i = v(N) \\ & \sum_{i \in C} x_i \geq v(C) - \epsilon, \text{ for any } C \subseteq N \end{aligned}$$

- A minimization program, rather than a feasibility program

# Computing Shapley Value

- $\Phi_i(\Gamma) = \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$
- $\Phi_i(\Gamma)$  is the **expected marginal contribution** of player  $i$  to the coalition of his predecessors

# Computing Shapley Value

- $\Phi_i(\Gamma) = \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$
- $\Phi_i(\Gamma)$  is the **expected marginal contribution** of player  $i$  to the coalition of his predecessors
- Quick and dirty way:

# Computing Shapley Value

- $\Phi_i(\Gamma) = \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$
- $\Phi_i(\Gamma)$  is the **expected marginal contribution** of player  $i$  to the coalition of his predecessors
- Quick and dirty way:  
Use Monte-Carlo method to compute  $\Phi_i(\Gamma)$



# Computing Shapley Value

- $\Phi_i(\Gamma) = \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$
- $\Phi_i(\Gamma)$  is the **expected marginal contribution** of player  $i$  to the coalition of his predecessors
- Quick and dirty way:  
Use Monte-Carlo method to compute  $\Phi_i(\Gamma)$   
Convergence guaranteed by Law of Large Numbers