

Stochastic Network Modeling (SNM)

Llorenç Cerdà-Alabern

llorenc@ac.upc.edu

Office: C6-213

Web page <http://docencia.ac.upc.edu/master/MIRI/SNM>

Parts

- I Introduction
- II Discrete Time Markov Chains (DTMC)
- III Continuous Time Markov Chains (CTMC)
- IV Queuing Theory

Evaluation

- $NF = 0.1 * NP + 0.30 * \max(EF, C) + 0.60 * EF$
where:
 - NF = final mark
 - EF = final theory exam
 - NP = Problems delivered by the students
 - C = average assessments mark: $C = 0.5 * C1 + 0.5 * C2$

Part I

Introduction

Outline

- Probability
- Stochastic Process (SP)



Introduction

Probability

Introduction

Probability

Ingredients of Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional Probability and Bayes Formula

Law of total probability

Probability in \mathbb{R}^k



Stochastic Process (SP)

Ingredients of Probability


- **Random experiment**, e.g. toss a die.
- **Outcome**, ω , e.g. tossing a die can be $\omega = 2$, choosing a fruit can be $\omega = \text{orange}$.
- **Sample space or Universal set**, U , set of all possible outcomes. E.g. tossing a die $U = \{1,2,3,4,5,6\}$.
- **Event**, A , any subset of U (e.g. tossing a die $A = \{1,2,3\}$). We say the event A occurs if the outcome of the experiment $\omega \in A$. U is the **sure event**, and we represent by the empty set \emptyset an **impossible outcome**.

Venn Diagrams


Graphical representation of events




Intersection $A \cap B$



Union $A \cup B$



Complement of A in U
 $A^c = U \setminus A$



Complement of A in B
(B minus A) $A^c \cap B = B \setminus A$

source: wikipedia



Introduction

Probability

Introduction

Probability

Ingredients of
Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional
Probability and Bayes
Formula

Law of total
probability

Probability in \mathbb{R}^k

Stochastic
Process (SP)

Random Variable

- For simplicity it is defined a **random variable (RV)**, X as a function that assigns a real number to each outcome in the sample space U , i.e.:

$$X: U \rightarrow \mathbb{R}$$

- We will represent the experiment by a RV, X , and the possible outcomes by its values. **$X = x_i$ is the outcome $X(\omega_i) = x_i$.**
- Using RVs the sample space is mapped in a subset of \mathbb{R} . So, in terms of X , **U is a set of points of \mathbb{R} . The same for any event.**
- Normally the definition of **X comes naturally from the experiment**, e.g. tossing a die: $X = \{\text{number in the toss}\}$.
- RVs can be **discrete** (e.g. tossing a die) or **continuous** (e.g. waiting time of a packet in a queue).



Introduction

Probability

Introduction

Probability

Ingredients of
Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional
Probability and Bayes
Formula

Law of total
probability

Probability in \mathbb{R}^k

Stochastic
Process (SP)

Probability Measure of Discrete RV

- If the sample space U of the RV X is **finite (discrete RV)**, $U = \{x_1, \dots, x_n\}$, a **probability measure** is an assignment of numbers $P(x_i)$, referred to as **probabilities**, to each **outcome x_i** such that:

$$0 \leq P(x_i) \leq 1$$

$$P(A) = \sum_{x_i \in A} P(x_i)$$

$$P(U) = 1$$

E.g. tossing a fair die,


$$P(x_i) = 1/6$$

$$P(X \in \{2, 4, 6\}) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2}$$

Conditional Probability and Bayes Formula

- Given the the sample space U and the **events** $A, B \in U$ with $P(B) > 0$ the **probability of A conditioned by B** is defined as:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$



Intersection $A \cap B$

NOTE: It's common to use commas to denote set intersection, and write $P(A \cap B)$ as $P(A, B)$.

- Bayes Formula**

$$P(A|B) P(B) = P(B|A) P(A) \Rightarrow P(A|B) = \frac{P(B|A) P(A)}{P(B)}$$



Introduction

Probability

Introduction

Probability

Ingredients of
Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional
Probability and Bayes
Formula

Law of total
probability

Probability in \mathbb{R}^k

Stochastic
Process (SP)

Law of total probability

- Let B_i a **partition** of the sample space U ($\cup_i B_i = U$, $B_i \cap B_j = \emptyset, \forall i \neq j$), then

$$P(A) = \sum_i P(A|B_i) P(B_i)$$

- For **conditional probabilities**:

$$P(A|C) = \sum_i P(A|C \cap B_i) P(B_i|C)$$

- If C is **independent** of any of the B_i

$$P(A|C) = \sum_i P(A|C \cap B_i) P(B_i)$$



Probability Measure of Continuous RV

- If the sample space of the RV X is continuous (**continuous RV**), the events are intervals of \mathbb{R} . The probability measure is defined by means of the **cumulative distribution function, CDF**:

$$F(x) = P(X \in (-\infty, x]) = P(X \leq x)$$

- X is called absolutely continuous^a if there exists the **probability density function, PDF**, such that for any interval $I = \{x \mid a \leq x \leq b\}$:

$$\int_a^b f(x) dx = P(X \in I) = F(b) - F(a)$$

^aSome special distributions, called singular, do not have a PDF. One example is the Cantor distribution (see Wikipedia).



Introduction

Probability

Introduction

Probability

Ingredients of
Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional
Probability and Bayes
Formula

Law of total
probability

Probability in \mathbb{R}^k

Stochastic
Process (SP)

Expected value

- Given the discrete $N \in \mathbb{Z}$, respectively continuous $X \in \mathbb{R}$ RV, the **expected value** is:

$$E[N] = \sum_{k=-\infty}^{\infty} k P(N = k)$$

$$E[X] = \int_{-\infty}^{\infty} x f(x) dx$$

Variance

- The amount of dispersion of a RV X with expected value $\mu = E[X]$ is measured by the **Variance**:

$$\text{Var}(X) = \sigma^2 = E[(X - \mu)^2] = E[X^2] - \mu^2$$

- Often it is used the **standard deviation** $\sigma = \sqrt{\text{Var}(X)}$.



Introduction

Probability

Introduction

Probability

Ingredients of
Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional
Probability and Bayes
Formula

Law of total
probability

Probability in \mathbb{R}^k

Stochastic
Process (SP)

Indicator Function

$$I(A) = \begin{cases} 1, & \text{if } A \text{ occurs,} \\ 0, & \text{otherwise.} \end{cases}$$

Therefore:

$$\mathbf{E}[I(A)] = 0 \times P(I(A) = 0) + 1 \times P(I(A) = 1) = \mathbf{P(A)}$$

Expected value of non negative RVs

- For **non negative** RVs, $N \geq 0$ discrete and $X \geq 0$ continuous:

$$E[N] = \sum_{k=0}^{\infty} k P(N = k) = \sum_{k=0}^{\infty} P(N > k)$$

$$E[X] = \int_0^{\infty} x f(x) dx = \int_0^{\infty} P(X > x) dx = \int_0^{\infty} (1 - F(x)) dx$$

Proof

$$N = \sum_{k=0}^{N-1} 1 = \sum_{k=0}^{\infty} I(N > k)$$

$$X = \int_0^X dx = \int_0^{\infty} I(X > x) dx$$

and take expectations.



Introduction

Probability

Introduction

Probability

Ingredients of

Probability

Venn Diagrams

Random Variable

Probability Measure

Conditional
Probability and Bayes
Formula

Law of total
probability

Probability in \mathbb{R}^k

Stochastic
Process (SP)

Wald's Equation

- Definition:** An positive integer RV $N > 0$ is a **stopping time** of a sequence X_1, X_2, \dots if the event $N = n$ is independent of X_{n+1}, X_{n+2}, \dots .

E.g. toss a die until you get 6. Let N be the number of tosses. N does not depend on the values obtained after getting 6.

- Wald's Equation** If X_1, X_2, \dots are independent and identically distributed and N is a stopping time:

$$E \left[\sum_{n=1}^N X_n \right] = E[X] E[N]$$

Wald's Equation

- Wald's Equation** If X_1, X_2, \dots are independent and identically distributed and N is a stopping time:

$$E \left[\sum_{n=1}^N X_n \right] = E[X] E[N]$$

- Proof**

$$E \left[\sum_{n=1}^N X_n \right] = E \left[\sum_{n=1}^{\infty} X_n I(n \leq N) \right] = \sum_{n=1}^{\infty} E[X_n] E[I(n \leq N)] =$$

$$E[X] \sum_{n=1}^{\infty} P(n \leq N) = E[X] \sum_{n=0}^{\infty} P(N > n) = E[X] E[N]$$

