

# Combinatorial Optimization Games

Maria

Fall 2020

- 1 Induced subgraph games
- 2 Minimum cost spanning tree games
- 3 References

- 1 Induced subgraph games
- 2 Minimum cost spanning tree games
- 3 References

# Induced subgraph games

# Induced subgraph games

- A game is described by an undirected, weighted graph  $G = (N, E)$  with  $|N| = n$  and  $|E| = m$  and an integer edge weight function  $w$ .

# Induced subgraph games

- A game is described by an undirected, weighted graph  $G = (N, E)$  with  $|N| = n$  and  $|E| = m$  and an integer edge weight function  $w$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .

# Induced subgraph games

- A game is described by an undirected, weighted graph  $G = (N, E)$  with  $|N| = n$  and  $|E| = m$  and an integer edge weight function  $w$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .
- In the game  $\Gamma(G, w) = (N, v)$  the set of players is  $N$ , and the value  $v$  of a coalition  $C \subseteq N$  is

$$v(C) = \sum_{e \in E(G[C])} w_e$$

# Induced subgraph games

- A game is described by an undirected, weighted graph  $G = (N, E)$  with  $|N| = n$  and  $|E| = m$  and an integer edge weight function  $w$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .
- In the game  $\Gamma(G, w) = (N, v)$  the set of players is  $N$ , and the value  $v$  of a coalition  $C \subseteq N$  is

$$v(C) = \sum_{e \in E(G[C])} w_e$$

- Usually **self-loops** are allowed when we want that the value of a singleton is different from 0.



# Induced subgraph games

- A game is described by an undirected, weighted graph  $G = (N, E)$  with  $|N| = n$  and  $|E| = m$  and an integer edge weight function  $w$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .
- In the game  $\Gamma(G, w) = (N, v)$  the set of players is  $N$ , and the value  $v$  of a coalition  $C \subseteq N$  is

$$v(C) = \sum_{e \in E(G[C])} w_e$$

- Usually **self-loops** are allowed when we want that the value of a singleton is different from 0.
- Observe that  $v(\emptyset) = 0$  and  $v(N) = w(E)$ .

# Induced subgraph games

# Induced subgraph games

- Induced subgraph games model aspects of social networks.

# Induced subgraph games

- Induced subgraph games model aspects of social networks.
- The value of each coalition (team, club) is determined by the relationships among its members: a player assigns a positive utility to being in a coalition with his friends and a negative utility to being in a coalition with his enemies.

# Induced subgraph games

- Induced subgraph games model aspects of social networks.
- The value of each coalition (team, club) is determined by the relationships among its members: a player assigns a positive utility to being in a coalition with his friends and a negative utility to being in a coalition with his enemies.
- The representation is succinct as long as the number of bits required to encode edge weights is polynomial in  $|N|$ : using an adjacency matrix to represent the graph requires only  $n^2$  entries.

# Induced subgraph games

- Induced subgraph games model aspects of social networks.
- The value of each coalition (team, club) is determined by the relationships among its members: a player assigns a positive utility to being in a coalition with his friends and a negative utility to being in a coalition with his enemies.
- The representation is succinct as long as the number of bits required to encode edge weights is polynomial in  $|N|$ : using an adjacency matrix to represent the graph requires only  $n^2$  entries.
- Weights can be exponential in  $n$  and still have polynomial size.

# Completeness?

# Completeness?

- Is this is a **complete** representation?



# Completeness?

- Is this is a **complete** representation?  
All simple games can be represented as induced subgraph games?

# Completeness?

- Is this is a **complete** representation?  
All simple games can be represented as induced subgraph games? **NO**

# Completeness?

- Is this is a **complete** representation?

All simple games can be represented as induced subgraph games? **NO**

Consider the game  $\Gamma = (N, v)$ , where  $n = \{1, 2, 3\}$  and

$$v(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

# Completeness?

- Is this is a **complete** representation?

All simple games can be represented as induced subgraph games? **NO**

Consider the game  $\Gamma = (N, v)$ , where  $n = \{1, 2, 3\}$  and

$$v(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .

# Completeness?

- Is this a **complete** representation?

All simple games can be represented as induced subgraph games? **NO**

Consider the game  $\Gamma = (N, v)$ , where  $n = \{1, 2, 3\}$  and

$$v(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .
  - By the first condition all self-loops must have weight 0.

# Completeness?

- Is this is a **complete** representation?

All simple games can be represented as induced subgraph games? **NO**

Consider the game  $\Gamma = (N, v)$ , where  $n = \{1, 2, 3\}$  and

$$v(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .
  - By the first condition all self-loops must have weight 0.
  - By the second condition any pair of different vertices must be connected by an edge with weight 1. So **G must be a triangle**.

# Completeness?

- Is this a **complete** representation?

All simple games can be represented as induced subgraph games? **NO**

Consider the game  $\Gamma = (N, v)$ , where  $n = \{1, 2, 3\}$  and

$$v(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .
  - By the first condition all self-loops must have weight 0.
  - By the second condition any pair of different vertices must be connected by an edge with weight 1. So  **$G$  must be a triangle.**
  - But then  $v(\{1, 2, 3\}) = 3 \neq 6$ .

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .



# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.
- Since we allow for negative edge weights, induced subgraph games are not necessarily monotone.

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.
- Since we allow for negative edge weights, induced subgraph games are not necessarily monotone.
- However, when all edge weights are non-negative, induced subgraph games are **convex**.

# Can the core be empty?

# Can the core be empty?

The **core** of  $\Gamma(N, v)$  is the set of all imputations  $x$  such that  $v(S) \leq x(S)$ , for each coalition  $S \subseteq N$ .

# Can the core be empty?

# Can the core be empty?

## Theorem

*If  $\Gamma = (N, v)$  is a convex game, then  $\Gamma$  has a non-empty core.*

# Can the core be empty?

## Theorem

*If  $\Gamma = (N, v)$  is a convex game, then  $\Gamma$  has a non-empty core.*

- Fix an arbitrary permutation  $\pi$ , and let  $x_i$  be the marginal contribution of  $i$  with respect to  $\pi$ .



# Can the core be empty?

## Theorem

*If  $\Gamma = (N, v)$  is a convex game, then  $\Gamma$  has a non-empty core.*

- Fix an arbitrary permutation  $\pi$ , and let  $x_i$  be the marginal contribution of  $i$  with respect to  $\pi$ .
- Let us show that  $(x_1, \dots, x_n)$  is in the core of  $\Gamma$ .

# Can the core be empty?

## Theorem

*If  $\Gamma = (N, v)$  is a convex game, then  $\Gamma$  has a non-empty core.*

- Fix an arbitrary permutation  $\pi$ , and let  $x_i$  be the marginal contribution of  $i$  with respect to  $\pi$ .
- Let us show that  $(x_1, \dots, x_n)$  is in the core of  $\Gamma$ .
  - For  $C \subseteq N$ , we can assume that  $C = \{i_1, \dots, i_s\}$  where  $\pi(i_1) < \dots < \pi(i_s)$ .
  - So,  $v(C) = v(\{i_1\}) - v(\emptyset) + v(\{i_1, i_2\}) - v(\{i_1\}) + \dots + v(C) - v(C \setminus \{i_s\})$ .
  - By supermodularity we have,  $v(\{i_1, \dots, i_{j-1}, i_j\}) - v(\{i_1, \dots, i_{j-1}\}) \leq v(\{1, \dots, i_j\}) - v(\{1, \dots, i_{j-1}\})$ .
  - Therefore  $v(C) \leq x(C)$  and  $v(N) = x(N)$ .

# Can the core be empty?

## Theorem

*If  $\Gamma = (N, v)$  is a convex game, then  $\Gamma$  has a non-empty core.*

- Fix an arbitrary permutation  $\pi$ , and let  $x_i$  be the marginal contribution of  $i$  with respect to  $\pi$ .
- Let us show that  $(x_1, \dots, x_n)$  is in the core of  $\Gamma$ .
  - For  $C \subseteq N$ , we can assume that  $C = \{i_1, \dots, i_s\}$  where  $\pi(i_1) < \dots < \pi(i_s)$ .
  - So,  $v(C) = v(\{i_1\}) - v(\emptyset) + v(\{i_1, i_2\}) - v(\{i_1\}) + \dots + v(C) - v(C \setminus \{i_s\})$ .
  - By supermodularity we have,  $v(\{i_1, \dots, i_{j-1}, i_j\}) - v(\{i_1, \dots, i_{j-1}\}) \leq v(\{1, \dots, i_j\}) - v(\{1, \dots, i_{j-1}\})$ .
  - Therefore  $v(C) \leq x(C)$  and  $v(N) = x(N)$ .
- Observe that we have shown that the vector formed by the Shapley value is in the core of a convex game.

# Shapley value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself

# Shapley value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself
- $\Pi(N)$  denote the set of all permutations of  $N$

# Shapley value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself
- $\Pi(N)$  denote the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$

# Shapley value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself
- $\Pi(N)$  denote the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$
- For  $C \subseteq N$ , let  $\delta_i(C) = v(C \cup \{i\}) - v(C)$

# Shapley value

- A permutation of  $\{1, \dots, n\}$  is a one-to-one mapping from  $\{1, \dots, n\}$  to itself
- $\Pi(N)$  denote the set of all permutations of  $N$
- Let  $S_\pi(i)$  denote the set of predecessors of  $i$  in  $\pi \in \Pi(N)$
- For  $C \subseteq N$ , let  $\delta_i(C) = v(C \cup \{i\}) - v(C)$
- The **Shapley value of player  $i$**  in a game  $\Gamma = (N, v)$  with  $n$  players is

$$\Phi_i(\Gamma) = \frac{1}{n!} \sum_{\pi \in \Pi(N)} \delta_i(S_\pi(i))$$



# Shapley value: Axiomatic Characterization

Properties of the Shapley value:

- Efficiency:  $\Phi_1 + \dots + \Phi_n = v(N)$
- Dummy: if  $i$  is a dummy,  $\Phi_i = 0$
- Symmetry: if  $i$  and  $j$  are symmetric,  $\Phi_i = \Phi_j$
- Additivity:  $\Phi_i(\Gamma_1 + \Gamma_2) = \Phi_i(\Gamma_1) + \Phi_i(\Gamma_2)$

# Shapley value: Axiomatic Characterization

Properties of the Shapley value:

- Efficiency:  $\Phi_1 + \dots + \Phi_n = v(N)$
- Dummy: if  $i$  is a dummy,  $\Phi_i = 0$
- Symmetry: if  $i$  and  $j$  are symmetric,  $\Phi_i = \Phi_j$
- Additivity:  $\Phi_i(\Gamma_1 + \Gamma_2) = \Phi_i(\Gamma_1) + \Phi_i(\Gamma_2)$

## Theorem

*The Shapley value is the only payoff distribution scheme that has properties (1) - (4)*

# Shapley value

# Shapley value

## Theorem

*The Shapley value of player  $i$  in  $\Gamma(G, w)$  is*

$$\Phi(i) = \frac{1}{2} \sum_{(i,j) \in E} w_{i,j}.$$

# Shapley value: Computation

# Shapley value: Computation

- Let  $\{e_1, \dots, e_m\}$  be the set of edges in  $G$ .

## Shapley value: Computation

- Let  $\{e_1, \dots, e_m\}$  be the set of edges in  $G$ .
- We can decompose the graph  $G$  into  $m$  graphs  $G_1, \dots, G_m$ , where for  $1 \leq j \leq m$  the graph  $G_j = (V, \{e_j\})$ .

## Shapley value: Computation

- Let  $\{e_1, \dots, e_m\}$  be the set of edges in  $G$ .
- We can decompose the graph  $G$  into  $m$  graphs  $G_1, \dots, G_m$ , where for  $1 \leq j \leq m$  the graph  $G_j = (V, \{e_j\})$ .
- Considering the same weight as in the original graph, let  $\Gamma_j = \Gamma(G_j, w)$ .



## Shapley value: Computation

- Let  $\{e_1, \dots, e_m\}$  be the set of edges in  $G$ .
- We can decompose the graph  $G$  into  $m$  graphs  $G_1, \dots, G_m$ , where for  $1 \leq j \leq m$  the graph  $G_j = (V, \{e_j\})$ .
- Considering the same weight as in the original graph, let  $\Gamma_j = \Gamma(G_j, w)$ .
- According to the definitions:

$$\Gamma = \Gamma(G, w) = \Gamma_1 + \dots + \Gamma_m.$$

## Shapley value: Computation

- Let  $\{e_1, \dots, e_m\}$  be the set of edges in  $G$ .
- We can decompose the graph  $G$  into  $m$  graphs  $G_1, \dots, G_m$ , where for  $1 \leq j \leq m$  the graph  $G_j = (V, \{e_j\})$ .
- Considering the same weight as in the original graph, let  $\Gamma_j = \Gamma(G_j, w)$ .
- According to the definitions:

$$\Gamma = \Gamma(G, w) = \Gamma_1 + \dots + \Gamma_m.$$

- By the additivity axiom, for each player  $i \in N$  we have

# Shapley value: Computation

- Let  $\{e_1, \dots, e_m\}$  be the set of edges in  $G$ .
- We can decompose the graph  $G$  into  $m$  graphs  $G_1, \dots, G_m$ , where for  $1 \leq j \leq m$  the graph  $G_j = (V, \{e_j\})$ .
- Considering the same weight as in the original graph, let  $\Gamma_j = \Gamma(G_j, w)$ .
- According to the definitions:

$$\Gamma = \Gamma(G, w) = \Gamma_1 + \dots + \Gamma_m.$$

- By the additivity axiom, for each player  $i \in N$  we have

$$\Phi_i(\Gamma) = \sum_{j=1}^m \Phi_i(\Gamma_j).$$

# Shapley value: Computation

# Shapley value: Computation

- We have to compute  $\Phi_i(\Gamma_j)$ .

# Shapley value: Computation

- We have to compute  $\Phi_i(\Gamma_j)$ .
- When  $i$  is not incident to  $e_j$ ,  $i$  is a dummy in  $\Gamma_j$  and  $\Phi_i(\Gamma_j) = 0$ .

# Shapley value: Computation

- We have to compute  $\Phi_i(\Gamma_j)$ .
- When  $i$  is not incident to  $e_j$ ,  $i$  is a dummy in  $\Gamma_j$  and  $\Phi_i(\Gamma_j) = 0$ .
- When  $e_j = (i, \ell)$  for some  $\ell \in N$ , players  $i$  and  $\ell$  are symmetric in  $\Gamma_j$ .

# Shapley value: Computation

- We have to compute  $\Phi_i(\Gamma_j)$ .
- When  $i$  is not incident to  $e_j$ ,  $i$  is a dummy in  $\Gamma_j$  and  $\Phi_i(\Gamma_j) = 0$ .
- When  $e_j = (i, \ell)$  for some  $\ell \in N$ , players  $i$  and  $\ell$  are symmetric in  $\Gamma_j$ .
- Since the value of the grand coalition in  $\Gamma_j$  equals  $w(i, \ell)$ , by efficiency and symmetry we get  $\Phi_i(\Gamma_j) = w(i, \ell)/2$ .



# Shapley value

# Shapley value

## Theorem

*The Shapley value of player  $i$  in  $\Gamma(G, w)$  is*

$$\Phi_i = \frac{1}{2} \sum_{(i,j) \in E} w_{i,j}.$$

# Shapley value

## Theorem

*The Shapley value of player  $i$  in  $\Gamma(G, w)$  is*

$$\Phi_i = \frac{1}{2} \sum_{(i,j) \in E} w_{i,j}.$$

## Corollary

*The Shapley values of induced subgraph games can be computed in polynomial time.*

# Can the core be empty?

## Theorem

*Consider a game  $\Gamma(G, w)$ , the following are equivalent*

- *The vector of Shapley values is in the core*
- *$(G, w)$  has no negative cut*
- *The core is non-empty*

# Can the core be empty?

# Can the core be empty?

The Shapley value is in the core iff  $G$  has no negative cut.

# Can the core be empty?

The Shapley value is in the core iff  $G$  has no negative cut.

- Let  $e(S, x) = v(S) - x(S)$  be the excess of coalition  $S$  at the imputation  $x$ .

# Can the core be empty?

The Shapley value is in the core iff  $G$  has no negative cut.

- Let  $e(S, x) = v(S) - x(S)$  be the excess of coalition  $S$  at the imputation  $x$ .
- Thus,  $x$  is in the core iff  $e(x, S) \leq 0 \ \forall S \subseteq N$ .



# Can the core be empty?

The Shapley value is in the core iff  $G$  has no negative cut.

- Let  $e(S, x) = v(S) - x(S)$  be the excess of coalition  $S$  at the imputation  $x$ .
- Thus,  $x$  is in the core iff  $e(x, S) \leq 0 \ \forall S \subseteq N$ .
- For the Shapley values,  $e(S, \Phi)$  is  $-\frac{1}{2}$  times the weight of the edges going from  $S$  to  $N \setminus S$ .

# Can the core be empty?

The Shapley value is in the core iff  $G$  has no negative cut.

- Let  $e(S, x) = v(S) - x(S)$  be the excess of coalition  $S$  at the imputation  $x$ .
- Thus,  $x$  is in the core iff  $e(x, S) \leq 0 \ \forall S \subseteq N$ .
- For the Shapley values,  $e(S, \Phi)$  is  $-\frac{1}{2}$  times the weight of the edges going from  $S$  to  $N \setminus S$ .
- Hence the Shapley value is in the core if and only if there is no negative cut  $(S, N \setminus S)$ .

# Can the core be empty?

# Can the core be empty?

The core is nonempty iff  $G$  has no negative cut.

# Can the core be empty?

The core is nonempty iff  $G$  has no negative cut.

- If  $G$  has no negative cut, the vector of Shapley values is in the core (by the previous proof).

# Can the core be empty?

The core is nonempty iff  $G$  has no negative cut.

- If  $G$  has no negative cut, the vector of Shapley values is in the core (by the previous proof).
- We have seen that if the core is non-empty, then the vector of Shapley values is in the core.

# Can the core be empty?

- **NEGATIVE-CUT:** Given a weighted graph  $(G, w)$ , determine whether there is a negative cut in  $G$ .

# Can the core be empty?

- **NEGATIVE-CUT:** Given a weighted graph  $(G, w)$ , determine whether there is a negative cut in  $G$ .

## NEGATIVE-CUT is NP-complete

- **W-MAX-CUT:** Given a weighted graph  $(G, w)$  with non-negative weights and an integer  $k$ , determine whether there is a cut of size at least  $k$  in  $G$ , is NP-complete.



# Can the core be empty?

- **NEGATIVE-CUT:** Given a weighted graph  $(G, w)$ , determine whether there is a negative cut in  $G$ .

## NEGATIVE-CUT is NP-complete

- **W-MAX-CUT:** Given a weighted graph  $(G, w)$  with non-negative weights and an integer  $k$ , determine whether there is a cut of size at least  $k$  in  $G$ , is NP-complete.
- Let  $(G, w)$  with non-negative weights and an integer  $k$ .  $G'$  is obtained as the disjoint union of  $G$  and the graph  $(\{a, b\}, \{(a, b)\})$ . Define  $w'$  as  $w'(e) = w(e)$  for  $e \in E(G)$  and  $w'((a, b)) = -k$ .

# Can the core be empty?

- **NEGATIVE-CUT:** Given a weighted graph  $(G, w)$ , determine whether there is a negative cut in  $G$ .

## NEGATIVE-CUT is NP-complete

- **W-MAX-CUT:** Given a weighted graph  $(G, w)$  with non-negative weights and an integer  $k$ , determine whether there is a cut of size at least  $k$  in  $G$ , is NP-complete.
- Let  $(G, w)$  with non-negative weights and an integer  $k$ .  $G'$  is obtained as the disjoint union of  $G$  and the graph  $(\{a, b\}, \{(a, b)\})$ . Define  $w'$  as  $w'(e) = w(e)$  for  $e \in E(G)$  and  $w'((a, b)) = -k$ .
- $G$  has a cut of size at least  $k$  iff  $G'$  has a negative cut.

# Complexity of core related problems

## Theorem

*The following problems are NP-complete:*

- *Given  $(G, w)$  and an imputation  $x$ , is it not in the core of  $\Gamma(G, w)$ ?*
- *Given  $(G, w)$ , is the vector of Shapley values of  $\Gamma(G, w)$  not in the core of  $\Gamma(G, w)$ ?*
- *Given  $(G, w)$ , is the core of  $\Gamma(G, w)$  empty?*

# Complexity of core related problems

## Theorem

*Given  $(G, w)$ , when all weights are non-negative, we can test in polynomial time*

- *whether the core is non-empty.*
- *whether an imputation  $x$  is in the core of  $\Gamma(G, w)$ .*

# Complexity of core related problems

## Theorem

*Given  $(G, w)$ , when all weights are non-negative, we can test in polynomial time*

- *whether the core is non-empty.*
- *whether an imputation  $x$  is in the core of  $\Gamma(G, w)$ .*

The first question is trivial as the vector of Shapley values belong to the core. The second problem can be solved by a reduction to MAX-FLOW.

- 1 Induced subgraph games
- 2 Minimum cost spanning tree games**
- 3 References

# MST Games

## Minimum cost spanning tree games

# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.



# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.
- $V(G) = \{v_0, \dots, v_n\}$ .

# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.
- $V(G) = \{v_0, \dots, v_n\}$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .  
We assume  $w_{i,j} \geq 0$

# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.
- $V(G) = \{v_0, \dots, v_n\}$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .  
We assume  $w_{i,j} \geq 0$
- In the game  $\Gamma(G, w) = (N, c)$  the set of players is  $N = \{v_1, \dots, v_n\}$ , and the **cost**  $c$  of a coalition  $C \subseteq N$  is  
 $c(C) =$  the weight of a minimum spanning tree of  $G[\cup\{v_0\}]$

# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.
- $V(G) = \{v_0, \dots, v_n\}$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .  
 We assume  $w_{i,j} \geq 0$
- In the game  $\Gamma(G, w) = (N, c)$  the set of players is  $N = \{v_1, \dots, v_n\}$ , and the **cost**  $c$  of a coalition  $C \subseteq N$  is  

$$c(C) = \text{the weight of a minimum spanning tree of } G[\cup\{v_0\}]$$
- Self-loops are not allowed.

# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.
- $V(G) = \{v_0, \dots, v_n\}$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .  
We assume  $w_{i,j} \geq 0$
- In the game  $\Gamma(G, w) = (N, c)$  the set of players is  $N = \{v_1, \dots, v_n\}$ , and the **cost**  $c$  of a coalition  $C \subseteq N$  is  
 $c(C) =$  the weight of a minimum spanning tree of  $G[Su\{v_0\}]$
- Self-loops are not allowed.
- The cost of a singleton coalition  $\{i\}$  is  $c(\{i\}) = w_{0,i}$ .

# MST Games

## Minimum cost spanning tree games

- A game is described by a weighted complete graph  $(G, w)$  with  $n + 1$  vertices.
- $V(G) = \{v_0, \dots, v_n\}$ .
- The weight of edge  $(i, j) \in E$  is denoted by  $w_{i,j}$ .  
 We assume  $w_{i,j} \geq 0$
- In the game  $\Gamma(G, w) = (N, c)$  the set of players is  $N = \{v_1, \dots, v_n\}$ , and the **cost**  $c$  of a coalition  $C \subseteq N$  is  

$$c(C) = \text{the weight of a minimum spanning tree of } G[S \cup \{v_0\}]$$
- Self-loops are not allowed.
- The cost of a singleton coalition  $\{i\}$  is  $c(\{i\}) = w_{0,i}$ .
- Observe that  $v(\emptyset) = 0$  and  $v(N) = w(T)$  where  $T$  is a MST of  $G$ .

# MST Games

# MST Games

- MST games model situations where a number of users must be connected to a common supplier, and the cost of such connection can be modeled as a minimum spanning tree problem.



# MST Games

- MST games model situations where a number of users must be connected to a common supplier, and the cost of such connection can be modeled as a minimum spanning tree problem.
- The representation is succinct as long as the number of bits required to encode edge weights is polynomial in  $|N|$ : using an adjacency matrix to represent the graph requires only  $n^2$  entries.

# Completeness?

# Completeness?

- Is this is a **complete** representation?

# Completeness?

- Is this is a **complete** representation? All simple games can be represented as MST games?

# Completeness?

- Is this is a **complete** representation? All simple games can be represented as MST games? **NO**

# Completeness?

- Is this is a **complete** representation? All simple games can be represented as MST games? **NO**

Consider the game  $\Gamma = (N, c)$ , where  $n = \{1, 2, 3\}$  and

$$c(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

# Completeness?

- Is this is a **complete** representation? All simple games can be represented as MST games? **NO**

Consider the game  $\Gamma = (N, c)$ , where  $n = \{1, 2, 3\}$  and

$$c(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .  $V(G) = \{0, 1, 2, 3\}$

# Completeness?

- Is this a **complete** representation? All simple games can be represented as MST games? **NO**

Consider the game  $\Gamma = (N, c)$ , where  $n = \{1, 2, 3\}$  and

$$c(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .  $V(G) = \{0, 1, 2, 3\}$ 
  - By the first condition  $w_{0,i} = 0$ , for  $i \in \{1, 2, 3\}$ .



# Completeness?

- Is this a **complete** representation? All simple games can be represented as MST games? **NO**

Consider the game  $\Gamma = (N, c)$ , where  $n = \{1, 2, 3\}$  and

$$c(C) = \begin{cases} 0 & \text{if } |C| \leq 1 \\ 1 & \text{if } |C| = 2 \\ 6 & \text{if } |C| = 3 \end{cases}$$

- Assume that  $\Gamma(G, w)$  realizes  $\Gamma$ .  $V(G) = \{0, 1, 2, 3\}$ 
  - By the first condition  $w_{0,i} = 0$ , for  $i \in \{1, 2, 3\}$ .
  - Thus, a coalition with  $|C| = 2$  has a MST with zero cost and the second condition cannot be met.

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **subadditive**  $v(C \cup D) \leq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **subadditive**  $v(C \cup D) \leq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.
- MST games are not necessarily monotone.

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **subadditive**  $v(C \cup D) \leq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.
- MST games are not necessarily monotone. Consider a triangle on  $V = \{0, 1, 2\}$  and weights  $w_{0,1} = 1$ ,  $w_{0,2} = 10$  and  $w_{1,2} = 1$   
 $c(N) = 2$  and  $c(\{1\}) = 1$  and  $c(\{2\}) = 10$

# Properties of valuations

- **monotone** if  $v(C) \leq v(D)$  for  $C \subseteq D \subseteq N$ .
- **superadditive** if  $v(C \cup D) \geq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **subadditive**  $v(C \cup D) \leq v(C) + v(D)$ , for every pair of disjoint coalitions  $C, D \subseteq N$ .
- **supermodular**  $v(C \cup D) + v(C \cap D) \geq v(C) + v(D)$ .
- A game  $(N, v)$  is **convex** iff  $v$  is supermodular.
- MST games are not necessarily monotone. Consider a triangle on  $V = \{0, 1, 2\}$  and weights  $w_{0,1} = 1$ ,  $w_{0,2} = 10$  and  $w_{1,2} = 1$   
 $c(N) = 2$  and  $c(\{1\}) = 1$  and  $c(\{2\}) = 10$
- $c$  is **subadditive**.

# Can the core be empty?

## Theorem

*Consider a MST game  $\Gamma(G, w)$ . Let  $T^*$  be a MST of  $(G, w)$  obtained using Prim's algorithm. The vector  $x = (x_1, \dots, x_n)$  that allocates to player  $i \in N$  the weight of the first edge  $i$  encounters on the (unique path) from  $v_i$  to  $v_0$  in  $T^*$  belongs to the core of  $\Gamma$ .*

Such an allocation is called **standard core allocation**

# Can the core be empty?

A standard allocation  $x$  belongs to the core

# Can the core be empty?

A standard allocation  $x$  belongs to the core

- Clearly  $\sum_{i=1}^n x_i = w(T^*) = c(N)$ .



# Can the core be empty?

A standard allocation  $x$  belongs to the core

- Clearly  $\sum_{i=1}^n x_i = w(T^*) = c(N)$ .
- We need to show that  $\sum_{i=1}^n x_i \leq c(S)$ .

# Can the core be empty?

A standard allocation  $x$  belongs to the core

- Clearly  $\sum_{i=1}^n x_i = w(T^*) = c(N)$ .
- We need to show that  $\sum_{i=1}^n x_i \leq c(S)$ .
- Consider a coalition  $S$  and let  $T$  be a MST obtained using Prim's algorithm of  $G[S \cup \{v_0\}]$ .

# Can the core be empty?

A standard allocation  $x$  belongs to the core

- Clearly  $\sum_{i=1}^n x_i = w(T^*) = c(N)$ .
- We need to show that  $\sum_{i=1}^n x_i \leq c(S)$ .
- Consider a coalition  $S$  and let  $T$  be a MST obtained using Prim's algorithm of  $G[S \cup \{v_0\}]$ .
- For  $j$  in  $S$ , let  $e_j$  be the first edge  $j$  encounters on the path from  $v_j$  to  $v_0$  in  $T$  and let  $y_j = w(e_j)$ .

# Can the core be empty?

A standard allocation  $x$  belongs to the core


- Clearly  $\sum_{i=1}^n x_i = w(T^*) = c(N)$ .
- We need to show that  $\sum_{i=1}^n x_i \leq c(S)$ .
- Consider a coalition  $S$  and let  $T$  be a MST obtained using Prim's algorithm of  $G[S \cup \{v_0\}]$ .
- For  $j$  in  $S$ , let  $e_j$  be the first edge  $j$  encounters on the path from  $v_j$  to  $v_0$  in  $T$  and let  $y_j = w(e_j)$ .
- The selected edge corresponds to the point in which Prim's algorithm connects the vertex to the component including  $v_0$ , i.e., it is a minimum weight edge in the allowed cut.

# Can the core be empty?

## A standard allocation $x$ belongs to the core

- Clearly  $\sum_{i=1}^n x_i = w(T^*) = c(N)$ .
- We need to show that  $\sum_{i=1}^n x_i \leq c(S)$ .
- Consider a coalition  $S$  and let  $T$  be a MST obtained using Prim's algorithm of  $G[S \cup \{v_0\}]$ .
- For  $j$  in  $S$ , let  $e_j$  be the first edge  $j$  encounters on the path from  $v_j$  to  $v_0$  in  $T$  and let  $y_j = w(e_j)$ .
- The selected edge corresponds to the point in which Prim's algorithm connects the vertex to the component including  $v_0$ , i.e., it is a minimum weight edge in the allowed cut.
- Analyzing carefully both executions it can be shown that  $x_j \leq y_j$  as the edges considered in one partition are a subset of the other.

# How fair are standard core allocations?



- Most of the cost is charged to player 1.
- How to find more appropriate core allocations?

## More appropriate core allocations?

- There are many proposals to try to get more appropriate core allocations.

## More appropriate core allocations?

- There are many proposals to try to get more appropriate core allocations.
- Granot and Huberman [1984] propose the **weak demand allocation** and **strong demand allocation** procedures. Which rectify standard allocations by transferring cost (whenever possible) from one node to their children.



## More appropriate core allocations?

- There are many proposals to try to get more appropriate core allocations.
- Granot and Huberman [1984] propose the **weak demand allocation** and **strong demand allocation** procedures. Which rectify standard allocations by transferring cost (whenever possible) from one node to their children.
- Norde, Moretti and Tijs [2001] show how to find a **population monotonic allocation scheme** (PMAS), which is an allocation scheme that provides a core element for the game and all its subgames and which, moreover, satisfies a monotonicity condition in the sense that players have to pay less in larger coalitions.

# Complexity of core related problems

# Complexity of core related problems

## Theorem

*The following problem is NP-complete:*

- *Given  $(G, w)$  and an imputation  $x$ , is it not in the core of  $\Gamma(G, w)$ ?*

# Complexity of core related problems

## Theorem

*The following problem is NP-complete:*

- *Given  $(G, w)$  and an imputation  $x$ , is it not in the core of  $\Gamma(G, w)$ ?*

The proof follows by a reduction from EXACT COVER BY 3-SETS  
[Faigle et al., Int. J. Game Theory 1997]

- 1 Induced subgraph games
- 2 Minimum cost spanning tree games
- 3 References**

# References

- X. Deng and C. Papadimitriou. *On the complexity of cooperative solution concepts*. Mathematics of Operations Research, 19(2):257–266, 1994.
- C. G. Bird. *On cost allocation for a spanning tree: A game theory approach*. Networks, 6:335–350, 1976.
- U. Faigle, W. Kern, S. P. Fekete, and W. Hochstättler. *On the complexity of testing membership in the core of min-cost spanning tree games*. International Journal of Game Theory, 26:361–366, 1997.

# References

- G. Chalkiadakis, E. Elkind, M. Wooldridge. [Computational Aspects of Cooperative Game Theory](#) Synthesis Lectures on Artificial Intelligence and Machine Learning, Morgan & Claypool, October 2011.