

Simple Games

Fall 2020

- 1 Simple Games
- 2 Problems on simple games

1 Simple Games

2 Problems on simple games

Simple Games

- Simple Games were introduced by (Taylor & Zwicker, 1999)

Simple Games

- Simple Games were introduced by (Taylor & Zwicker, 1999)
- A **simple game** is a cooperative game (N, v) such that $v : \mathcal{C}_N \rightarrow \{0, 1\}$ and it is monotone.

Simple Games

- Simple Games were introduced by (Taylor & Zwicker, 1999)
- A **simple game** is a cooperative game (N, v) such that $v : \mathcal{C}_N \rightarrow \{0, 1\}$ and it is monotone.
- A simple game can be described by a pair (N, \mathcal{W}) :
 - N is a set of players,
 - $\mathcal{W} \subseteq \mathcal{P}(N)$ is a monotone set of **winning coalitions**, those coalitions X with $v(X) = 1$.
 - $\mathcal{L} = \mathcal{C}_N \setminus \mathcal{W}$ is the set of **losing coalitions** those coalitions X with $v(X) = 0$.

Simple Games

- Simple Games were introduced by (Taylor & Zwicker, 1999)
- A **simple game** is a cooperative game (N, v) such that $v : \mathcal{C}_N \rightarrow \{0, 1\}$ and it is monotone.
- A simple game can be described by a pair (N, \mathcal{W}) :
 - N is a set of players,
 - $\mathcal{W} \subseteq \mathcal{P}(N)$ is a monotone set of **winning coalitions**, those coalitions X with $v(X) = 1$.
 - $\mathcal{L} = \mathcal{C}_N \setminus \mathcal{W}$ is the set of **losing coalitions** those coalitions X with $v(X) = 0$.
- Members of $N = \{1, \dots, n\}$ are called **players** or **voters**.

Simple games: Representation

Due to monotonicity, any one of the following families of coalitions define a simple game on a set of players N :

- *winning coalitions* \mathcal{W} .
- *losing coalitions* \mathcal{L} .
- *minimal winning coalitions* \mathcal{W}^m
$$\mathcal{W}^m = \{X \in \mathcal{W}; \forall Z \in \mathcal{W}, Z \not\subseteq X\}$$
- *maximal losing coalitions* \mathcal{L}^M
$$\mathcal{L}^M = \{X \in \mathcal{L}; \forall Z \in \mathcal{L}, X \not\subseteq Z\}$$

This provides us with many representation forms for simple games.

Weighted voting games

Weighted voting games

- Weighted voting games (WVG)

Weighted voting games

- Weighted voting games (WVG)

A simple game for which there exists a **quota** q and it is possible to assign to each $i \in N$ a **weight** w_i , so that

$$X \in \mathcal{W} \text{ iff } \sum_{i \in X} w_i \geq q.$$

Weighted voting games

- Weighted voting games (WVG)

A simple game for which there exists a **quota** q and it is possible to assign to each $i \in N$ a **weight** w_i , so that

$$X \in \mathcal{W} \text{ iff } \sum_{i \in X} w_i \geq q.$$

- WVG can be represented by a tuple of integers

$(q; w_1, \dots, w_n)$.

as **any weighted game admits such an integer realization**,

[Carreras and Freixas, Math. Soc.Sci., 1996]

Vector weighted voting games

Vector weighted voting games

Vector weighted voting games

- Given two simple games $\Gamma_1 = (N, \mathcal{W}_1)$ and $\Gamma_2 = (N, \mathcal{W}_2)$ we can define their

Vector weighted voting games

- Given two simple games $\Gamma_1 = (N, \mathcal{W}_1)$ and $\Gamma_2 = (N, \mathcal{W}_2)$ we can define their
 - union

Vector weighted voting games

- Given two simple games $\Gamma_1 = (N, \mathcal{W}_1)$ and $\Gamma_2 = (N, \mathcal{W}_2)$ we can define their
 - union** $\Gamma_1 \cup \Gamma_2 = (N, \mathcal{W}_1 \cup \mathcal{W}_2)$

Vector weighted voting games

- Given two simple games $\Gamma_1 = (N, \mathcal{W}_1)$ and $\Gamma_2 = (N, \mathcal{W}_2)$ we can define their
 - union** $\Gamma_1 \cup \Gamma_2 = (N, \mathcal{W}_1 \cup \mathcal{W}_2)$
 - intersection**

Vector weighted voting games

- Given two simple games $\Gamma_1 = (N, \mathcal{W}_1)$ and $\Gamma_2 = (N, \mathcal{W}_2)$ we can define their
 - union** $\Gamma_1 \cup \Gamma_2 = (N, \mathcal{W}_1 \cup \mathcal{W}_2)$
 - intersection** $\Gamma_1 \cap \Gamma_2 = (N, \mathcal{W}_1 \cap \mathcal{W}_2)$

Both are simple games

Vector weighted voting games

- Given two simple games $\Gamma_1 = (N, \mathcal{W}_1)$ and $\Gamma_2 = (N, \mathcal{W}_2)$ we can define their
 - union** $\Gamma_1 \cup \Gamma_2 = (N, \mathcal{W}_1 \cup \mathcal{W}_2)$
 - intersection** $\Gamma_1 \cap \Gamma_2 = (N, \mathcal{W}_1 \cap \mathcal{W}_2)$

Both are simple games

- A simple game Γ is a **vector weighted voting game** if there are WVGs $\Gamma_1, \dots, \Gamma_k$, for some $k \geq 1$, so that $\Gamma = \Gamma_1 \cap \dots \cap \Gamma_k$.

Representability

Representability

- There are simple games that are not WVGs

Representability

- There are simple games that are not WVGs

The game with $N = \{1, 2, 3, 4\}$ where the minimal winning coalitions are the sets $\{1, 2\}$ and $\{3, 4\}$ is not a WVG.

Representability

- There are simple games that are not WVGs

The game with $N = \{1, 2, 3, 4\}$ where the minimal winning coalitions are the sets $\{1, 2\}$ and $\{3, 4\}$ is not a WVG.

- Assume it is given by $(q; w_1, w_2, w_3, w_4)$.

Representability

- There are simple games that are not WVGs

The game with $N = \{1, 2, 3, 4\}$ where the minimal winning coalitions are the sets $\{1, 2\}$ and $\{3, 4\}$ is not a WVG.

- Assume it is given by $(q; w_1, w_2, w_3, w_4)$.
- We have $w_1 + w_2 \geq q$ and $w_3 + w_4 \geq q$.
- Thus $\max\{w_1, w_2\} \geq q/2$ and $\max\{w_3, w_4\} \geq q/2$,
- So, $\max\{w_1, w_2\} + \max\{w_3, w_4\} \geq q$ which cannot be.

Representability

Representability

- Any simple game Γ is a VWVGs

Representability

- Any simple game Γ is a VWVGs
 - Take a losing coalition C and consider the game in which players in C have weight 0 and players outside C 1, set the quote to 1.
Any set that is not contained in C wins!

Representability

- Any simple game Γ is a VWVGs
 - Take a losing coalition C and consider the game in which players in C have weight 0 and players outside C 1, set the quote to 1.
Any set that is not contained in C wins!
 - The intersection of the above games describes Γ .

Representability

- Any simple game Γ is a VWVGs
 - Take a losing coalition C and consider the game in which players in C have weight 0 and players outside C 1, set the quote to 1.
Any set that is not contained in C wins!
 - The intersection of the above games describes Γ .
A winning coalition cannot be a subset of any losing coalition.

Representability

- Any simple game Γ is a VWVGs
 - Take a losing coalition C and consider the game in which players in C have weight 0 and players outside C 1, set the quote to 1.
Any set that is not contained in C wins!
 - The intersection of the above games describes Γ .
A winning coalition cannot be a subset of any losing coalition.
- The **dimension** of a simple games is the minimum number of WVGs that allows its representation as VWVG

A representation as WVGs

A representation as WVGs

The game Γ with $N = \{1, 2, 3, 4\}$ where the minimal winning coalitions are the sets $\{1, 2\}$ and $\{3, 4\}$ is not a WVG.

A representation as WVGs

The game Γ with $N = \{1, 2, 3, 4\}$ where the minimal winning coalitions are the sets $\{1, 2\}$ and $\{3, 4\}$ is not a WVG.

- The maximal losing coalitions are $\{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$

A representation as WVGs

The game Γ with $N = \{1, 2, 3, 4\}$ where the minimal winning coalitions are the sets $\{1, 2\}$ and $\{3, 4\}$ is not a WVG.

- The maximal losing coalitions are $\{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$
- This gives four WVG, according to the previous construction

$$\Gamma = [1; 0, 1, 0, 1] \cap [1; 0, 1, 1, 0] \cap [1; 1, 0, 0, 1] \cap [1; 1, 0, 1, 0].$$

Input representations

- Simple Games
 (N, \mathcal{W}) : extensive winning, (N, \mathcal{W}^m) : minimal winning
 (N, \mathcal{L}) : extensive losing, (N, \mathcal{L}^M) maximal losing
 (N, C) : monotone circuit winning
 (N, F) : monotone formula winning,
- Weighted voting games: $(q; w_1, \dots, w_n)$
- Vector weighted voting games:
 $(q_1; w_1^1, \dots, w_n^1), \dots, (q_k; w_1^k, \dots, w_n^k)$

All numbers are integers

The core of simple games

The core of simple games

- It is standard to assume that the grand coalition forms, even if the simple game is not superadditive.
- A player is a **veto player** if $v(C) = 0$, for any $C \subseteq N \setminus \{i\}$.
- Ex: Consider the unanimity game (N, v) where $v(C) = 0$, if $C \neq N$ and $v(N) = 1$.

The game indeed is a simple game and can be described in (minimal) winning form by $(N, \{N\})$.

In the unanimity game all players are veto players.

The core of simple games

Theorem

A simple game has non-empty core iff it has a veto player.

The core of simple games

Theorem

A simple game has non-empty core iff it has a veto player.

- If Γ has a veto player i .
 - Consider the payoff $x_i = 1$ and $x_j = 0$, for $j \neq i$
 - For C with $i \in C$, $v(C) = 1$ and $x(C) = 1$.
 - For C with $i \notin C$, $v(C) = 0$ and $x(C) = 0$.
 - Thus, x is in the core.
- If Γ does not have a veto player and non-empty core.
 - Consider a payoff x that is in the core.
 - $x(N) = v(N) = 1$, so there exists i with $x_i > 0$.
 - So, $x(N \setminus \{i\}) < 1$. But, $v(N \setminus \{i\}) = 1$ as i is not a veto player.
 - Thus, x is not in the core.

Is the core empty?

- Determining if the core is empty or not can be done by checking for every player whether it is a veto player or not.
- For this it is enough to check whether $v(N \setminus \{i\}) = 0$.
- For reasonable v , polynomial time computable, this can be done in poly time

Shapley value and Banzhaf index

- Player i is **pivotal** for coalition C if $v(C) = 1$ and $v(C \setminus \{i\}) = 0$.
- The sum counts those the terms for which the player is pivotal.

$$\varphi_i(\Gamma) = \frac{1}{n!} |\{i \text{ is pivotal for } S_\pi(i)\}|$$

- $\varphi_i(\Gamma)$ is the probability that the arrival of player i turns a losing coalition into a winning one.
- The Banzhaf value gives the probability of this fact over **random** coalitions.
Players in $N \setminus \{i\}$ select to be or not in the coalition tossing a fair coin.

1 Simple Games

2 Problems on simple games

Problems on simple games

In general we state a property P , for simple games, and consider the associated decision problem which has the form:

Name: IsP

Input: A simple game/WVG/VWVG Γ

Question: Does Γ satisfy property P ?

Four properties

A simple game (N, \mathcal{W}) is

- **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$.
- **proper** if $S \in \mathcal{W}$ implies $N \setminus S \notin \mathcal{W}$.
- a **weighted voting game**.
- a **vector weighted voting game**.

IsStrong: Simple Games

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

IsStrong: Simple Games

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

Theorem

The ISSTRONG problem, when Γ is given in explicit winning or losing form or in maximal losing form can be solved in polynomial time.

IsStrong: Simple Games

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

Theorem

The ISSTRONG problem, when Γ is given in explicit winning or losing form or in maximal losing form can be solved in polynomial time.

- First observe that, given a family of subsets F , we can check, for any set in F , whether its complement is not in F in polynomial time.
- Therefore, the ISSTRONG problem, when the input is given in explicit winning or losing form is polynomial time solvable.

IsStrong: Simple Games losing forms

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

- A simple game is not strong iff

$$\exists S \subseteq N : S \in \mathcal{L} \wedge N \setminus S \in \mathcal{L}$$

IsStrong: Simple Games losing forms

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

- A simple game is not strong iff

$$\exists S \subseteq N : S \in \mathcal{L} \wedge N \setminus S \in \mathcal{L}$$

which is equivalent to

$$\exists S \subseteq N : \exists L_1, L_2 \in \mathcal{L}^M : S \subseteq L_1 \wedge N \setminus S \subseteq L_2$$

IsStrong: Simple Games losing forms

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

- A simple game is not strong iff

$$\exists S \subseteq N : S \in \mathcal{L} \wedge N \setminus S \in \mathcal{L}$$

which is equivalent to

$$\exists S \subseteq N : \exists L_1, L_2 \in \mathcal{L}^M : S \subseteq L_1 \wedge N \setminus S \subseteq L_2$$

- which is equivalent to there are two maximal losing coalitions L_1 and L_2 such that $L_1 \cup L_2 = N$.

IsStrong: Simple Games losing forms

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

- A simple game is not strong iff

$$\exists S \subseteq N : S \in \mathcal{L} \wedge N \setminus S \in \mathcal{L}$$

which is equivalent to

$$\exists S \subseteq N : \exists L_1, L_2 \in \mathcal{L}^M : S \subseteq L_1 \wedge N \setminus S \subseteq L_2$$

- which is equivalent to there are two maximal losing coalitions L_1 and L_2 such that $L_1 \cup L_2 = N$.
- This can be checked in polynomial time, given \mathcal{L}^M .



IsStrong: minimal winning forms

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

Theorem

The ISSTRONG problem is coNP-complete when the input game is given in explicit minimal winning form.

IsStrong: minimal winning forms

Γ is **strong** if $S \notin \mathcal{W}$ implies $N \setminus S \in \mathcal{W}$

Theorem

The ISSTRONG problem is coNP-complete when the input game is given in explicit minimal winning form.

- The property can be expressed as

$$\forall S [(S \in \mathcal{W}) \text{ or } (S \notin \mathcal{W} \text{ and } N \setminus S \in \mathcal{W})]$$

- Observe that the property $S \in \mathcal{W}$ can be checked in polynomial time given S and \mathcal{W}^m .
- Thus the problem belongs to coNP.

IsStrong: minimal winning forms

- We provide a polynomial time reduction from the complement of the NP-complete **set splitting** problem.
- An instance of the **set splitting problem** is a collection C of subsets of a finite set N . The question is whether it is possible to partition N into two subsets P and $N \setminus P$ such that no subset in C is entirely contained in either P or $N \setminus P$.

IsStrong: minimal winning forms

- We provide a polynomial time reduction from the complement of the NP-complete **set splitting** problem.
- An instance of the **set splitting problem** is a collection C of subsets of a finite set N . The question is whether it is possible to partition N into two subsets P and $N \setminus P$ such that no subset in C is entirely contained in either P or $N \setminus P$.
- We have to decide whether $P \subseteq N$ exists such that

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

IsStrong: minimal winning forms

- We provide a polynomial time reduction from the complement of the NP-complete **set splitting** problem.
- An instance of the **set splitting problem** is a collection C of subsets of a finite set N . The question is whether it is possible to partition N into two subsets P and $N \setminus P$ such that no subset in C is entirely contained in either P or $N \setminus P$.
- We have to decide whether $P \subseteq N$ exists such that

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

We associate to a set splitting instance (N, C) the simple game in explicit minimal winning form (N, C^m) .

IsStrong: minimal winning forms

- C^m can be computed in polynomial time, given C . Why?

IsStrong: minimal winning forms

- C^m can be computed in polynomial time, given C . Why?
- Now assume that $P \subseteq N$ satisfies

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

IsStrong: minimal winning forms

- C^m can be computed in polynomial time, given C . Why?
- Now assume that $P \subseteq N$ satisfies

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

- This means that P and $N \setminus P$ are losing coalitions in the game (N, C^m) .

IsStrong: minimal winning forms

- C^m can be computed in polynomial time, given C . Why?
- Now assume that $P \subseteq N$ satisfies

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

- This means that P and $N \setminus P$ are losing coalitions in the game (N, C^m) .
- So, $S \not\subseteq P$ and $S \not\subseteq N \setminus P$, for any $S \in C^m$.

IsStrong: minimal winning forms

- C^m can be computed in polynomial time, given C . Why?
- Now assume that $P \subseteq N$ satisfies

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

- This means that P and $N \setminus P$ are losing coalitions in the game (N, C^m) .
- So, $S \not\subseteq P$ and $S \not\subseteq N \setminus P$, for any $S \in C^m$.
- This implies $S \not\subseteq P$ and $S \not\subseteq N \setminus P$, for any $S \in C$ since any set in C contains a set in C^m .

IsStrong: minimal winning forms

- C^m can be computed in polynomial time, given C . Why?
- Now assume that $P \subseteq N$ satisfies

$$\forall S \in C : S \not\subseteq P \wedge S \not\subseteq N \setminus P$$

- This means that P and $N \setminus P$ are losing coalitions in the game (N, C^m) .
- So, $S \not\subseteq P$ and $S \not\subseteq N \setminus P$, for any $S \in C^m$.
- This implies $S \not\subseteq P$ and $S \not\subseteq N \setminus P$, for any $S \in C$ since any set in C contains a set in C^m .
- Therefore, (N, C) has a set splitting iff (N, C^m) is not proper.

IsProper: winning forms

Γ is **proper** if $S \in \mathcal{W}$ implies $N \setminus S \notin \mathcal{W}$.

Theorem

The ISPROPER problem, when the game is given in explicit winning or losing form or in minimal winning form, can be solved in polynomial time.

IsProper: winning forms

Γ is **proper** if $S \in \mathcal{W}$ implies $N \setminus S \notin \mathcal{W}$.

Theorem

The ISPROPER problem, when the game is given in explicit winning or losing form or in minimal winning form, can be solved in polynomial time.

- As before, given a family of subsets F , we can check, for any set in F , whether its complement is not in F in polynomial time.

Taking into account the definitions, the ISPROPER problem is polynomial time solvable for the explicit forms

Is Proper: winning forms

- Γ is not proper iff

$$\exists S \subseteq N : S \in \mathcal{W} \wedge N \setminus S \in \mathcal{W}$$

Is Proper: winning forms

- Γ is not proper iff

$$\exists S \subseteq N : S \in \mathcal{W} \wedge N \setminus S \in \mathcal{W}$$

- which is equivalent to

$$\exists S \subseteq N : \exists W_1, W_2 \in \mathcal{W}^m : W_1 \subseteq S \wedge W_2 \subseteq N \setminus S.$$

Is Proper: winning forms

- Γ is not proper iff

$$\exists S \subseteq N : S \in \mathcal{W} \wedge N \setminus S \in \mathcal{W}$$

- which is equivalent to

$$\exists S \subseteq N : \exists W_1, W_2 \in \mathcal{W}^m : W_1 \subseteq S \wedge W_2 \subseteq N \setminus S.$$

- equivalent to there are two minimal winning coalitions W_1 and W_2 such that $W_1 \cap W_2 = \emptyset$.

Is Proper: winning forms

- Γ is not proper iff

$$\exists S \subseteq N : S \in \mathcal{W} \wedge N \setminus S \in \mathcal{W}$$

- which is equivalent to

$$\exists S \subseteq N : \exists W_1, W_2 \in \mathcal{W}^m : W_1 \subseteq S \wedge W_2 \subseteq N \setminus S.$$

- equivalent to there are two minimal winning coalitions W_1 and W_2 such that $W_1 \cap W_2 = \emptyset$.
- Which can be checked in polynomial time when \mathcal{W}^m is given.

IsProper: maximal losing form

Γ is **proper** if $S \in \mathcal{W}$ implies $N \setminus S \notin \mathcal{W}$.

Theorem

The IsProper problem is coNP-complete when the input game is given in extensive maximal losing form.

IsProper: maximal losing form

Γ is **proper** if $S \in \mathcal{W}$ implies $N \setminus S \notin \mathcal{W}$.

Theorem

The ISPROPER problem is coNP-complete when the input game is given in extensive maximal losing form.

- A game is *not proper* iff

$$\exists S \subseteq N : S \notin \mathcal{L} \wedge N \setminus S \notin \mathcal{L}$$

- which is equivalent to

$$\exists S \subseteq N : \forall T_1, T_2 \in \mathcal{L}^M : S \not\subseteq T_1 \wedge N \setminus S \not\subseteq T_2$$

IsProper: maximal losing form

Γ is **proper** if $S \in \mathcal{W}$ implies $N \setminus S \notin \mathcal{W}$.

Theorem

The ISPROPER problem is coNP-complete when the input game is given in extensive maximal losing form.

- A game is *not proper* iff

$$\exists S \subseteq N : S \notin \mathcal{L} \wedge N \setminus S \notin \mathcal{L}$$

- which is equivalent to

$$\exists S \subseteq N : \forall T_1, T_2 \in \mathcal{L}^M : S \not\subseteq T_1 \wedge N \setminus S \not\subseteq T_2$$

- Therefore ISPROPER belongs to coNP.

IsProper: maximal losing form

To show that the problem is also coNP-hard we provide a reduction from the IsStrong problem for games given in extensive minimal winning form.

IsProper: maximal losing form

To show that the problem is also coNP-hard we provide a reduction from the IsStrong problem for games given in extensive minimal winning form.

- If a family C of subsets of N is minimal then the family $\{N \setminus L : L \in C\}$ is maximal.
- Given a game $\Gamma = (N, \mathcal{W}^m)$, in minimal winning form, we construct the game $\Gamma' = (N, \{N \setminus L : L \in \mathcal{W}^m\})$ in maximal losing form.
- Which can be obtained in polynomial time.

IsProper: maximal losing form

To show that the problem is also coNP-hard we provide a reduction from the IsStrong problem for games given in extensive minimal winning form.

- If a family C of subsets of N is minimal then the family $\{N \setminus L : L \in C\}$ is maximal.
- Given a game $\Gamma = (N, \mathcal{W}^m)$, in minimal winning form, we construct the game $\Gamma' = (N, \{N \setminus L : L \in \mathcal{W}^m\})$ in maximal losing form.
- Which can be obtained in polynomial time.
- Besides, Γ is strong iff Γ' is proper.

Weighted voting games

Weighted voting games

Some of the proofs are based on reductions from the NP-complete problem PARTITION:

Weighted voting games

Some of the proofs are based on reductions from the NP-complete problem PARTITION:

Name: PARTITION

Input: n integer values, x_1, \dots, x_n

Question: Is there $S \subseteq \{1, \dots, n\}$ for which

$$\sum_{i \in S} x_i = \sum_{i \notin S} x_i.$$

Weighted voting games

Some of the proofs are based on reductions from the NP-complete problem PARTITION:

Name: PARTITION

Input: n integer values, x_1, \dots, x_n

Question: Is there $S \subseteq \{1, \dots, n\}$ for which

$$\sum_{i \in S} x_i = \sum_{i \notin S} x_i.$$

Observe that, for any instance of the PARTITION problem in which the sum of the n input numbers is odd, the answer must be NO.

Weighted voting games

Theorem

The ISSTRONG and the ISPROPER problems, when the input is described by an integer realization of a weighted game $(q; w)$, are coNP-complete.

Weighted voting games

Theorem

The ISSTRONG and the ISPROPER problems, when the input is described by an integer realization of a weighted game $(q; w)$, are coNP-complete.

- From the definitions of strong, proper it is straightforward to show that both problems belong to coNP.

Weighted voting games

Theorem

The ISSTRONG and the ISPROPER problems, when the input is described by an integer realization of a weighted game $(q; w)$, are coNP-complete.

- From the definitions of strong, proper it is straightforward to show that both problems belong to coNP.
- Observe that the weighted game with integer representation $(2; 1, 1, 1)$ is both proper and strong.

Hardness

We transform an instance $x = (x_1, \dots, x_n)$ of PARTITION into a realization of a weighted game according to the following schema

$$f(x) = \begin{cases} (q(x); x) & \text{when } x_1 + \dots + x_n \text{ is even,} \\ (2; 1, 1, 1) & \text{otherwise.} \end{cases}$$

Hardness

We transform an instance $x = (x_1, \dots, x_n)$ of PARTITION into a realization of a weighted game according to the following schema

$$f(x) = \begin{cases} (q(x); x) & \text{when } x_1 + \dots + x_n \text{ is even,} \\ (2; 1, 1, 1) & \text{otherwise.} \end{cases}$$

- Function f can be computed in polynomial time provided q does.

Hardness

We transform an instance $x = (x_1, \dots, x_n)$ of PARTITION into a realization of a weighted game according to the following schema

$$f(x) = \begin{cases} (q(x); x) & \text{when } x_1 + \dots + x_n \text{ is even,} \\ (2; 1, 1, 1) & \text{otherwise.} \end{cases}$$

- Function f can be computed in polynomial time provided q does.
- Independently of q , when $x_1 + \dots + x_n$ is *odd*, x is a NO input for partition, but $f(x)$ is a YES instance of ISSTRONG or ISPROPER.

IsStrong

Assume that $x_1 + \cdots + x_n$ is even.

Let $s = (x_1 + \cdots + x_n)/2$ and $N = \{1, \dots, n\}$.

Set $q(x) = s + 1$.

IsStrong

Assume that $x_1 + \dots + x_n$ is even.

Let $s = (x_1 + \dots + x_n)/2$ and $N = \{1, \dots, n\}$.

Set $q(x) = s + 1$.

- If there is $S \subset N$ such that $\sum_{i \in S} x_i = s$, then $\sum_{i \notin S} x_i = s$, thus both S and $N \setminus S$ are losing coalitions and $f(x)$ is not strong.

IsStrong

Assume that $x_1 + \dots + x_n$ is even.

Let $s = (x_1 + \dots + x_n)/2$ and $N = \{1, \dots, n\}$.

Set $q(x) = s + 1$.

- If there is $S \subset N$ such that $\sum_{i \in S} x_i = s$, then $\sum_{i \notin S} x_i = s$, thus both S and $N \setminus S$ are losing coalitions and $f(x)$ is not strong.
- If S and $N \setminus S$ are losing coalitions in $f(x)$.
If $\sum_{i \in S} x_i < s$ then $\sum_{i \notin S} x_i \geq s + 1$, $N \setminus S$ should be winning.
Thus $\sum_{i \in S} x_i = \sum_{i \notin S} x_i = s$, and there exists a partition of x .

IsProper

Assume that $x_1 + \cdots + x_n$ is even.

Let $s = (x_1 + \cdots + x_n)/2$ and $N = \{1, \dots, n\}$.

Set $q(x) = s$.

Is Proper

Assume that $x_1 + \dots + x_n$ is even.

Let $s = (x_1 + \dots + x_n)/2$ and $N = \{1, \dots, n\}$.

Set $q(x) = s$.

- If there is $S \subset N$ such that $\sum_{i \in S} x_i = s$, then $\sum_{i \notin S} x_i = s$, thus both S and $N \setminus S$ are winning coalitions and $f(x)$ is not proper.

IsProper

Assume that $x_1 + \dots + x_n$ is even.

Let $s = (x_1 + \dots + x_n)/2$ and $N = \{1, \dots, n\}$.

Set $q(x) = s$.

- If there is $S \subset N$ such that $\sum_{i \in S} x_i = s$, then $\sum_{i \notin S} x_i = s$, thus both S and $N \setminus S$ are winning coalitions and $f(x)$ is not proper.
- When $f(x)$ is not proper

$$\exists S \subseteq N : \sum_{i \in S} x_i \geq s \wedge \sum_{i \notin S} x_i \geq s,$$

and thus $\sum_{i \in S} x_i = s$.