

# A Report of *Type Theory and Formal Proof*

Juan Pablo Royo Sales

Universitat Politècnica de Catalunya

February 9, 2021

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Untyped lambda calculus</b>	<b>1</b>
2.1	Definition . . . . .	2
	<b>References</b>	<b>2</b>

## 1 Introduction

This report is going to provide a summary over the book [NG14]. Alongside the different chapters of the book I am going to describe briefly the most important parts of each chapter and, at the same time, I am going to solve 1 or 2 of the exercises proposed by the authors.

The organization of the report is going to be the same as the chapters of the book.

## 2 Untyped lambda calculus

In this first chapter the authors define and describe Lambda Calculus ( $\lambda$ -*calculus*) system which encapsulates the formalization of basic aspects of mathematical functions, in particular construction and use. In  $\lambda$ -*calculus* formalization system there are *typed* and *untyped* formalization of the same system. In this first case authors introduced the first basic and simple formalization which is *untyped*.

## 2.1 Definition

There are *two constructions principles* and *one evaluation rule*

### Construction principles:

- *Abstraction:* Given an expression  $M$  and a variable  $x$  we can construct the expression:  $\lambda x.M$ . This is abstraction of  $x$  over  $M$  Example:  $\lambda y.(\lambda x.x - y)$  Abstraction of  $y$  over  $\lambda x.x - y$
- *Application:* Given 2 expressions  $M$  and  $N$  we can construct the expression:  $M N$ . This is the application of  $M$  to  $N$ . Example:  $(\lambda x.x^2 + 1)(3)$  Application of 3 over  $\lambda x.x^2 + 1$

**Evaluation Rule:** Formalization of this process is called Beta Reduction ( $\beta$ -reduction). Makes use of substitution, expressed by  $[ ]$ .  $\beta$ -reduction: An expression  $(\lambda x.M)N$  can be rewritten to  $M[x := N]$ , which means every  $x$  should be replaced by  $N$  in  $M$ . This process is called  $\beta$ -reduction of  $(\lambda x.M)N$  to  $M[x := N]$ .

Example:  $(\lambda x.x^2 + 1)(3)$  reduces to  $(x^2 + 1)[x := 3]$ , which is  $3^2 + 1$ .

## References

- [NG14] Rob Nederpelt and Herman Geuvers. *Type Theory and Formal Proof*. Cambridge University Press, Cambridge CB2 8BS, United Kingdom, 2014.