

# Stochastic Network Modeling

## Homework 7 - Solutions

Juan Pablo Royo Sales  
Universitat Politècnica de Catalunya

October 8, 2020

### Problem 7.1

#### 7.1.1

The period of the chain is  $d = 3$  and the cyclic classes are:  $C_0 = \{a, d, e, f\}$ ,  $C_1 = \{b, c\}$ ,  $C_2 = \{g, h\}$

#### 7.1.2

$$P = \begin{bmatrix} & a & b & c & d & e & f & g & h \\ a & 0 & 0.2 & 0.8 & 0 & 0 & 0 & 0 & 0 \\ b & 0 & 0 & 0 & 0.3 & 0.7 & 0 & 0 & 0 \\ c & 0 & 0 & 0 & 0 & 0.3 & 0.7 & 0 & 0 \\ d & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ e & 0 & 0 & 0 & 0 & 0 & 0 & 0.5 & 0.5 \\ f & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ g & 0.8 & 0 & 0 & 0.2 & 0 & 0 & 0 & 0 \\ h & 0.8 & 0 & 0 & 0 & 0 & 0.2 & 0 & 0 \end{bmatrix}$$

$$P^7 = \begin{bmatrix} 0.00000 & 0.03200 & 0.12800 & 0.00000 & 0.00000 & 0.00000 & 0.21000 & 0.63000 \\ 0.67200 & 0.00000 & 0.00000 & 0.05480 & 0.06080 & 0.21240 & 0.00000 & 0.00000 \\ 0.67200 & 0.00000 & 0.00000 & 0.05080 & 0.06080 & 0.21640 & 0.00000 & 0.00000 \\ 0.00000 & 0.13440 & 0.53760 & 0.00000 & 0.00000 & 0.00000 & 0.08800 & 0.24000 \\ 0.00000 & 0.13440 & 0.53760 & 0.00000 & 0.00000 & 0.00000 & 0.08400 & 0.24400 \\ 0.00000 & 0.13440 & 0.53760 & 0.00000 & 0.00000 & 0.00000 & 0.08000 & 0.24800 \\ 0.26240 & 0.00000 & 0.00000 & 0.05792 & 0.25536 & 0.42432 & 0.00000 & 0.00000 \\ 0.26240 & 0.00000 & 0.00000 & 0.05632 & 0.25536 & 0.42592 & 0.00000 & 0.00000 \end{bmatrix}$$

$$P^{15} = \begin{bmatrix} 0.00000 & 0.06559 & 0.26235 & 0.00000 & 0.00000 & 0.00000 & 0.16802 & 0.50405 \\ 0.53765 & 0.00000 & 0.00000 & 0.05328 & 0.12462 & 0.28445 & 0.00000 & 0.00000 \\ 0.53765 & 0.00000 & 0.00000 & 0.05328 & 0.12462 & 0.28446 & 0.00000 & 0.00000 \\ 0.00000 & 0.10753 & 0.43012 & 0.00000 & 0.00000 & 0.00000 & 0.11560 & 0.34675 \\ 0.00000 & 0.10753 & 0.43012 & 0.00000 & 0.00000 & 0.00000 & 0.11559 & 0.34676 \\ 0.00000 & 0.10753 & 0.43012 & 0.00000 & 0.00000 & 0.00000 & 0.11558 & 0.34676 \\ 0.36988 & 0.00000 & 0.00000 & 0.05538 & 0.20431 & 0.37044 & 0.00000 & 0.00000 \\ 0.36988 & 0.00000 & 0.00000 & 0.05538 & 0.20431 & 0.37044 & 0.00000 & 0.00000 \end{bmatrix}$$

## Problem 7.2

### 7.2.1

Period is  $d = 2$  and the cycles are 2  $C_0 = \{a, c\}, C_1 = \{b\}$

### 7.2.2

$$P = \begin{bmatrix} & a & b & c \\ a & 0 & 1 & 0 \\ b & \alpha & 0 & 1 - \alpha \\ c & 0 & 1 & 0 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} & a & b & c \\ a & 0 & 1 & 0 \\ b & \alpha^2 & 0 & (1 - \alpha)^2 \\ c & 0 & 1 & 0 \end{bmatrix}$$

$$P^{20} = \begin{bmatrix} a & b & c \\ a & 0 & 1 & 0 \\ b & \alpha^{20} & 0 & (1-\alpha)^{20} \\ c & 0 & 1 & 0 \end{bmatrix}$$

$$P^\infty = \begin{bmatrix} a & b & c \\ a & 0 & 1 & 0 \\ b & 0 & 0 & 0 \\ c & 0 & 1 & 0 \end{bmatrix}$$

### 7.2.3

Given the chain the eigenvalues are  $\lambda_1 = 1, \lambda_2 = -1, \lambda_3 = 0$

Therefore for  $\pi_a(n)$  we have:

$$\pi_a(n) = a\lambda_1^n + b\lambda_2^n \quad (1a)$$

$$\pi_a(0) = a + b = (\pi(0)P^0)_a = 1 \quad (1b)$$

$$\pi_a(1) = a - b = (\pi(0)P^1)_a = 0 \quad (1c)$$

$$\pi_a(2) = a + b = (\pi(0)P^2)_a = \alpha \quad (1d)$$

$$(1e)$$

Solving the equation system we have that  $a = \frac{2-\alpha}{2}, b = \frac{\alpha}{2}$ .

Therefore,  $\pi_a(n) = \frac{2-\alpha}{2} + (-1)^n \frac{\alpha}{2}$

Therefore for  $\pi_b(n)$  we have:

$$\pi_b(n) = a\lambda_1^n + b\lambda_2^n \quad (2a)$$

$$\pi_b(0) = a + b = (\pi(0)P^0)_b = 0 \quad (2b)$$

$$\pi_b(1) = a - b = (\pi(0)P^1)_b = 1 \quad (2c)$$

$$\pi_b(2) = a + b = (\pi(0)P^2)_b = 0 \quad (2d)$$

$$(2e)$$

Solving the equation system we have that  $a = \frac{1}{2}, b = -\frac{1}{2}$ .

Therefore,  $\pi_b(n) = \frac{1}{2} + (-1)^n - \frac{1}{2}$

Therefore for  $\pi_c(n)$  we have:

$$\pi_c(n) = a\lambda_1^n + b\lambda_2^n \quad (3a)$$

$$\pi_c(0) = a + b = (\pi(0)P^0)_c = 0 \quad (3b)$$

$$\pi_c(1) = a - b = (\pi(0)P^1)_c = 0 \quad (3c)$$

$$\pi_c(2) = a + b = (\pi(0)P^2)_c = 1 - \alpha \quad (3d)$$

$$(3e)$$

Solving the equation system we have that  $a = \frac{1-\alpha}{2}, b = \frac{1-\alpha}{2}$ .  
Therefore,  $\pi_c(n) = \frac{1-\alpha}{2} + (-1)^n \frac{1-\alpha}{2}$

### Problem 7.3

I havent follow this problem. How to compute the stationary distribution without values? On the other hand i havent followed the problem properly.