

Stochastic Network Modeling (SNM)

Llorenç Cerdà-Alabern

Universitat Politècnica de Catalunya

Departament d'Arquitectura de Computadors

llorenc@ac.upc.edu

Parts

- I Introduction
- II Discrete Time Markov Chains (DTMC)
- III **Continuous Time Markov Chains (CTMC)**
- IV Queuing Theory

Part III

Continuous Time Markov Chains (CTMC)

Outline

- Definition of a CTMC
- Transient Solution
- Embedded MC of a CTMC
- Classification of States
- Steady State
- Semi-Markov Process
- Finite Absorbing Chains

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains



Definition of a CTMC

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transition Matrix

State Transition
Diagram

Sojourn Time

Exponential jumps
Description of a CTMC

Example: Pure Aloha
System

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Finite
Absorbing

Properties of a continuous time MC

- The states must be a numerable set.
- Let $X(t)$ be the event {at time t the system is in state i }, then it must hold the **memoryless property**:

$$P(X(t_n) = i \mid X(t_1) = j, X(t_2) = k, \dots) =$$

$$P(X(t_n) = i \mid X(t_1) = j) \text{ for any } t_n > t_1 > t_2 > t_3 \dots$$



Definition of a CTMC

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transition Matrix

State Transition
Diagram

Sojourn Time

Exponential Jumps
Description of a CTMC

Example: Pure Aloha
System

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Finite
Absorbing

Transition Matrix

- **Transition probabilities:**

$$p_{ij}(t_1, t_2) = P(X(t_2) = j \mid X(t_1) = i)$$

- For an **homogeneous chain**:

$$\begin{aligned} p_{ij}(t) &= P(X(t_1 + t) = j \mid X(t_1) = i) = \\ &= P(X(t) = j \mid X(0) = i), \forall t_1 \end{aligned}$$

- In matrix form (**transition probability matrix**):

$$\mathbf{P}(t) = \begin{bmatrix} p_{11}(t) & p_{12}(t) & \cdots \\ p_{21}(t) & p_{22}(t) & \cdots \\ \cdots & \cdots & \cdots \end{bmatrix}, t \geq 0$$

- Notes:

- Compare with the n-step prob. matrix of a DTMC: $\mathbf{P}(n)$.
- $\mathbf{P}(t)$ must be a **stochastic matrix** (all rows add to 1).

Transition Matrix

$$p_{ij}(t) = P(X(t) = j \mid X(0) = i), t \geq 0$$

- We look for an equivalent 1-step prob. matrix **P** of DTMCs.
- For consistency: $\lim_{t \rightarrow 0} p_{ij}(t) = \delta_{ij}$. In matrix form:

$$\lim_{t \rightarrow 0} \mathbf{P}(t) = \mathbf{I}.$$

- And assume that the following **transition rates** exist:

$$q_{ij} = \lim_{t \rightarrow 0} \frac{p_{ij}(t)}{t}, i \neq j; \quad q_{ii} = \lim_{t \rightarrow 0} \frac{p_{ii}(t) - 1}{t}$$

- In matrix form: $\mathbf{Q} = \lim_{t \rightarrow 0} \frac{\mathbf{P}(t) - \mathbf{I}}{t}$
- Note that $\sum_j p_{ij}(t) = 1 \Rightarrow p_{ii}(t) = 1 - \sum_{j \neq i} p_{ij}(t)$, thus:


$$q_{ii} = \lim_{t \rightarrow 0} \frac{p_{ii}(t) - 1}{t} = \lim_{t \rightarrow 0} \frac{-\sum_{j \neq i} p_{ij}(t)}{t} = -\sum_{j \neq i} q_{ij}$$

Transition Matrix

- The matrix **Q** is called the **transition rate or infinitesimal generator** of the chain.
- Since $q_{ii} = -\sum_{j \neq i} q_{ij}$, **all the rows of Q add to 0**.
- The rate q_{ij} , $i \neq j$ measures “how fast” the chain moves from state i to j : the higher is q_{ij} , the faster it moves from i to j .
- For $q_{ii} = -\sum_{j \neq i} q_{ij}$, the higher $-q_{ii}$ is, the faster the chain leaves state i .
- If $q_{ij} = 0$, $\forall j \Rightarrow q_{ii} = 0$, then i is an **absorbing state**: the chain “moves with rate 0 from i to other states”, i.e. never leaves i .

State Transition Diagram


- A continuous MC is characterized by the transition rate or infinitesimal generator: the Q-matrix.
- The state transition diagram is now represented as:



- Note that now we have **transition rates** ($0 \leq q_{ij} < \infty, i \neq j$) **and not probabilities**.
- The **rates** q_{ii} are not written in the diagram, **no self transitions**.

Sojourn Time

- Sojourn or holding time: Is the RV H_k equal to the sojourn time in state k :



- The Markov property implies that **the sojourn time is exponentially distributed with parameter q_{ii}** :

$$P(H_i \leq x) = 1 - e^{q_{ii}x} \Rightarrow P(H_i > x) = e^{q_{ii}x}, \quad q_{ii} = -\sum_{j \neq i} q_{ij}, \quad x \geq 0$$

The exponential distribution satisfies the Markov property

- Markov property (**memoryless**):

$$P(X(t_2) = i \mid X(t_1) = i, X(0) = i) =$$

$$P(X(t_2) = i \mid X(t_1) = i), t_2 > t_1 > 0$$

- In terms of the sojourn time:

$$P(H_i > t_2 \mid H_i > t_1) = P(H_i > t_2 - t_1)$$


- But:

$$P(H_i > t_2 \mid H_i > t_1) =$$

$$\frac{P(H_i > t_2, H_i > t_1)}{P(H_i > t_1)} = \frac{P(H_i > t_2)}{P(H_i > t_1)} = \frac{e^{q_{ii} t_2}}{e^{q_{ii} t_1}} = e^{q_{ii}(t_2 - t_1)} =$$

$$P(H_i > t_2 - t_1) \quad \square$$


- The **exponential distribution is the only one satisfying the memoryless property.**



Exponential Jumps Description of a CTMC

Assume a process built as follows:

- Upon reaching a state i
 - the process can jump to a state $j \in \{1, 2, \dots, l\}$.
 - A set of **independent exponential RVs**, $\{H_{i1}, H_{i2}, \dots, H_{il}\}$, with parameters $\{q_{i1}, q_{i2}, \dots, q_{il}\}$ are triggered. That is, $P(H_{ij} \leq t) = 1 - e^{-q_{ij}t}$, $t \geq 0$.
- If $\min\{H_{i1}, H_{i2}, \dots, H_{il}\} = H_{ij} \Rightarrow$ the process jumps to the state j . In other words, a transition occurs to state j if the RV H_{ij} is the minimum of $\{H_{i1}, H_{i2}, \dots, H_{il}\}$.



Theorem: This process is a CTMC with transition rates q_{ij} .

Exponential Jumps Description of a CTMC

$$P(H_{ij} \leq t) = 1 - e^{-q_{ij}t}.$$

Theorem: This process is a CTMC with transition rates q_{ij} .

Proof:

- The RV $H_i = \min\{H_{i1}, H_{i2}, \dots, H_{il}\}$ (sojourn time in state i) is **exponentially distributed** with parameter $q_i = \sum_j q_{ij}$:
 $P(H_i \leq t) = 1 - e^{-q_i t}$.
- $P(\min\{H_{i1}, H_{i2}, \dots, H_{il}\} = H_{ij}) = q_{ij} / \sum_j q_{ij}$. Thus, the **transition rate to state j** is:

$$\lim_{t \rightarrow 0} \frac{p_{ij}(t)}{t} = \lim_{t \rightarrow 0} \frac{P(\min\{H_{i1}, H_{i2}, \dots, H_{il}\} = H_{ij}) \times P(H_i \leq t)}{t} =$$

$$\frac{q_{ij}}{\sum_j q_{ij}} \left. \frac{\partial P(H_i \leq t)}{\partial t} \right|_{t=0} = \frac{q_{ij}}{\sum_j q_{ij}} \sum_j q_{ij} = q_{ij} \quad \square$$



Definition of a CTMC

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transition Matrix

State Transition
Diagram

Sojourn Time

Exponential Jumps
Description of a CTMC

Example: Pure Aloha
System

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Finite
Absorbing


Example: Pure Aloha System

- Consider a **Pure Aloha System** with **2 nodes**:
 - Nodes in **thinking state** Tx a packet in a time exponentially distributed with rate λ .
 - Transmission time** is exponentially distributed with rate μ .
 - If two transmissions overlap, the packet is lost and stations become backlogged (after the packet transmission) until the packet is successfully transmitted.
 - Nodes in **backlogged state** Tx a packet in a time exponentially distributed with rate α .

Questions

- Build the state **transition diagram**.

Example: Pure Aloha System



State	Condition	Legend
1	T, T	T Thinking
2	X, T	X Transmitting
3	C, C	C Collided transmission
4	B, C	B Backlogged
5	B, B	
6	X, B	
7	T, B	



Part III

Continuous Time Markov Chains (CTMC)

Outline

- Definition of a CTMC
- Steady State
- **Transient Solution**
- Semi-Markov Process
- Embedded MC of a CTMC
- Finite Absorbing Chains
- Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Chapman-Kolmogorov Equations

- **Chapman-Kolmogorov:** $p_{ij}(t) = \sum_k p_{ik}(t - \alpha) p_{kj}(\alpha), 0 \leq \alpha \leq t$

- Thus:

$$\frac{p_{ij}(t + \Delta t) - p_{ij}(t)}{\Delta t} = \sum_k \left\{ \frac{p_{ik}(t + \Delta t - \alpha) - p_{ik}(t - \alpha)}{\Delta t} p_{kj}(\alpha) \right\}$$

- Taking the limit

$$\alpha \rightarrow t, \Delta t \rightarrow 0 \Rightarrow \begin{cases} p_{ik}(t - \alpha) \rightarrow 0, & i \neq k \\ p_{ik}(t - \alpha) \rightarrow 1, & i = k \end{cases}$$

and using:

we have:

$$\begin{cases} q_{ij} = \lim_{t \rightarrow 0} \frac{p_{ij}(t)}{t}, & i \neq j \\ q_{ij} = \lim_{t \rightarrow 0} \frac{p_{ij}(t) - 1}{t}, & i = j \end{cases}$$

$$\frac{\partial p_{ij}(t)}{\partial t} = \sum_k q_{ik} p_{kj}(t), \quad t \geq 0, \forall i, j$$

Chapman-Kolmogorov Equations (cont)

- we have: $\frac{\partial p_{ij}(t)}{\partial t} = \sum_k q_{ik} p_{kj}(t), t \geq 0, \forall i, j$
- In matrix form: $\frac{\partial \mathbf{P}(t)}{\partial t} = \mathbf{Q} \mathbf{P}(t), t \geq 0$ known as the master equations of a CTMC.
- The solution of the previous matrix differential equation is the **exponential matrix**:

$$\mathbf{P}(t) = e^{\mathbf{Q}t} = \sum_{i=0}^{\infty} \frac{(\mathbf{Q}t)^i}{i!} = \mathbf{I} + \mathbf{Q}t + \frac{\mathbf{Q}^2 t^2}{2!} + \frac{\mathbf{Q}^3 t^3}{3!} + \dots, t \geq 0$$
- Due to rounding errors, the previous series is difficult to compute numerically (the powers of \mathbf{Q} have positive and negative entries).



Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

State Probabilities

- Define the probability of being in state i at time t :

$$\pi_i(t) = P(X(t) = i)$$

- In vector form (row vector)

$$\boldsymbol{\pi}(t) = (\pi_1(t), \pi_2(t), \dots).$$

- Clearly:

$$\pi_i(t) = \sum_k P(X(0) = k) P(X(t) = i \mid X(0) = k) = \sum_k \pi_k(0) p_{ki}(t)$$

- In matrix form:

$$\boldsymbol{\pi}(t) = \boldsymbol{\pi}(0) \mathbf{P}(t) = \boldsymbol{\pi}(0) \mathbf{e}^{\mathbf{Q}t}, t \geq 0$$

where $\boldsymbol{\pi}(0)$ is the **initial distribution**.

- NOTE:** Compare with **DTMC**

$$\boldsymbol{\pi}(n) = \boldsymbol{\pi}(0) \mathbf{P}^n, n \geq 0$$



Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Transient Solution

- If we are interested in the **transient evolution** we shall study $\boldsymbol{\pi}(t) = \boldsymbol{\pi}(0) \mathbf{P}(t) = \boldsymbol{\pi}(0) e^{\mathbf{Q}t}$, $t \geq 0$.
- Assume a **finite CTMC** with N states (infinitesimal generator $\mathbf{Q}^{N \times N}$).
- Assume that \mathbf{Q} can be **diagonalized**: $\mathbf{Q} = \mathbf{L}^{-1} \boldsymbol{\Lambda} \mathbf{L}$, where $\boldsymbol{\Lambda}$ is the diagonal matrix $\boldsymbol{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_N)$, with λ_l , $l = 1, \dots, N$ the eigenvalues of \mathbf{Q} .
- **NOTE**: the **eigenvalues** λ_l of a matrix \mathbf{A} are scalars that satisfy: $\mathbf{l}\mathbf{A} = \lambda_l \mathbf{l}$ (or $\mathbf{A}\mathbf{r} = \lambda_l \mathbf{r}$) for some row vectors \mathbf{l} (column vectors \mathbf{r}), referred to as **left and right eigenvectors**, respectively. Thus, solve the **characteristic polynomial** $\det(\lambda \mathbf{I} - \mathbf{A}) = 0$.

Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Transient Solution

... assume that \mathbf{Q} can be **diagonalized**: $\mathbf{Q} = \mathbf{L}^{-1} \mathbf{\Lambda} \mathbf{L}$

- Since

$$\mathbf{P}(t) = \mathbf{e}^{\mathbf{Q}t} = \sum_{i=0}^{\infty} \frac{(\mathbf{Q}t)^i}{i!} = \sum_{i=0}^{\infty} \frac{(\mathbf{L}^{-1} \mathbf{\Lambda} \mathbf{L}t)^i}{i!} =$$

$$\mathbf{L}^{-1} \text{diag} \left(\sum_{i=0}^{\infty} \frac{(\lambda_1 t)^i}{i!}, \dots \right) \mathbf{L} = \mathbf{L}^{-1} \text{diag}(e^{\lambda_1 t}, \dots) \mathbf{L}$$

we have that

$$\boldsymbol{\pi}(t) = \boldsymbol{\pi}(0) \mathbf{e}^{\mathbf{Q}t} =$$

$$\boldsymbol{\pi}(0) \mathbf{L}^{-1} \text{diag}(e^{\lambda_1 t}, e^{\lambda_2 t}, \dots, e^{\lambda_L t}) \mathbf{L}$$



Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Transient Solution

... we have that $\boldsymbol{\pi}(t) = \boldsymbol{\pi}(0) \mathbf{L}^{-1} \text{diag}(e^{\lambda_1 t}, e^{\lambda_2 t}, \dots, e^{\lambda_L t}) \mathbf{L}$

- Thus, the probability of being in state i is given by:

$$\pi_i(t) = (\boldsymbol{\pi}(t))_i = \sum_{l=1}^N a_i^{(l)} e^{\lambda_l t}, t \geq 0$$

where the **unknown coefficients** $a_i^{(l)}$ can be obtained solving the system of equations:

$$\left. \frac{\partial^n \pi_i(t)}{\partial t^n} \right|_{t=0} = (\boldsymbol{\pi}(0) \mathbf{Q}^n)_i, n = 0, \dots, N-1$$

NOTE: Compare with $(\boldsymbol{\pi}(n))_i = (\boldsymbol{\pi}(0) \mathbf{P}^n)_i, n = 0, \dots, N-1$



Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process


Eigenvalues of an Infinitesimal Generator

- \mathbf{Q} has **an eigenvalue equal to 0** ($\mathbf{Q}\mathbf{x} = \lambda\mathbf{x}$, for $\lambda = 0$, $\mathbf{x} \neq \mathbf{0}$).

Proof: $\mathbf{Q}\mathbf{e} = \mathbf{0}$, where $\mathbf{e} = (1, 1, \dots)^T$ is a column vector of 1 (all rows of \mathbf{Q} add to 0). □

- The eigenvalue $\lambda = 0$ is **single** if \mathbf{Q} is **irreducible** (Perron-Frobenius theorem). \mathbf{Q} is irreducible if all states communicate: for $t > 0$, $p_{ij}(t) > 0$, $\forall i, j$.
- All eigenvalues of \mathbf{Q} are $\lambda_l \leq 0$.

Proof: Using Gerschgorin's theorem and the fact that the rows of \mathbf{Q} add to 0. □



Gerschgorin discs
of a rate matrix



Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Example

- Assume a CTMC with

$$\mathbf{Q} = \begin{bmatrix} -1 & 1 \\ 1/2 & -1/2 \end{bmatrix}$$

- We want the probability of being in state 2 at time t starting from state 1: $\pi_2(t)$ with $\boldsymbol{\pi}(0) = \begin{bmatrix} 1 & 0 \end{bmatrix}$.



Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Solution

- It can be easily found that the **eigenvalues** of \mathbf{Q} are $\lambda_1 = 0$ and $\lambda_2 = -3/2$.

$$\pi_2(t) = ae^{\lambda_1 t} + be^{\lambda_2 t} = a + be^{-(3/2)t}$$

- Imposing the **boundary conditions**:

$$\pi_2(0) = a + b = (\boldsymbol{\pi}(0) \mathbf{Q}^0)_2 = (\boldsymbol{\pi}(0) \mathbf{I})_2 = (\boldsymbol{\pi}(0))_2 = 0$$

$$\left. \frac{\partial \pi_2(t)}{\partial t} \right|_{t=0} = b(-3/2) = (\boldsymbol{\pi}(0) \mathbf{Q})_2 = \mathbf{Q}_{12} = 1$$

we have that $a = 2/3$, $b = -2/3$, thus:

$$\pi_2(t) = 2/3 (1 - e^{-(3/2)t}), \quad t \geq 0$$

Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC

Classification
of States

Steady State

Semi-Markov
Process

Chain with a Defective Matrix

- What if \mathbf{Q} cannot be diagonalized? (**defective** matrix).
- Let λ_l , $l = 1, \dots, L$ be the eigenvalues of $\mathbf{Q}^{N \times N}$, each with multiplicity k_l ($k_l \geq 1$, $\sum_l k_l = N$). Then [1]:

$$\pi_j(t) = \sum_{l=1}^L e^{\lambda_l t} \sum_{m=0}^{k_l-1} a_j^{(l,m)} t^m$$

where $a_j^{(l,m)}$ are constants. So, exponentials associated with eigenvalues λ_l of multiplicity $k_l > 1$ are multiplied by polynomials in t of degree $k_l - 1$.

- [1] Llorenç Cerdà-Alabern. *Transient Solution of Markov Chains Using the Uniformized Vandermonde Method*. Tech. rep. UPC-DAC-RR-XCSD-2010-2. Universitat Politècnica de Catalunya, Dec. 2010. URL: https://www.ac.upc.edu/app/research-reports/html/research_center_index-XCSD-2010,en.html.

Transient Solution

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Chapman-Kolmogorov
Equations

State Probabilities

Transient Solution

Eigenvalues of an
Infinitesimal
Generator

Example

Chain with a Defective
Matrix

Example

Embedded MC
of a CTMC


Classification
of States

Steady State

Semi-Markov
Process

Example

- Assume the CTMC:



$$\pi(0) = [1 \quad 0 \quad 0]$$

$$\mathbf{Q} = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

- We have $\lambda_1 = 0$ and $\lambda_2 = -1$ with multiplicity 2. Thus:

$$\pi_3(t) = a + e^{-t}(b + ct)$$

- We have that $a = 1$, because state 3 is absorbing. Imposing $\pi_3(0) = 0$ and $\pi'_3(0) = 0$, we have $b = c = -1$, and

$$\pi_3(t) = 1 - e^{-t}(1 + t), \quad t \geq 0$$



Part III

Continuous Time Markov Chains (CTMC)

Outline

- Definition of a CTMC
- Transient Solution
- Embedded MC of a CTMC
- Classification of States
- Steady State
- Semi-Markov Process
- Finite Absorbing Chains

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Definition

Proof

Example


Classification
of States

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains

Definition



- We form a discrete time process $X^e(n)$, called the **Embedded MC (EMC)**, by looking at a CTMC at the transition time instants.

Theorem: This process is a **DTMC** with transition probabilities:

$$p_{ij}^e = \begin{cases} 0, & i = j \\ \frac{q_{ij}}{\sum_{j \neq i} q_{ij}}, & i \neq j \end{cases}$$

- NOTE:** If i is **absorbing** ($q_{ii} = 0$), we define $p_{ii}^e = 1$.


Proof

Theorem: This process is a DTMC with transition probabilities:

$$p_{ij}^e = \begin{cases} 0, & i = j \\ \frac{q_{ij}}{\sum_{j \neq i} q_{ij}}, & i \neq j \end{cases}$$

- The EMC satisfies the **memoryless** property.
- Since we look the system only upon transition to a different state: $p_{ii}^e = 0$. NOTE: it might be $p_{ii}^e \neq 0$ if we look at transitions that end up in the same state.
- The probability that there is a transition from state i to j in the CTMC is the probability that the exponentially distributed RV with parameter q_{ij} is the **minimum from the independent exponentially distributed RVs** with parameters $\{q_{ik}\}_{k \neq i}$. This probability is $q_{ij} / \sum_{k \neq i} q_{ik}$. \square

Example



$$\mathbf{Q} = \begin{bmatrix} -7 & 4 & 3 \\ 0 & -2 & 2 \\ 5 & 3 & -8 \end{bmatrix} \Rightarrow \mathbf{P}_e = \begin{bmatrix} 0 & 4/7 & 3/7 \\ 0 & 0 & 1 \\ 5/8 & 3/8 & 0 \end{bmatrix}$$

- Each **transition** in the CTMC is a transition in the EMC.
- One step in ***i*** in the EMC is a **sojourn time H_i** in the CTMC.



Part III

Continuous Time Markov Chains (CTMC)

Outline

- Definition of a CTMC
- Steady State
- Transient Solution
- Semi-Markov Process
- Embedded MC of a CTMC
- Finite Absorbing Chains
- Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Irreducibility
Transient and
Recurrent
Mean recurrence time
of the CTMC

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains



Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Irreducibility

Transient and
Recurrent

Mean recurrence time
of the CTMC

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains

Irreducibility

- A state j is said to **communicate** with i , $i \leftrightarrow j$, if $p_{ij}(t_1) > 0$, $p_{ji}(t_2) > 0$ for some $t_1 \geq 0$, $t_2 \geq 0$.
- We define an **irreducible closed set, ICS** C_k as a set where all states communicate with each other, and have no transitions to other states out of the set:
 $i \leftrightarrow j, \forall i, j \in C_k$ and $q_{ij} = 0, \forall i \in C_k, j \notin C_k$
- An **absorbing state** form an ICS of only one element. This state, i , must have $q_{ij} = 0 \forall i, j$.
- **Transient states** do not belong to any ICS.
- A MC is **irreducible** if all the states form a unique ICS.

Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Irreducibility

Transient and
Recurrent

Mean recurrence time
of the CTMC

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains

Irreducibility

- Assume a MC has **M ICSs**: By properly numbering the states, we can write \mathbf{P} as an M block diagonal matrix with the probabilities of the transient states in the last rows.
- Example**, if $M = 3$:

$$\mathbf{Q} = \begin{array}{|c|c|c|} \hline \mathbf{Q}_1 & & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{Q}_2 & \\ \hline & & \mathbf{Q}_3 \\ \hline \text{at least} & & \\ \text{one} > 0 & & \mathbf{T} \\ \hline \end{array}$$

$$\Rightarrow \boldsymbol{\pi}(t) = \boldsymbol{\pi}(0) e^{\mathbf{Q}t} = \boldsymbol{\pi}(0)$$

$$\begin{array}{|c|c|c|} \hline e^{\mathbf{Q}_1 t} & & \mathbf{0} \\ \hline \mathbf{0} & e^{\mathbf{Q}_2 t} & \\ \hline & & e^{\mathbf{Q}_3 t} \\ \hline \text{at least} & & \\ \text{one} > 0 & & e^{\mathbf{T}t} \\ \hline \end{array}$$

- Note that **the M sub-matrices are infinitesimal generators** (their rows add to 0).

Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Irreducibility

Transient and
Recurrent

Mean recurrence time
of the CTMC


Steady State

Semi-Markov
Process

Finite
Absorbing
Chains

Transient and Recurrent

- **Recurrent:** States that, being visited, have a probability > 0 of being visited again. They are visited an infinite number of times when $t \rightarrow \infty$.
- **Transient:** States that, being visited, have a probability > 0 of never being visited again. They are visited a finite number of times when $t \rightarrow \infty$.
- **Absorbing:** A single (recurrent) state where the chain remains with probability $= 1$.



State 1 is **transient**
States 2 and 3 are **recurrent**
State 4 is **absorbing**



Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Irreducibility

Transient and
Recurrent

Mean recurrence time
of the CTMC

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains

Transient and Recurrent

- To derive a classification criteria, we shall study the **embedded MC (EMC)**, and proceed as in DTMC: Let $f_{ij}^e(n)$ the first passage prob. of the EMC, and $f_{ij}^e = \sum_{n=1}^{\infty} f_{ij}^e(n)$.
- If $f_{ii}^e = 1$ we say i is a **recurrent state**.
- If $f_{ii}^e < 1$ we say i is a **transient state**.
- When $f_{ii}^e = 1$, we define the **mean recurrence time of the EMC** $m_{ii}^e = \sum_{n=1}^{\infty} n f_{ii}^e(n)$. **NOTE:** in **steps**, not time units.
- If $m_{ii}^e = \infty$ the state is **null recurrent**.
- If $m_{ii}^e < \infty$ the state is **positive recurrent**.
- **NOTES:** (i) Even if the EMC is periodic, **there are not periodic CTMC** (it has no sense). (ii) f_{ij}^e and m_{ij}^e can be computed using the **recursive equations**.

Classification of States

Continuous
Time Markov
Chains (CTMC)

Definition of a
CTMC

Transient
Solution

Embedded MC
of a CTMC

Classification
of States

Irreducibility

Transient and
Recurrent

Mean recurrence time
of the CTMC

Steady State

Semi-Markov
Process

Finite
Absorbing
Chains

Mean recurrence time of the CTMC

- If the chain is in i at a time t , it takes an **expected time to leave i** equal to $1/(-q_{ii}) = 1/\sum_{j \neq i} q_{ij}$ (**sojourn time exponentially distributed** with rate $q_i = -q_{ii} = \sum_{j \neq i} q_{ij}$).
- Thus, if the chain is **in state i** , it takes a **mean time to enter state j** (**mean first passage time**):

$$m_{ij} = \frac{1}{q_i} + \sum_{k \neq i} p_{ik}^e m_{kj}$$

- Since: $p_{ij}^e = \begin{cases} 0, & i = j \\ \frac{q_{ij}}{\sum_{j \neq i} q_{ij}} = \frac{q_{ij}}{q_i}, & i \neq j \end{cases}$ we have:

$$m_{ij} = \frac{1}{q_i} + \sum_{k \neq i} p_{ik}^e m_{kj} = \frac{1}{q_i} + \sum_{k \neq i} \frac{q_{ik}}{q_i} m_{kj} \text{ [time units]}$$