


Crossing the x -axis

MOPSS



Suggested readings

- Evan Chen's advice [On reading solutions](https://blog.evanchen.cc/2017/03/06/on-reading-solutions/), available at <https://blog.evanchen.cc/2017/03/06/on-reading-solutions/>.
- Evan Chen's [Advice for writing proofs/Remarks on English](https://web.evanchen.cc/handouts/english/english.pdf), available at <https://web.evanchen.cc/handouts/english/english.pdf>.
- [Notes on proofs](#) by Evan Chen from [OTIS Excerpts](#) [[Che25](#), Chapter 1].
- [Tips for writing up solutions](https://www.math.utoronto.ca/barbeau/writingup.pdf) by Edward Barbeau, available at <https://www.math.utoronto.ca/barbeau/writingup.pdf>.
- Evan Chen discusses why [math olympiads](#) are a valuable experience for [high schoolers](#) in the post on [Lessons from math olympiads](#), available at <https://blog.evanchen.cc/2018/01/05/lessons-from-math-olympiads/>.

List of problems and examples

1.1	Example	2
1.2	Exercise (All-Russian Mathematical Olympiad 2013 Grade 9 Day 1 P1, AoPS)	2
1.3	Exercise (All-Russian Mathematical Olympiad 2007 Grade 8 Day 1 P1, AoPS, by O. Podlipsky)	4
1.4	Example	5
1.5	Example	5
1.6	Example	5
1.7	Exercise (China TST 1995 Day 2 P2, AoPS)	6

§1 Crossing the x -axis

Here are a few problems from [this notes](#), and [this one](#).

Example 1.1. Suppose $P(x)$ is a polynomial with real coefficients such that $P(x) = x$ has no real solution. Show that $P(P(x)) = x$ has no real solutions.

Walkthrough —
(a)

Solution 1. Since $x \mapsto P(x)$ defines a continuous map from $\mathbb{R} \rightarrow \mathbb{R}$, by the intermediate value theorem, it follows that $P(x) > x$ holds for all $x \in \mathbb{R}$ or $P(x) < x$ holds for all $x \in \mathbb{R}$. If $P(x) > x$ holds for all $x \in \mathbb{R}$, then $P(P(x)) > P(x) > x$ holds for all $x \in \mathbb{R}$, and hence $P(P(x)) = x$ has no real solutions. Similarly, if $P(x) < x$ holds for all $x \in \mathbb{R}$, then $P(P(x)) = x$ has no real solutions. ■

Exercise 1.2 (All-Russian Mathematical Olympiad 2013 Grade 9 Day 1 P1, AoPS). Given three distinct real numbers a , b , and c , show that at least two of the three following equations

$$\begin{aligned}(x - a)(x - b) &= x - c, \\(x - b)(x - c) &= x - a, \\(x - c)(x - a) &= x - b\end{aligned}$$

have real solutions.

Walkthrough —
(a)

Solution 2. Without loss of generality, let us consider the case that the equations

$$(x - a)(x - b) = x - c, (x - b)(x - c) = x - a$$

do not have any real root. It follows that their discriminants are negative, that is,

$$(a + b + 1)^2 - 4(ab + c), (b + c + 1)^2 - 4(bc + a)$$

are negative. Note that

$$\begin{aligned} (a + b + 1)^2 - 4(ab + c) &= (a - b)^2 + 1 + 2(a + b - 2c) \\ &= (b - a + 1)^2 + 4(a - c), \\ (b + c + 1)^2 - 4(bc + a) &= (b - c)^2 + 1 + 2(b + c - 2a) \\ &= (b - c + 1)^2 + 4(c - a). \end{aligned}$$

Since a and c are distinct real numbers, one of $a - c$ and $c - a$ is positive. Thus, the discriminant of one of the equations

$$(x - a)(x - b) = x - c, (x - b)(x - c) = x - a$$

is non-negative, which is a contradiction. Therefore, at least two of the three equations have real solutions. ■

Solution 3. Let $f(x)$, $g(x)$, and $h(x)$ denote the polynomials defined by

$$\begin{aligned} f(x) &= (x - a)(x - b) - (x - c), \\ g(x) &= (x - b)(x - c) - (x - a), \\ h(x) &= (x - c)(x - a) - (x - b). \end{aligned}$$

Note that

$$\begin{aligned} f(x) + g(x) &= (x - b)(2x - a - c) - (2x - a - c) \\ &= (2x - a - c)(x - b - 1). \end{aligned}$$

This implies that $f(x) + g(x)$ admits a real root. Hence, at least one of $f(x)$ and $g(x)$ is non-positive at some real number. If one of them admits a real root, then we are done. Otherwise, at least one of them is negative at some real number. Also note that if k is a real number satisfying $k \geq |a| + |b| + |c| + 2$, then

$$\begin{aligned} f(k + |a| + |b|) &= (k + |a| + |b| + a)(k + |a| + |b| + b) - (k + |a| + |b| + c) \\ &\geq k^2 - k - |a| - |b| - |c| \\ &\geq k - |a| - |b| - |c| \\ &> 0 \end{aligned}$$

holds, and similarly,

$$g(k + |a| + |b|) > 0$$

hold too. By the intermediate value theorem, at least one of $f(x)$ and $g(x)$ admits a real root. Interchanging the roles of f , g , and h , it follows that at least one of any two of the polynomials f , g , and h admits a real root. Therefore, at least two of the three equations have real solutions for x . ■

Exercise 1.3 (All-Russian Mathematical Olympiad 2007 Grade 8 Day 1 P1, AoPS, by O. Podlipsky). Given reals numbers a , b , c , prove that at least one of three equations

$$\begin{aligned}x^2 + (a - b)x + (b - c) &= 0, \\x^2 + (b - c)x + (c - a) &= 0, \\x^2 + (c - a)x + (a - b) &= 0\end{aligned}$$

has a real root.

Walkthrough —

(a)

Solution 4. Note that the sum of the discriminants of the quadratic polynomials

$$x^2 + (a - b)x + (b - c), x^2 + (b - c)x + (c - a), x^2 + (c - a)x + (a - b)$$

are equal to

$$\begin{aligned}(a - b)^2 - 4(b - c) + (b - c)^2 - 4(c - a) + (c - a)^2 - 4(a - b) \\= (a - b)^2 + (b - c)^2 + (c - a)^2,\end{aligned}$$

which is non-negative, and hence, at least one of the discriminants is non-negative. This implies that at least one of the three equations has a real root. ■

Solution 5. Consider the three quadratic polynomials defined by

$$\begin{aligned}f(x) &= x^2 + (a - b)x + (b - c), \\g(x) &= x^2 + (b - c)x + (c - a), \\h(x) &= x^2 + (c - a)x + (a - b).\end{aligned}$$

Note that

$$f(0) + g(0) + h(0) = 0,$$

which implies that at least one of $f(0)$, $g(0)$, $h(0)$ is non-positive. Without loss of generality, assume that $f(0) \leq 0$. If $f(0)$ is zero, then $x = 0$ is a root of $f(x)$. If $f(0) < 0$, then noting that

$$\begin{aligned}
 & f(2|a - b| + 2|b - c| + 1) \\
 &= \frac{1}{2}(2|a - b| + 2|b - c| + 1)^2 + (a - b)(2|a - b| + 2|b - c| + 1) \\
 &\quad + \frac{1}{2}(2|a - b| + 2|b - c| + 1)^2 + (b - c) \\
 &= \frac{1}{2}(2|a - b| + 2|b - c| + 1)(2|a - b| + 2|b - c| + 1 - 2(a - b)) \\
 &\quad + \frac{1}{2}(2|a - b| + 2|b - c| + 1)^2 + (b - c) \\
 &\geq 1
 \end{aligned}$$

holds, it follows by the intermediate value theorem that $f(x)$ has a real root. ■

Example 1.4. Any polynomial of odd degree with real coefficients has a real root.

Example 1.5. Let $P(x)$ and $Q(x)$ be monic polynomials of degree 10 having real coefficients. Assume that the equation $P(x) = Q(x)$ has no real roots. Prove that the equation $P(x + 1) = Q(x - 1)$ has at least one real root.

Walkthrough —

- (a) Consider the difference $P(x) - Q(x)$ to show that the coefficient of x^9 in these polynomials are equal.
- (b) Prove that the polynomial $P(x + 1) - Q(x - 1)$ is of degree 9.

Solution 6. Note that $P(x) - Q(x)$ is a polynomial of degree at most 9 having real coefficients. Since $P(x) - Q(x)$ has no real root, it follows that it has degree at most 8. In other words, the coefficients of x^9 in $P(x), Q(x)$ are the same. Note that $P(x + 1) - Q(x - 1)$ is of degree ≤ 9 , and the coefficient of x^9 in $P(x + 1) - Q(x - 1)$ is equal to the coefficient of x^9 in $(x + 1)^{10} - (x - 1)^{10}$, which is equal to 20. This shows that $P(x + 1) - Q(x - 1)$ is a polynomial of degree 9 with real coefficients. Consequently, it has at least one real root. ■

Example 1.6. Let $P(x)$ be a nonconstant polynomial with real coefficients having a real root. Suppose it does not vanish at 0. Show that the monomial terms appearing in $P(x)$ can be erased one by one to obtain its constant term such that the intermediate polynomial have at least one real root.

Walkthrough —

(a)

Solution 7. Write $P(x) = a_n x^n + \cdots + a_0$ with $a_n, \dots, a_0$ lying in $\mathbb{R}$, and $a_n a_0 \neq 0$. Since any polynomial of odd degree has a real root, it follows that if the degree of $P(x)$ is odd, then the nonconstant monomials, other than the leading term, can be erased one by one, and then, the leading term can be erased to obtain a_0 , and the intermediate polynomials have a real root. If a_n, a_0 are of opposite signs and n is even, then the same process can be followed, and note that any of the intermediate polynomials takes values of opposite signs at 0 and at a large enough integer, and hence has a real root.

Let us assume that n is even, and a_n, a_0 are of the same sign. Let α denote a real root of $P(x)$. Let $Q(x)$ denote the polynomial $P(x) - a_n x^n$. Note that $Q(0) = a_0$ and $Q(\alpha) = -a_n \alpha^n$. Since a_0 is nonzero, it follows that $\alpha \neq 0$, and hence, $Q(0), Q(\alpha)$ are of opposite signs. This shows that $Q(x)$ is a nonconstant polynomial with real coefficients having a real root, and it does not vanish at 0. Since the degree of $Q(x)$ is smaller than that of $P(x)$, by induction, we are done. ■

Exercise 1.7 (China TST 1995 Day 2 P2, AoPS). Alice and Bob play a game with a polynomial of degree at least 4:

$$x^{2n} + \square x^{2n-1} + \square x^{2n-2} + \cdots + \square x + 1.$$

They take turns to fill the empty boxes. If the resulting polynomial has no real root, Alice wins, otherwise, Bob wins. If Alice goes first, who has a winning strategy?

Walkthrough —

- (a) There are more odds than evens among the integers $1, 2, \dots, 2n-1$.
- (b) Can Bob have a winning strategy using the odds in his favour?

Solution 8. Bob has a winning strategy, as described below.

Bob makes sure that at the end of each of his turns except the last one, the number of even powers of x whose coefficients have been provided by some of them is equal to the number of odd powers of x whose coefficients have been provided by some of them. This can be done, for instance, if during a turn of Bob, other than the last turn, Bob provides the coefficient of an odd (resp. even) power of x if Alice has provided the coefficient of an even (resp. odd) power of x in the preceeding turn.

Since $n \geq 2$, it follows that Bob gets at least one turn. At the beginning of the final turn of Bob, there are two powers of x whose coefficients are to be determined, denote them by x^i, x^j , their coefficients by c_i, c_j respectively. Let

$Q(x)$ denote the polynomial, obtained by the erasing the terms corresponding to x^i, x^j from the polynomial that Bob had at the beginning of his final turn. Note that

$$P(x) = Q(x) + c_i x^i + c_j x^j.$$

Note that at least of i, j is odd. Interchanging i, j if necessary, let us assume that i is odd. We describe the strategy that Bob follows in the two cases below.

Let us consider the case that j is even. Bob determines c_j in such a way that for any choice of c_i , the completed polynomial $P(x)$ is guaranteed to have at least one real root. This can be done, for instance, by taking c_j satisfying

$$Q(1) + Q(-1) + 2c_j = 0.$$

For any choice of c_i , the above choice of c_j shows that $P(1) + P(-1) = 0$, which implies that $P(x)$ has a root in the interval $[-1, 1]$.

Let us consider the case that j is odd. Bob determines c_j in such a way that for any choice of c_i by Alice in the next turn, the completed polynomial $P(x)$ is guaranteed to have at least one real root. This can be done, for instance, by taking c_j satisfying

$$Q(2) + c_j 2^j + 2^i Q(-1) - c_j 2^i = 0.$$

Since $i \neq j$, the above holds for some $c_j \in \mathbb{R}$. For any choice of c_i , note that

$$P(2) + 2^i P(-1) = 0$$

holds, which implies that $P(x)$ has a root in $[-1, 2]$. ■

References

- [Che25] EVAN CHEN. *The OTIS Excerpts*. Available at <https://web.evanchen.cc/excerpts.html>. 2025, pp. vi+289 (cited p. 1)