

Symmetric groups

Chapter 1. Symmetric groups

§1.1 eqn test

$$a = b. \tag{1}$$

Study the equation in [Equation 1](#).

$$c = d. \tag{2}$$

Study the equation in [Equation 2](#).

§1.2 Core Definitions

In this session, we analyze how linear transformations scale specific vector subspaces.

§1.2.1 some defn

Definition 2.1. Let $T : V \rightarrow V$ be a linear operator on a vector space V over a field F . A non-zero vector $v \in V$ is called an **eigenvector** of T if there exists a scalar $\lambda \in F$ such that:

$$T(v) = \lambda v$$

The scalar λ is called the **eigenvalue** corresponding to the eigenvector v .

See [Definition 2.0](#) for the formal definition of eigenvalues and eigenvectors.

§1.3 Fundamental Theorems

Theorem 2.2. Let $T : V \rightarrow V$ be a linear operator. If $v_1, v_2, \dots, v_m$ are eigenvectors corresponding to **distinct** eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$, then the set $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

By [Theorem 2.1](#), we conclude that eigenvectors corresponding to distinct eigenvalues are linearly independent.

§1.4 Homework Practice

Exercise 2.3. Consider the 2×2 matrix A representing a linear transformation on $\mathbb{R}^2$:

$$A = \begin{pmatrix} 4 & 2 \\ 1 & 3 \end{pmatrix}$$

1. Compute the characteristic polynomial $p(\lambda) = \det(A - \lambda I)$.
2. Find the distinct eigenvalues λ_1 and λ_2 by solving:

$$\lambda^2 - 7\lambda + 10 = 0$$

By [Exercise 2.2](#).

Lemma 2.4. lemma

by [Lemma 2.3](#).