Salting the Earth: Shrimp Farming and Agricultural Externalities in Bangladesh

Krittika Ray* Jack Thiemel (Job Market Paper) †

6 November 2026

This paper is updated regularly; please find the latest version here.

Abstract

Climate change is raising groundwater salinity across coastal regions globally, threatening smallholder livelihoods. In southern Bangladesh, a common adaptation is to convert freshwater rice paddies into saltwater shrimp ponds. However, this adaptation comes with an externality: saltwater from shrimp ponds may seep into neighboring fields, damaging rice. This paper considers whether such private adaptation generates negative externalities on neighboring rice farmers and whether those externalities accelerate further adoption. Combining a novel household survey with a satellite panel of 73,300 plots over nine years, we document that a neighbor's conversion reduces vegetation health on adjacent rice plots and raises own conversion probability by 18.0 percentage points. Interpreted through a model of land use based on comparative advantage and a boundary externality, these results imply an adaptation cascade wherein 51.5 percent of ponds exist only because a neighbor converted, and a tax on conversion would raise aggregate farm profit by 11 percent. Thus, adaptation that benefits individual farmers can spread through costs imposed on their neighbors: private decisions that promote climate resilience for some can erode it for others.

We are grateful to Oriana Bandiera, Robin Burgess, Gharad Bryan, Alison Andrew, Clare Balboni, Stefano Caria, Dennis Egger, Rohini Kamal, Niclas Moneke, Michelle Rao, and Chris Woodruff for comments and suggestions. We also thank seminar participants at BRAC University, the 2026 Interdisciplinary PhD Workshop on Sustainable Development, the UCL/IFS/LSE Development Economics Work-in-Progress Seminar, OxDev 2026, and the Heartland Environmental and Resource Economics Workshop at Illinois 2026 for helpful comments. We acknowledge funding from the International Growth Centre (IGC), the Hub for Equal Representation (HER), the LSE Global School of Sustainability (GSoS), the George Webb Medley Fund, and the Suntory and Toyota International Centres for Economics and Related Disciplines (STICERD). This project received approval from the LSE Research Ethics Review Board #350691. Any errors are our own.

*Department of Economics, University of Oxford. krittika.ray@economics.ox.ac.uk

†Department of Economics, London School of Economics. j.p.thiemel@lse.ac.uk

1 Introduction

It is well established that human activity is altering the global climate (IPCC, 2021). A large body of evidence shows this is harming livelihoods, especially in the world's poorest countries (Dell et al., 2012; Burke et al., 2015; Carleton et al., 2022). The extent to which these changes affect human populations depends, in large part, on people's ability to adapt to new conditions (IPCC, 2022).

One setting where adaptation may be slow is agriculture. The cost of new technology, barriers to learning, and heterogeneous returns are just a few well-documented obstacles to take-up.[1] Since agricultural workers are especially exposed to the climate through both physical exposure and the relationship between heat and yields, slow adoption may be especially costly in this sector. We examine a countervailing force: a production externality imposed on farmers who do not adopt. If adopting a new technology harms non-users who can avoid that damage by adopting it themselves, take-up may be faster than in the absence of the externality.

We study this dynamic in the context of southwestern Bangladesh, where farmers can respond to climate-change-induced groundwater salinity by switching from rice cultivation (the old technology) to inland shrimp farming (the new technology) in saltwater ponds. During the dry season, farmers can flood their plots with salty water from groundwater, canals or slow-flowing rivers, which protects them from the damage that saltwater intrusion would otherwise have done to their rice. fields. Neighbors can therefore face a choice between continuing to cultivate rice under worsening conditions and converting to shrimp themselves.

The choice facing an individual farmer may therefore depend not only on the salinity of their own plot, but also on the land-use decisions of adjoining farmers. A plot that would otherwise remain profitable in rice may become less attractive for rice after a neighbor converts. We first ask whether this exposure affects subsequent adoption. We then ask a separate question: whether the induced conversions improve or reduce aggregate farm profits.

This distinction is central to the paper. A farmer who converts because shrimp has become the more profitable use of their own land is adapting to an environmental constraint. A farmer who converts because a neighbor's pond has reduced the return to rice faces a different incentive: one household's response alters the options available to another. The first process may lead to a change in the composition of agriculture without any inefficiency; the second creates the possibility that privately rational choices reinforce one another even when the resulting allocation is not best

[1] For surveys of this literature, see Magruder (2018); Suri and Udry (2022); Adamopoulos et al. (2026).

for farmers collectively. Our empirical analysis examines whether such dependence on neighboring choices exists, while the model is needed to quantify its extent and assess its welfare implications.

First, using a novel survey that combines a household questionnaire (N = 739) with ground-truth salinity readings (N = 1,478), we document three stylized facts that support the paper's argument that the salinity externality exists and affects patterns of adoption. *Fact 1:* We begin by showing that farmers believe shrimp farming generates a production externality. In a hypothetical plot exercise, farmers expect that, in the year following their neighbors' conversion from rice to shrimp, their own plots will experience both higher salinity and lower yields equivalent to a 48 percent decrease in output if they do not adopt shrimp farming as well. This stated loss is much larger than estimates of the average annual decline in rice yields from long-run salinity trends alone, suggesting that farmers perceive neighboring conversions as a more immediate threat (Dasgupta et al., 2015). *Fact 2:* Farmers' beliefs about salinity are supported by our ground-truth salinity survey. Measured salinity is also higher near shrimp ponds. Rice plots bordering shrimp ponds have salinity readings 20.6 percent (0.6 decisiemens per meter) higher than those of plots that do not border shrimp ponds. We take readings in May, immediately before the onset of the annual rainy period, and show that shrimp-adjacent plots have higher readings than their non-shrimp-adjacent counterparts. *Fact 3:* Finally, these saltier rice plots that border shrimp ponds produce lower rice yields (171 kg/acre less) than plots that do not border shrimp farms. While this is not causal evidence of an externality, it is notable that even in equilibrium, when farmers are responding to their neighbors, they are unable to offset these losses and equalize yields across adjacent plots, even if they are able to choose salt-tolerant rice varieties. Taken together, these facts suggest that there is scope for farmers to make land-use decisions with this production externality in mind.

Second, to study whether farmers' beliefs about this externality affect patterns of adoption, we construct a new plot-level panel of land use covering 73,300 plots in coastal Bangladesh from 2017 to 2025. We apply Delineate Anything, a deep-learning model for field-boundary delineation, to high-resolution PlanetScope imagery, allowing us to recover individual plot polygons and the adjacency network through which spillovers may propagate. We then classify each plot during *Rabi* (the dry season, which lasts from November to February) as agriculture or aquaculture using a water index derived from repeated Sentinel-2 observations and a data-driven classification threshold.[2] To our knowledge, this is the first plot-level, seasonal record of shrimp

[2]We apply the Modified Normalized Difference Water Index (MNDWI) of Xu (2006). This index

aquaculture adoption at this scale.

Third, we use the resulting panel to estimate the effects of a neighbor's conversion to aquaculture in an event-study framework, using the estimator of Sun and Abraham (2021) to account for staggered conversion timing and heterogeneous treatment effects across cohorts. A key identification challenge is spatially correlated shocks: both plots may be responding to the same rising underlying salinity. To address the risk of correlated shocks, we tile the study region with hexagons that average 20 plots per hexagon and control for hexagon-by-time fixed effects. Using this approach, we show that the conversion of a plot to aquaculture induces an 18.0 percentage point increase in the likelihood that a bordering plot adopts aquaculture as well. This effect persists for at least five years until the end of the sample. Using the same empirical strategy, we further show that among plots that remain in agriculture, a satellite-derived measure of crop health falls by 1 percent, or 3 percent of the baseline mean, in the year following a neighbor's conversion, before returning to pre-conversion levels in subsequent years. Because this estimate is obtained only from plots that themselves never convert to aquaculture, it likely understates the average penalty: if the most severely affected plots are the most likely to exit agriculture, the remaining sample is positively selected. Together, these results are consistent with an environmental externality that not only damages neighboring agriculture but also reshapes land-use decisions, accelerating the spread of aquaculture.

Fourth, to interpret these effects, we develop a model of land-use choice under a production externality. We build on the Roy (1951) model of selection on comparative advantage in which profits from either shrimp or rice depend on observable plot characteristics and a Fréchet-distributed productivity for every crop à la Costinot et al. (2016); Sotelo (2020) and Farrokhi and Pellegrina (2023). However, in these models, a parcel's returns do not depend on neighbors' crop choices. We add an environmental externality that acts only at the boundary between two adjoining plots to mirror salinity spillovers through groundwater. This lowers the threshold at which a plot switches to aquaculture, creating the potential for a spatial cascade. We model the cascade as a static repeated problem because the empirical analysis gives no evidence of anticipation and land-use decisions are year-over-year reversible; thus, there are no option value gains to waiting to adopt. Last, because beliefs about damages are two orders of magnitude larger than what climate change alone would imply, we do not explicitly model a global salinity trend. The event study estimate of the share of plots that adopt shrimp farming due to neighbors' behavior identifies

better separates water from built-up land and is more reliable in detecting shallow water compared to the unmodified NDWI.

the size of the net externality favoring shrimp farming under the assumption that profits observed in the household survey pin down the dispersion of plots' productivity draws. We estimate the model from the cross-section of rice and shrimp farms in the household survey and the conversion event study from the satellite panel. The dispersion of realized output identifies the shape of the distribution of comparative advantage, while plots' land-use choices identify how size and distance to the river shift relative suitability, with the baseline rate of aquaculture anchored to the satellite data. Equipped with the estimated model, we can compare the take-up of shrimp among plots where we set the externality to zero. On a stylized landscape, we find that the externality generates a substantial cascade: 51.5 percent of the plots that take up aquaculture do so only because of the externality from their neighbors.

The model and event studies together only allow us to speak to whether or not there is a cascade. To assess whether the observed extent of shrimp farming represents over- or under-adaptation, we decompose the net spillover estimated from land-use choices into damage to neighboring rice and benefits to neighboring ponds, using farmers' elicited beliefs about the rice yield loss a neighbor's conversion would cause, and compare the resulting allocation with the planner's allocation. The result of this exercise is that the recently announced subsidy regime for shrimp in Bangladesh (8% for exports) is in the wrong direction—our model implies that the level of shrimp farming adoption in southern Bangladesh is too high and that a tax on the conversion of agricultural land to aquaculture would increase welfare by 11 percent.

Contributions: A large literature studies the drivers of technology adoption in agriculture. Economists have examined this question since at least Griliches (1957), who showed that the diffusion of hybrid corn tracked its profitability, and subsequent work has identified many reasons adoption may be slow even when new technologies are profitable on average. Farmers must learn about a technology's returns, either through their own experience or from their neighbors (Foster and Rosenzweig, 1995; Conley and Udry, 2010); returns are heterogeneous, so low adoption may partly reflect low returns for some farmers (Suri, 2011); and farmers may fail to notice relevant dimensions of a technology or learn from misspecified models (Hanna et al., 2014). In developing countries, a further set of frictions can slow adoption: missing credit and insurance markets (Karlan et al., 2014; Mobarak and Rosenzweig, 2013; Casaburi and Willis, 2018), exposure to downside risk (Emerick et al., 2016), factor market failures (Jones et al., 2022), behavioral biases (Duflo et al., 2011), and limited access to information (Beaman et al., 2021; Aker and Jack, 2025). We study a setting in which an externality imposed on non-adopters instead accelerates adoption. We show that plots only respond to other plots with which they share a direct border, farmers

also report that they discuss farming with their direct neighbors no more often than with other households in the village, and rice greenness declines on neighboring plots that never convert. Together, these patterns suggest the results are not driven by learning alone. In this setting, the new technology (i.e., shrimp farming) acts as a force that speeds adoption, even though these other frictions would, on net, lead us to expect slow uptake.

Second, we contribute to the literature on adaptation to climate change in low- and middle-income countries.[3] This literature has largely focused on whether and how households adapt and on the constraints that limit adaptation. We highlight a different dimension of heterogeneity in the consequences of adaptation: adaptive behavior by some households can make others worse off. In our setting, conversion to aquaculture accelerates further adoption, while also imposing costs on those who do not convert. In this respect, our paper relates to work showing that adaptation can generate costs for others, such as groundwater depletion from competitive pumping (Ryan and Sudarshan, 2022; Blakeslee et al., 2020) and conflict arising from pastoralist migration in response to drought (McGuirk and Nunn, 2025). Relative to this work, we observe the externality directly at the level of individual plots and their neighbors, allowing us to trace how damage to one plot affects its owner's subsequent decisions and how adoption spreads across a landscape. We show that in responding to climate shocks, people can exacerbate the very shocks for others that they themselves sought to avoid.

Third, within the climate adaptation literature, we contribute to a growing body of work on responses to salinity. Rising salinity in coastal Bangladesh has been linked to migration (Chen and Mueller, 2018), economic precarity (Chen et al., 2022), early-life health (Guimbeau et al., 2024), and lower returns to high-yielding rice varieties (Dasgupta et al., 2015), while recent work studies the adoption of salt-tolerant rice (Patel, 2025). Closest to our work, Haque (2025) shows that fragmented land markets and religious homogeneity can impede the adoption of shrimp aquaculture, but that land may be reallocated toward shrimp through the rental market. A related study documents higher salinity and lower paddy profits on plots closer to shrimp ponds (Morshed et al., 2020). We complement this work with a plot-level seasonal panel tracking land use across 73,300 plots over nine years, which allows us to study adoption dynamically rather than at a single point in time. By observing plot boundaries and the adjacency network between plots, we show that the externality from conversion can generate a cascade in which each adoption raises the likelihood of further conversions nearby.

[3] See Kala et al. (2026) and Carleton et al. (2024) for reviews.

The remainder of the paper proceeds as follows. Section 2 introduces the setting. Section 3 discusses the data. Section 4 introduces the model. Section 5 presents the stylized facts. Section 6 presents the empirical strategy and event-study results. Section 7 discusses estimation of the model and the counterfactuals. Finally, Section 8 concludes.

2 Context/Setting

This section provides background information on the key features of southwestern Bangladesh where this study takes place. It discusses the salinity conditions of the setting and the extent to which saltwater shrimp farming can be accurately described as climate adaptation.

2.1 Climate Change in Bangladesh

This project takes place in the Khulna and Satkhira districts of southwestern Bangladesh. Appendix Figure B1 shows the locations of unions (the smallest administrative unit of Bangladesh) in which we collect survey data; unions used in the satellite analysis are shown in pink. Both districts are bounded by mangrove forests and the saltwater Bay of Bengal to the south, and inland Bangladesh to the north. These districts sit within the Ganges-Brahmaputra-Meghna delta and are low-lying and densely populated. A large share of households work in agriculture, and rice dominates (Bangladesh Bureau of Statistics, 2022). In addition to other climate-induced shocks such as heatwaves, droughts, and flooding that affect the entire country, people living in these districts face disparate additional threats such as storm surges and saltwater intrusion (World Bank Group, 2024). Among these threats, salinity is notable in that it builds gradually and directly reduces agricultural productivity.

Two different processes may govern the evolution of soil salinity, operating on two different time scales. The first is a slow regional change driven by climate and upstream water use. Sea-level rise and storm surges push saline water further inland, while reduced freshwater flows in the dry season allow tidal salinity to travel further up coastal rivers and permeate groundwater (Dasgupta et al., 2015). According to the Bangladesh Ministry of Environment, Forest, and Climate Change, the area under critical saltwater intrusion of 5–10 parts per thousand is expected to increase by 68% by 2050 (Ministry of Environment, Forest and Climate Change, Government of the People's Republic of Bangladesh, 2018). This is a long-run, ongoing process: the Soil Resource Development Institute in the Bangladesh Ministry of Agriculture has found that from 1973 to 2009, salt-affected areas increased by 26.7% (Soil Resource

Development Institute, 2010). The second process is both faster and more localized. When a plot is converted to aquaculture, saline water is brought onto the plot and can spread to neighboring fields within a season, as we show in Section 6. This process is shaped, in part by distance to the coast and by infrastructure: the system of polders and embankments built from the 1960s was designed to keep saline water out of agricultural land, but shrimp farmers have been documented breaching embankments to bring saline water inland (Brammer, 1990). The regional trend thus gradually changes which plots are close to the margin between rice and aquaculture, while the local externality determines how conversion spreads among neighbors within much shorter periods.

Policies that may limit salinity exist, but often require coordinated management that has been difficult to achieve in coastal Bangladesh. A key tool to address groundwater intrusion is the enforcement of limits on over-pumping informed by regular water balance assessments (Stein et al., 2023). Other approaches that facilitate recharge of groundwater include shifting demand to other water sources, reducing water consumption, switching to salt-tolerant crops, or otherwise changing production processes. Because groundwater is a common-pool resource, regulation requires central authority; however, de facto water governance suffers from weak coordination among agencies and ineffective local water management institutions (Esha et al., 2026). Even if central policy were agreed upon, smallholder water users may also resist limits, which drives up the cost of monitoring and enforcement.

2.2 Shrimp Farming as Climate Adaptation

Shrimp farming in Bangladesh involves flooding ponds with saltwater, generally drawn from the sea or saline rivers, at high tide and retaining it at low tide (Morshed et al., 2020). This takes place largely on paddy fields or salt pans on tidal flats, ringed by low earthen levees that are often built inside government flood-control polders. At the end of a growing season, salt ponds may be pumped out to allow for agriculture or to introduce new saline water for a new season of shrimp. In this setting, mixed-use plots are common; representative sampling suggests approximately two-thirds of shrimp plots are used intermittently for rice cultivation (Mondal et al., 2013). The most commonly farmed species is the black tiger shrimp (Penaeus monodon, locally *bagda*), which is stocked at low densities, with little or no external feed and reliance on tidal water exchange, so the shrimp largely subsist on the natural plankton and organisms that develop in the pond. The critical input is the seed stock, which is purchased from a network of hatcheries across coastal areas.

Figure 1 shows benchmark ranges for salt tolerance of the crops considered in

this paper. For non-tolerant rice, we use the standard threshold of 3 dS/m (soil saturation extract), above which yield declines by about 12 percent per additional dS/m (Maas and Hoffman, 1977). These non-salt-tolerant varieties remain the most common varieties in the dry season in the area of our study (Islam et al., 2024). For salt-tolerant rice, we use 8 dS/m, the upper end of the whole-season tolerance of BRRI dhan67 (Paul et al., 2024). Newer rice varieties such as BRRI dhan97 and BRRI dhan99 are available at higher salinity tolerances, but produce lower average yields and are still rarely adopted, making up less than 1 percent of current use (Debsharma et al., 2024).




Figure 1. Salinity Range for Rice and Shrimp

Black tiger prawns (*Penaeus monodon*) tolerate water salinity of roughly 5–35 ppt, and farming in Bangladesh typically takes place at 5–20 ppt (Rahi et al., 2021). In controlled trials with Bangladeshi stock, growth and survival were highest at 20 ppt and lower but viable at 5 ppt (Rahi et al., 2021). Converting parts per thousand measures into decisiemens for consistency with the ground-truth salinity measures we will take below, we convert to electrical conductivity at 25 °C using the Practical Salinity Scale (UNESCO, 1981), which places this range at approximately 9–46 dS/m. On these benchmarks, shrimp aquaculture becomes feasible at a salinity just above the whole-season tolerance of the most widely grown salt-tolerant rice varieties.

The primary inputs for saltwater shrimp farming are post-larval fry, saltwater, and feed. Fry are sourced from approximately 50 hatcheries in the southeastern district of Cox's Bazar and then transported and distributed to retailers throughout coastal areas. Shrimp yields in Bangladesh are low by international standards: about 0.35 tonnes per hectare, compared with roughly 3.9 tonnes in India and 4.3 tonnes in Thailand (Nurullah et al., 2024). This gap reflects the extensive nature of farming in Bangladesh, its reliance on the low-yielding black tiger shrimp at a time when major producers have shifted to the higher-yielding Pacific white shrimp (*Litopenaeus van-*

namei), and recurrent disease outbreaks, particularly white spot syndrome (Nurullah et al., 2024).[4] After farming, shrimp are sold at local fish markets to consumers or middlemen who transport to processing and packaging centers where up to 80% of shrimp is then exported (Islam et al., 2014). The primary export partners are the European Union, the United States, Japan, and Russia (Kabir et al., 2023).

International development efforts have also sought to expand shrimp farming in the region. The World Bank-financed Sustainable Coastal and Marine Fisheries Project, valued at US$281.6 million and running until 2023, supported the fisheries sector as a whole while targeting shrimp farmers in particular. It aimed to organize marginal shrimp farmers into production clusters and offered matching grants in the form of feed in exchange for farmers' investment in deepening their ponds, alongside training in good aquaculture practices. The Climate-Smart Agriculture and Water Management Project, valued at US$155.3 million and running from 2021 to 2026, improved flood control and drainage infrastructure, established cold storage facilities, and strengthened farmers' links with buyers, among other support to the sector (Kabir et al., 2023).

3 Data

This section describes the data used in the paper.

3.1 Household Survey

We conducted a household agricultural survey between May and June of 2025 covering 739 households across fourteen unions from the Khulna and Satkhira districts of Bangladesh. To select the sample, enumerators first identified marginal plot triples within villages where they could observe an active saltwater shrimp pond adjacent to two agricultural plots. This construction (i.e., a central agricultural plot with a saltwater shrimp pond on one side and a traditional agricultural plot on the other side) allowed us to collect salinity readings that could provide a gradient at both an agriculture–agriculture boundary and an agriculture–aquaculture boundary. After selecting the plots, enumerators identified the plot operators to interview. After obtaining consent and completing the modules with the operators, they returned to the plots to take salinity readings.

[4] In *extensive* systems, shrimp are stocked at low densities and depend mainly on the natural productivity of the pond, with little or no supplemental feed, minimal aeration, and water exchanged with the tides. *Intensive* systems stock shrimp at much higher densities and rely on formulated feed, mechanical aeration, and active management of water quality.

3.2 Salinity Readings

Enumerators collected salinity readings using a portable salinity meter with an electrical conductivity range of 0–10 mS/cm. The meter has a metal probe that is inserted directly into moist soil, where it passes a small electrical current between its tips and measures how easily that current flows. Because dissolved salts carry electricity, saltier soil conducts the current more readily, and the meter converts this into a salinity reading shown on its screen, which is then entered into the survey by an enumerator. The primary advantage of this type of meter is that it provides a nearly instantaneous reading and requires minimal training to operate. However, it is important to note that the meters measure overall electrical conductivity, which will depend on soil moisture, texture, and compactness. These sensors are less accurate than lab tests, but take very little time (about 15 seconds per reading), and have close to zero marginal cost. To improve quality, we calibrated all meters using a standardized saline solution before the survey was conducted and took each reading twice. We averaged the two readings before using them in the analyses presented in this paper. The soil salinity reading protocol for enumerators is reported in Appendix D.

May was chosen for the timing of the survey because it was immediately before the beginning of the monsoon season and thus gave us an opportunity to observe salinity at or near its peak, following its accumulation throughout the dry winter period. The salinity data presented here are therefore best interpreted as showing a gradient between saline and non-saline plots at or near its greatest extent (Rahman et al., 2025).

3.3 River Data

River locations come from OpenStreetMap (OpenStreetMap, 2026), from which we keep only features tagged as [waterway=river], excluding smaller channels such as streams, canals, drains and ditches. For each plot in our satellite analysis we compute the shortest distance from the plot's boundary to the nearest river segment in a projected metric coordinate system (EPSG:3106).

3.4 Satellite Data

The analysis of remote sensing data in this paper combines two data sources to first delineate plots, and then categorize those plots as either agricultural land or aquaculture.

Field boundaries are delineated using preprocessed high-resolution Planet basemaps,

which are single seamless images that aggregate cloud-free pixels from repeated satellite passes over a common area. Because this imagery is essentially a mosaic of different time periods, it can be used to recognize permanent and semi-permanent features of the landscape, such as the boundaries between plots, but cannot be used to categorize the use of the plot during any season. At Bangladesh's latitude, the pixels captured by the sensors on the satellite will correspond to 4.4 meters. We process this imagery by applying the Delineate Anything model, a machine learning approach that performs instance segmentation (i.e., identifies individual plots, rather than the lines between plots). The algorithm is pre-trained on a sample of 73 million plots across the world and is discussed further in Appendix C. Because the mosaics are already cloud composites, they are an ideal sample on which to extract plot geometry. We retain the resulting field polygons as the fixed spatial units of the panel. Here, we abstract away from the division of one plot into multiple plots or the combination of multiple plots into one larger plot during the span of the satellite portion of our analysis (i.e., 2017–2025). Panels 1 and 2 of Figure 2 show the first step of the approach. Panel 1 displays the base map mosaic from Magurkhali Union, one of the 51 unions that make up the sample. Panel 2 shows the delineated plots—distinct polygons that outline the boundaries of recognized plots. We drop those polygons that fall above the 99th percentile of plot size.

Delineated polygons make up the unit of analysis of the satellite analysis in Section 6, and also provide the network over which we propose that the physical salinity externality may take place. It is important to note here that classical algorithms (such as those in the watershed family) which, rather than identifying individual polygons, divide regions into units that fully tile space would be insufficient here because they would mechanically give every 'plot' which was defined a neighbor, potentially over saturating the links through which spillovers may happen.




Figure 2. Illustrative Plot Classification - Magurkhali Union

To measure land use over time, we use Sentinel-2 satellite imagery, a lower-resolution but higher-frequency dataset from a European satellite program with a spatial resolution of 10 to 20 meters per pixel depending on the spectral band. For every plot from 2017 to 2025, we compute two indices from the Sentinel-2 bands. The first is the Modified Normalized Difference Water Index (MNDWI), a canonical formula combining a green and a shortwave-infrared band that produces high values over standing water and low values over vegetation and soil, which we use to detect ponds.[5] Intuitively, this index captures that areas which are covered in surface water will absorb shortwave infrared (SWIR) light and reflect relatively more green light. The second index constructed for each plot is the Normalized Difference Vegetation Index (NDVI), likewise a canonical greenness measure that is high over healthy vegetation and low over bare or waterlogged ground, which we use as a proxy for agricultural productivity.[6]

We discard any Sentinel-2 scene with more than 50 percent cloud cover. Within the retained scenes, we mask individual pixels flagged as cloud, cloud shadow, or cirrus – thin high-altitude cloud – using the imagery's built-in scene-classification band. We then combine the surviving clear-sky pixels within each season by taking the median value at each location, which suppresses any residual haze, and we then average the clear pixels falling inside each field boundary to obtain one MNDWI and one NDVI value per field during the Rabi (i.e., dry) season of each year, aggregating readings from 1 November to 28 February each year. We chose the dry season to gain the cleanest differentiation between plots that are flooded with saltwater and those that are not flooded, but may have residual water from other uses. Plot-seasons based on fewer than three clear pixels are flagged as unreliable and dropped. Finally, a plot is classified as in aquaculture in a given season when its MNDWI exceeds a season-specific cutoff chosen as follows: we apply Otsu's method to the distribution of MNDWI values within each season and year, which selects the threshold that best separates the two modes of the distribution rather than imposing a fixed cutoff of zero.[7] Appendix Figure B5 shows dry season histograms of the MNDWI readings for all plots in the sample, grouped by year. The vertical line shows that year's cutoff as

[5]The Modified Normalized Difference Water Index is defined as $\text{MNDWI} = \dfrac{\rho_{\text{Green}} - \rho_{\text{SWIR}}}{\rho_{\text{Green}} + \rho_{\text{SWIR}}}$, where ρ_{Green} and ρ_{SWIR} are reflectance in the green and shortwave-infrared bands.

[6]The Normalized Difference Vegetation Index is defined as $\text{NDVI} = \dfrac{\rho_{\text{NIR}} - \rho_{\text{Red}}}{\rho_{\text{NIR}} + \rho_{\text{Red}}}$, where ρ_{NIR} and ρ_{Red} are surface reflectance in the near-infrared and red bands, respectively.

[7]Otsu's method chooses the threshold t^* that minimizes the intensity variance within the two groups it creates (equivalently, maximizes the between-group variance): $t^* = \arg\max_t \ \omega_0(t)\,\omega_1(t)\left[\mu_0(t) - \mu_1(t)\right]^2$, where $\omega_0(t), \omega_1(t)$ are the proportions of observations below and above t and $\mu_0(t), \mu_1(t)$ are their respective means.

determined by the Otsu method to be the cutoff between agriculture and aquaculture.

3.4.1 Satellite plot detection and classification quality

To conduct meaningful analysis using the satellite-derived panel, we need two conditions to hold: (i) the plot recognition algorithm is able to detect enough plots at a density that permits them to border one another frequently enough that we have sufficient variation in whether or not an agricultural plot is bordered by an aquaculture plot for an event-study approach; and (ii) the land use classification algorithm is not merely noise, i.e., it contains a sufficiently large signal about the actual use of that plot. Among the 51 unions that make up the satellite sample, areas designated as plots cover 36.8% of total space (see Table B1). Estimates place the share of land under cultivation at approximately 60%, implying that our delineated plots cover approximately 60% of cultivated land (Bari et al., 2024).[8]

Accurately classifying plots as agriculture or aquaculture is difficult in this setting because the water we are attempting to detect is shallow and murky, so errors may arise from either under- or over-classification of wetness. Appendix Figure B5 shows the distribution of MNDWI readings for our sample by year. The vertical lines show the water threshold chosen each year by Otsu's method, which selects the cutoff that maximizes the between-class variance of the index. Although the index is "normalized", the selected threshold is above zero in every year, and many plots have intermediate values. Thus, although hereafter we refer to plots above the threshold as *ponds* or as being in *aquaculture*, the classification is more generally interpreted as placing a plot above a data-driven threshold of pond likelihood. As an alternative, we also classify plots using a single manual threshold, MNDWI > 0.241, held fixed across all years. We choose this value to minimize the balanced error rate—the average of the false-positive and false-negative rates—against land use reported in our household survey for plots we can match to the satellite data. Because it lies above the Otsu threshold in every year, the manual threshold applies a stricter definition of a pond. The main empirical result of the paper (i.e. the event-study estimate of the effect of a neighbor's adoption of aquaculture on one's own adoption) is robust to this reclassification, as shown in Appendix Figure B7.

One additional concern specific to satellite-delineated plots is that the segmentation may split a single pond into two adjacent polygons. If it does, the two halves would fill with water in the same season and appear as a plot converting at the same moment as its neighbor, mechanically generating a spillover. To test for this, we

[8]This percentage includes both the city of Khulna which has very little agricultural land as well as mangrove forests in the Khulna Division, so it is best interpreted as a lower bound.

compare plot pairs that convert in the same season with other adjacent pond pairs. If two polygons were in fact one pond, they should be ponds in the same years and, while both are ponds, have nearly identical MNDWI readings that move together from year to year; we measure this only in years when both plots are ponds, since any two plots that convert together will have correlated readings over the full panel simply because of the shared switch. Same-season pairs are slightly more likely than other adjacent ponds to look like a single pond: 11% versus 8% under the strictest similarity criteria, and 21% versus 19% under the strict criteria (Appendix Figure B6). Some simultaneous conversions may therefore reflect split ponds, but the excess is small, and we show below that our results are robust to dropping these plots.[9] A final concern with the classification exercise is that some ponds in our setting may be used for freshwater aquaculture rather than saline shrimp farming, and our water index cannot distinguish the two. Three features of the setting limit how much this matters. First, freshwater aquaculture is considerably less common than brackish shrimp farming in our study area: in Khulna and Satkhira districts combined, shrimp (*bagda*) farms cover about 91,300 hectares, against about 28,400 hectares for freshwater prawn (*golda*) farms (Department of Fisheries, 2025). Second, where it does occur, the most common freshwater system rotates prawns in the wet season with boro rice in the dry season (Jamal et al., 2023), so these plots are typically under rice, rather than open water, during the dry-season window we use for classification. Third, to the extent that misclassified freshwater ponds do not affect their neighbors' land use, because they generate no salinity spillover, they attenuate our estimates toward zero. If they do affect their neighbors, for example by waterlogging adjacent fields, they contribute to the net externality that our event study measures and that the model's parameter δ is designed to capture.

4 A model of land use with a border externality

This section proposes a model of land allocation between rice and saline aquaculture in which a plot's returns depend on whether its neighbors are in aquaculture. It separates selection on comparative advantage from the externality, and shows how the externality propagates across a landscape.

4.1 Environment

There are plots which are each operated by an individual farmer. Plots, indexed by i, are contiguous, and each is either in rice ($d_i = 0$) or in saline aquaculture ($d_i = 1$).

[9] See Appendix Figure B12.

Plot i has an unobserved productivity in each use, a_i^R and a_i^S which are independently drawn from Fréchet distributions. These capture everything that makes a plot more or less profitable in rice or shrimp, respectively beyond its observed characteristics: soil, drainage, access to fresh or saline water, and the skill and capital of whoever farms it.

We consider a plot to be exposed if it borders another plot that has adopted shrimp:

$$b_i = \{\text{plot } i \text{ shares a boundary with at least one shrimp plot}\}. \tag{1}$$

4.2 Payoffs and land use

Profit for each plot is productivity times observable shifters:

$$\pi_i^S = a_i^S \cdot \underbrace{e^{w_i'\gamma^S}}_{\text{characteristics}} \cdot \underbrace{e^{-c}}_{\text{conversion cost}}, \tag{2}$$

$$\pi_i^R = a_i^R \cdot \underbrace{e^{w_i'\gamma^R}}_{\text{characteristics}} \cdot \underbrace{e^{-\delta b_i}}_{\text{net damage}}, \tag{3}$$

where w_i contains plot size and distance to the river. In order to adopt shrimp, a plot pays a conversion cost which is common to all plots. If a plot shares a boundary with at least one shrimp plot (i.e. $b_i = 1$), then profit shifts by $e^{-\delta b_i}$, the relative damage to rice profits caused by the boundary externality. Each plot is allocated to the use that pays more:

$$d_i = 1 \iff \underbrace{\log a_i^S - \log a_i^R}_{\Delta_i} > c - \delta b_i - w_i'\gamma, \qquad \gamma \equiv \gamma^S - \gamma^R. \tag{4}$$

The breakeven condition is thus the relative productivity draws a_i^S, a_i^R such that profits from the new technology, shrimp, are equal to profits in the old technology, rice. Δ_i is the plot's comparative advantage in aquaculture. An aquaculture neighbor lowers the threshold by δ without changing Δ_i.

It is important to note that δ is a net damage effect: the loss of rice profit from an aquaculture neighbor plus any gain in shrimp profit from one. Both make aquaculture relatively more attractive, and land use responds only to their sum; writing δ on rice profit is a normalization. The multiplicative form makes the effect proportional, so a given exposure changes returns by the same fraction on every plot. The conversion cost c is common across plots. It cannot be separated from the intercept of $w_i'\gamma$ and is absorbed into it; plot-level variation in conversion costs is absorbed into Δ_i.

4.3 Types and allocation probabilities

Productivities are independent Fréchet draws with common shape θ, and unit scale since w_i enters explicitly:

$$\left(a_i^k \leq a\right) = \exp\left(-a^{-\theta}\right), \qquad k \in \{R, S\}. \tag{5}$$

Then Δ_i is logistic with scale $1/\theta$, and with Λ the logistic function,

$$P_i \equiv (d_i = 1 \mid b_i, w_i) = \Lambda\big(\theta(w_i'\gamma - c + \delta b_i)\big). \tag{6}$$

θ indexes the dispersion of comparative advantage. When θ is low, plots differ widely, and land use responds little to a change in returns. In other words, plot fundamentals are the main determinant of land use. Alternatively, when θ is high, plots are close substitutes and the same change reallocates more land.

As is standard with the Fréchet setup, realized profit $\pi_i = \max\{\pi_i^S, \pi_i^R\}$ has the same distribution whichever use is chosen,

$$(\pi_i \leq x \mid d_i = k) = \exp\Big[-\left(e^{\theta(w_i'\gamma^S - c)} + e^{\theta(w_i'\gamma^R - \delta b_i)}\right) x^{-\theta}\Big], \qquad k \in \{R, S\}, \tag{7}$$

so the dispersion of realized returns identifies θ.

Independence across plots is a substantive assumption. It rules out, for example, a single operator farming several adjoining plots, whose joint conversion the model would read as a cascade. In southern Bangladesh, land markets are fragmented, and it is common for a household operating multiple plots to be unable to rent or purchase them in a contiguous area. This is consistent with our household survey, which documents that farmers are operating plots an average of 153 meters apart as compared to 19 meters apart for true neighbors.

4.4 Interpretation

It is important to note that the model does not require the externality to be salinity. δ captures any channel through which an aquaculture neighbor changes a plot's relative returns, whether negative for rice, such as salinity, waterlogging or disease, or positive for shrimp, such as shared embankments and canals, information, or access to buyers and inputs. Changes in land use derived from the satellite data identify only the net effect. While this net externality is sufficient to simulate, the split between these channels matters for over- or underadoption and thus the correct policy response. We return to these concerns in Section 7.

The model also does not include time. An equilibrium describes where land use settles given current returns, not the path or speed of adjustment, and plots respond to their neighbors without anticipating future conversions. We justify this modeling approach as follows: Because the focus of our study, and the event study estimates that will be used to estimate the model, are both marginal plots, the plot represented by the model is not in an area where saltwater intrusion is so severe that plots are in aquaculture the entire year. As monsoon rains arrive in June each year, salt levels are reset, and at the beginning of the next dry season, plots must again make the decision of whether to adopt shrimp or rice. As such, year-over-year the decision is reversible; thus, there is no scope for option value of waiting to adopt in the setting or the model. Further, there is no anticipatory action, as supported in this setting, as evidenced by the lack of pre-treatment trends in the event study estimates in Section 6.

Even without pre-treatment trends, we might be concerned about anticipatory action if farmers believe salinity may increase in a future period, and so adapt to avoid the negative effect of climate change directly. While we cannot rule this out explicitly, the year-over-year changes in salinity in this setting account for $\approx 0.5\%$ decreases in rice yields per year (Dasgupta et al., 2015). This is two orders of magnitude less than the yield losses farmers report expecting from rice yields in the year following their neighbor's first adoption of shrimp farming (see Section 5), so we assume decision-making is driven by plot fundamentals and the externality alone. Thus, the model is best interpreted as a medium-run framework, net of climate change: productivity in both uses is held at current levels, so the model does not capture a secular decline in rice returns from rising salinity.

5 Stylized Facts

This section presents brief demographic information about the farmers and plots represented in the household survey. Then it introduces three stylized facts about the shrimp farming industry in Bangladesh.

Key features of the households surveyed are shown in Panel A of Table 1. Respondents are selected from among marginal plot operators (i.e., who are cultivating rice near other shrimp plots or cultivating shrimp near other rice plots).[10]. Both groups are of similar age, sex, and education. They are very concerned about climate change. On the 1–3 scale for climate change concern, almost all respondents in both groups considered climate change a "1 = very serious threat". Respondents engaged

[10]For greater detail about sample selection, see Section 3

Table 1. Farmer and Plot Demographics

Panel A: Households			
	Shrimp farming	Non shrimp farming	p-value
Age	45.7	47.0	0.151
	(0.6)	(0.7)	
Male	0.970	0.962	0.591
	(0.009)	(0.010)	
Individual income (1,000 taka)	282.5	217.3	0.000***
	(12.5)	(10.2)	
Completed secondary school	0.297	0.258	0.238
	(0.023)	(0.024)	
Muslim	0.670	0.568	0.004***
	(0.024)	(0.027)	
Number of non-homestead plots	2.46	2.02	0.000***
	(0.05)	(0.06)	
Climate change concern (1–3)	1.24	1.21	0.341
	(0.02)	(0.02)	
Panel B: Plots			
	Shrimp pond	Agriculture	p-value
Plot size (decimals)	131.5	69.6	0.000***
	(7.5)	(3.4)	
How acquired			
Bought	0.071	0.091	0.182
	(0.012)	(0.010)	
Rented	0.478	0.384	0.001***
	(0.025)	(0.018)	
Inherited	0.349	0.444	0.000***
	(0.024)	(0.019)	
Year acquired	2012.6	2012.1	0.459
	(0.6)	(0.5)	
Water source			
Rainfed	0.482	0.746	0.000***
	(0.026)	(0.017)	
River	0.367	0.082	0.000***
	(0.026)	(0.011)	
Canal	0.296	0.029	0.000***
	(0.022)	(0.006)	
Pond	0.002	0.023	0.000***
	(0.002)	(0.005)	
Groundwater	0.379	0.716	0.000***
	(0.025)	(0.020)	
Plot count	496	1138	
Household count	394	345	

Notes: Means are reported with standard errors in parentheses. Panel A: robust standard errors; a household is classified as shrimp farming if it has at least one shrimp pond. Climate change concern is coded 1 = very serious threat, 2 = somewhat serious threat, 3 = not a threat at all. Panel B: cultivable/arable plots and shrimp ponds only; standard errors clustered by household. The last column reports the p-value of the difference in means between columns 1 and 2. One decimal equals 1/100 acre. Acquisition categories are not exhaustive (user-right, gifted, government land, and other are omitted), so rows do not sum to one. Water source categories are not mutually exclusive. The household count in Panel B is the number of households with at least one plot of that type. * $p < 0.10$, ** $p < 0.05$, *** $p < 0.01$.

in shrimp farming report 30% higher individual income (282.5 thousand taka annually for shrimp farming households vs 217.3 for non-shrimp farming households). This gap is also reflected in the number of plots operated by each group: shrimp farming households operate 2.46 non-homestead plots relative to non-shrimp farming households' 2.02. Shrimp farming households are 10 percentage points more likely to be Muslim. Panel B shows plot differences across plots that operate as shrimp ponds at any time of the year and those that do not. Shrimp ponds are much larger, averaging 1.32 acres relative to 0.70 acres for agricultural plots. Land allocated to either use is rarely purchased outright (7.1% for shrimp ponds vs 9.1% for agriculture). The two dominant forms of plot acquisition are renting (47.8% for shrimp and 38.4% for agriculture) as well as inheritance (35% for shrimp and 44.4% for agriculture). These plots have been operated by their current owners for, on average, 15 years at the time of the survey in 2025. There is no dominant water source for shrimp ponds, which use a combination of rainfall (48.2%), rivers (36.7%), canals (29.6%) and groundwater (37.9%). Agricultural plots use rainfall (74.6%), rivers (8.2%), canals (2.9%) and groundwater (71.6%).

Fact 1: Farmers believe shrimp causes a salinity production externality

Figure 3 presents the distributions of two elicited outcomes—perceived water salinity (left panel) and subjective rice health (right panel)—under two scenarios. Both measures are reported on scales from 0 to 7. Salinity runs from 0 (no salt in the water) to 7 (the saltiest water the respondent has ever encountered). Rice health is an inverted scale: respondents are shown images corresponding to progressively poorer plant growth, so 0 denotes the healthiest rice and 7 the least healthy, with higher values reflecting the visible damage associated with greater salinity. The images shown to farmers are drawn from Grattan et al. (2002) and are shown in Appendix Figure B4. The green bars show the status-quo distribution: for each plot, respondents report their current water access and their current rice condition. The red bars show the distribution under a hypothetical in which the plot bordering theirs is converted to saline aquaculture. In both panels, the distribution shifts markedly to the right when a neighboring plot converts—perceived salinity moves from an average of 3.0 to 4.2, and rice health shifts from an average subjective score of 2.2 toward the damaged end of the scale at 4.9. In other words, farmers expect that a neighbor's conversion will both raise the salinity of their own water and worsen the health of their rice.

This is important for two reasons. First, it shows that farmers understand the spillover process, at least directionally: their subjective expectations move in exactly the direction the agronomic evidence on lateral salt transmission would predict (Yu




Figure 3. Salinity and Rice Outcomes After Neighbor Conversion

et al., 2024; Braaten and Flaherty, 2001), and they correctly link the intermediate channel (rising salinity) to the downstream outcome they care about (falling rice yields). Their beliefs are therefore not naive—they encode a coherent causal model of how a neighbor's land use propagates onto their own plot. Second, and consequently, these perceived spillovers are likely to enter farmers' own decisions. If a farmer anticipates that a neighbor's conversion will salinize their water and depress their rice yield, they may adjust their own cropping choices—or their own decision to convert to shrimp aquaculture—in response to what they expect their neighbors to do. This sets up a strategic, potentially self-reinforcing dynamic in which conversion decisions are interdependent across neighboring plots, which motivates the analysis of conversion behavior that follows.

In addition to this qualitative exercise, we ask farmers a hypothetical question in which they are told to consider a plot that is the same size as one of their randomly chosen existing plots that is growing rice and does not border any shrimp farms.

When asked whether yields from that plot will change (from a benchmark of 100 kg) in the year following a neighbor's conversion to shrimp farming, they report an expected decrease in yields of 48%. This shows that the qualitative results from Figure 3 map onto a large quantitative decrease. We will use this quantitative gap to estimate the model of crop choice in Section 7.

Fact 2: The ground-truth salinity gradient exists

Figure 4 reports average soil salinity (green bars, dS/m) alongside a reading taken directly from the saline aquaculture pond (blue bar), pooled across all sample plots and measured in May–June 2025. The four bars are arranged by distance from the saline source. The leftmost bar is the reading inside the saltwater aquaculture ("wet") plot itself, which averages 10.3 dS/m. The two central bars are boundary readings taken on a plot that simultaneously borders a saline aquaculture plot and an agricultural plot: the wet/dry boundary (4.3 dS/m) and the dry/dry boundary (3.9 dS/m). The rightmost bar is taken from an agricultural ("dry") plot that borders only other agricultural land (3.4 dS/m). Two features stand out. First, there is a steep gradient between the aquaculture reading and the agricultural readings—salinity more than halves crossing from the pond to the nearest agricultural boundary. Second, among the agricultural readings there is a smaller but consistent declining gradient as one moves further from the saline plot, from 4.3 to 3.9 to 3.4 dS/m.

This pattern is suggestive evidence that local salinity on a farmer's land may be shaped by the land-use decisions of neighboring farmers, rather than by that farmer's own choices alone. An important caveat is that this is cross-sectional evidence, so the direction of causation is not identified. One competing explanation is selection on a pre-existing, static salinity gradient: if some areas were already saltier, those plots may have been the ones converted to saline aquaculture, leaving the underlying spatial gradient—not the conversion—as the true driver of what we observe. That said, the agronomic literature makes lateral transmission of salt between adjacent plots physically plausible. Controlled monitoring and numerical modeling of onshore aquaculture farms document both vertical and lateral saltwater intrusion, showing that salt infiltration at ponds gradually invades neighboring farmland unless physically blocked by a guard wall, though evidence from Bangladesh is mixed (Ayers et al., 2017; Yu et al., 2024). Salt-balance studies of inland shrimp ponds similarly find that lateral seepage through pond walls and direct discharge of saline water into irrigation canals are the most significant salt-transfer pathways, raising salinity in adjacent canals and rice paddies above yield-damaging thresholds (Braaten and Flaherty, 2001). In this sense, salt can move across plot boundaries much as fertilizer




Figure 4. Salinity Readings Across Test Plots

or other soil inputs can spill over between neighboring fields. To distinguish this dynamic, spillover-based account from the static-gradient explanation, we supplement the cross-sectional comparison with the event studies below.

Fact 3: Saltier bordering plots have lower rice yields and profits relative to plots that do not border shrimp ponds, controlling for plot size

If proximity to a saline aquaculture pond raises the salinity of neighboring agricultural land, as Figure 4 suggests, then we should expect elevated salinity to depress the productivity of the affected plots. Figure 5 tests this directly, comparing annual rice yields—measured in kilograms per decimal (one decimal being 1/100 of an acre)—across agricultural plots according to whether they border a shrimp pond. Plots that do not border a pond yield approximately 23.5 kg/decimal, compared with 21.8 kg/decimal for plots sharing a boundary with a pond: a difference of -1.71 kg/decimal, or roughly seven percent, significant at the ten percent level. The salinity gradient documented in Figure 4 thus appears to carry through to agricultural outcomes—plots exposed to a neighboring pond are not only saltier but measurably less productive.

As with the salinity readings, this comparison is cross-sectional and does not identify a causal effect. Border and non-border plots may differ systematically for reasons unrelated to the pond—in unobserved soil quality, location, or the characteristics of the farmers who operate them—and the same selection concern raised for the salin-




Figure 5. Rice Yields Across the Plot Distribution

ity gradient applies here: if plots that were already lower-yielding were the ones near which aquaculture was established, the yield gap would reflect pre-existing differences rather than the pond's influence. The figure should therefore be read as descriptive corroboration that the salinity spillover has real economic consequences, motivating the panel research design developed below, which uses the timing of neighbors' conversions to isolate the causal effect on land use and productivity.

6 Empirical Strategy & Event Study

We estimate whether a plot's conversion to shrimp aquaculture (hereafter a *pond*) induces its neighbors to convert. The central challenge is that adoption is spatially clustered for reasons unrelated to peer influence—soil salinity, water access, and proximity to processing infrastructure vary smoothly across space and jointly determine the profitability of aquaculture for neighboring plots. A naive comparison of conversion rates between plots with and without pond neighbors would conflate genuine contagion with these shared locational fundamentals. Our strategy exploits the *timing* of neighbors' conversions within a difference-in-differences design, comparing a plot's own adoption trajectory before and after a neighbor converts with that of plots in the same small neighborhood whose neighbors have not converted.

The unit of observation is a delineated plot i observed in Rabi (dry) season t, for $t = 2017, \ldots, 2025$. Let $Y_{it} = 1$ if plot i is classified as a pond in period t and 0

otherwise. Each plot's treatment date E_i is the first period in which any plot sharing a boundary with i is classified as a pond. We estimate the event-study model:

$$Y_{it} = \alpha_i + \lambda_{h(i)t} + \sum_{k \neq -1} \beta_k \, D_{it}^k + X_i' \eta_t + \varepsilon_{it}, \tag{8}$$

where D_{it}^k indicates that plot i is k periods from the first period in which a neighbor is a pond, α_i are plot fixed effects, and $\lambda_{h(i)t}$ are hexagon-by-period fixed effects for the hexagon $h(i)$ containing plot i. The hexagons form a regular grid over the study area, sized so that each contains on average 20 plots of the estimation sample, which gives hexagons 1.03 km across; hexagons containing a single plot are dropped. X_i contains log distance to the nearest river and log plot area, each interacted with period so that their effects can vary over time. The period $k = -1$ is omitted as the reference, so each β_k measures the change in the probability that i is a pond at event time k relative to the period immediately preceding a neighbor's conversion. Extreme leads and lags are binned at ± 5 years, so that the endpoint coefficients are not identified off a single treatment cohort.

The plot fixed effects α_i absorb all time-invariant determinants of aquaculture suitability—soil type, elevation, baseline salinity, distance to water channels, and unobserved operator characteristics—so that identification comes entirely from within-plot changes over time. The hexagon-by-period fixed effects $\lambda_{h(i)t}$ absorb any shock common to a neighborhood of about 1 km in a given year, such as an advancing salinity front, local waterlogging, or shifts in local shrimp prices, as well as region-wide movements in prices and aggregate weather. Identification therefore comes only from differences in the timing of neighbors' conversions among plots in the same hexagon and year. The period interactions $X_i' \eta_t$ allow plots nearer rivers and larger plots to follow different trends. Because Y_{it} is binary, (8) is a linear probability model and the β_k are interpreted in percentage-point units.

A well-defined event study requires that treatment onset be observable within the panel. We therefore restrict the estimation sample in two ways. First, we exclude plots with no delineated neighbors, as neighbor exposure is undefined for them. Second, we drop plots that already border a pond in the first period of the panel. These plots are always-exposed: their treatment date is left-censored, and no clean pre-period exists against which to measure the effect of a neighbor's conversion. The retained sample consists of plots that begin the panel with no pond neighbor, for which the first appearance of a pond neighbor is an observed transition rather than a pre-existing state. The estimand is accordingly local to the expansion frontier—plots at the margin of the aquaculture landscape—and does not describe the already-saturated core.

Because treatment timing is staggered and treatment effects may be heterogeneous across adoption cohorts, the two-way fixed-effects estimator of the β_k can be contaminated by comparisons in which already-treated plots serve as effective controls for later-treated plots; with dynamic effects, these comparisons enter the estimate with potentially negative weights (Goodman-Bacon, 2021; de Chaisemartin and D'Haultfœuille, 2020). We therefore report the interaction-weighted estimator of Sun and Abraham (2021), which estimates a separate coefficient for each adoption cohort and event time and aggregates them using each cohort's share of observations at that event time, so that all weights are non-negative. Never-treated plots—those whose neighbors never convert during the panel—form the control group. Because both land use and its unobserved determinants are spatially correlated, we compute standard errors following Conley (1999), allowing for correlation across plots within 2 km of each other in the same period, with weights declining linearly in distance, and for arbitrary correlation within a plot across periods.

6.1 Identifying Assumption

The identifying assumption is parallel trends: absent a neighbor's conversion, treated plots would have followed the same own-adoption path as control plots. This is not directly testable, but its plausibility is assessed through the pre-treatment coefficients $\{\beta_k\}_{k<-1}$, which should be jointly indistinguishable from zero if treated and control plots were on common trajectories before treatment. The salient threat is a plot-specific, time-varying shock that drives both a neighbor's conversion and the plot's own—most plausibly a localized intrusion of saline water that renders a cluster of contiguous plots newly suitable for aquaculture in the same period. Such a shock would generate the appearance of contagion absent any causal peer effect. Time-invariant salinity differences are absorbed by α_i and region-wide salinity trends by λ_t; only *local, time-varying* salinity movements pose a threat.

6.2 Extensive Margin Effects

Figure 6 presents the event-study estimates for the probability that a plot is classified as a pond. Following a neighbor's conversion, a plot's own probability of being a pond rises sharply and immediately, by 18.0 percentage points in the period following conversion. The effect persists across all subsequent periods, ranging between 0.14 and 0.22 over the following five years, and shows no evidence of decay. Own-conversion thus appears to be a durable rather than transitory response: once a neighbor converts, the elevated likelihood of the plot itself becoming a pond is sustained rather

than fading over time. The pre-treatment coefficients are flat and statistically indistinguishable from zero, consistent with the parallel-trends assumption.




Figure 6. Aquaculture Event Study

Notes: The figure plots Sun and Abraham (2021) event-study estimates of own aquaculture status against years since a plot's first adjacent neighbor became a pond, for plots with no adjacent pond at baseline. The specification includes plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls, and shaded bands show 95% (dark) and 99% (light) confidence intervals using Conley standard errors with a 2 km cutoff.

The effect emerges at event time zero—in the same period as the neighbor's conversion—rather than building gradually over subsequent years. This timing is informative about the mechanism. A gradual, increasing response would be consistent with social learning, in which a farmer observes a neighbor's outcomes over several seasons before updating and eventually converting. The immediate response we observe instead suggests that the second farmer acts concurrently with, or very shortly after, the neighbor's conversion. Such timing is consistent with an anticipatory mechanism: if farmers believe that an adjacent pond will depress their own yields—for instance, through saline intrusion—the optimal response is to convert immediately rather than to wait and learn, since remaining in agriculture beside a pond becomes less viable as soon as the neighbor converts.

6.3 Intensive Margin Effects

Figure 7 presents the event-study estimates with NDVI—a satellite proxy for vegetative greenness, and hence crop productivity—as the outcome. This specification is estimated only on the sample of plots that had no pond neighbor at baseline and that never themselves convert to a pond at any point in the panel, so that NDVI always reflects standing agriculture rather than open water. Following a neighbor's conversion, greenness falls sharply and immediately: the effect is -0.019 at the time of conversion, indicating a contemporaneous decline in crop productivity. The effect then attenuates steadily over the following years, converging back toward zero by year five. The pre-treatment coefficients are flat and indistinguishable from zero, consistent with parallel trends. As with the extensive-margin result, the decline appears at event time zero rather than emerging gradually, indicating that the productivity cost of an adjacent pond—plausibly through saline intrusion—is felt immediately rather than accumulating slowly. The subsequent recovery toward baseline suggests that farmers who remain in rice may, over time, adopt mitigating practices that partially restore greenness to its pre-conversion level.

Two features of the sample warrant caution in interpreting these magnitudes. Because the estimation is restricted to plots that never convert, the sample is not representative of all plots initially exposed to a converting neighbor: precisely those plots that experience the largest productivity losses may be the ones most likely to abandon rice and convert to shrimp themselves, thereby exiting the sample. Conditioning on non-conversion therefore selects on an outcome that is itself affected by treatment, and it does so in a direction that biases the estimated harm toward zero—the worst-affected plots are absent. The coefficients reported here should thus be read as a conservative lower bound on the true productivity cost of a neighboring pond, capturing the effect among the subset of farmers who chose to remain in agriculture despite it.

6.4 Robustness

To examine how the effect of a neighboring conversion decays with distance, we re-estimate the event study separately by distance rings, assigning each plot to a ring according to the distance between the plot and the first pond to appear within 500 meters. Appendix Figure B11 reports the results for four rings: plots whose first nearby pond appears within 20 meters (panel a), between 20 and 50 meters (panel b), between 50 and 200 meters (panel c), and between 200 and 500 meters (panel d). In each panel, the outcome is an indicator for whether the plot is itself a pond, and event




Figure 7. Vegetation Index Event Study

Notes: The figure plots Sun and Abraham (2021) event-study estimates of Rabi-season NDVI against years since a plot's first adjacent neighbor became a pond, for plots that are never classified as aquaculture and have no adjacent pond at baseline. The specification includes plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls, and shaded bands show 95% (dark) and 99% (light) confidence intervals using Conley standard errors with a 2 km cutoff.

time is measured relative to the year the first pond appears within 500 meters. The effect is concentrated among plots closest to the new pond. For plots within 20 meters, the probability of conversion rises by 18.0 percentage points on average over the post-period, with no evidence of differential trends before the neighboring conversion. Beyond immediate neighbors, we find no meaningful spillover. The average post-period effects are 0.24 percentage points for non-adjacent plots within 50 meters (SE 0.54), 0.23 percentage points between 50 and 200 meters (SE 0.25), and 0.62 percentage points between 200 and 500 meters (SE 0.22). All three estimates are small, and their confidence intervals rule out effects larger than about one percentage point in either direction. The absence of an effect on nearby but non-adjacent plots is consistent with a spillover that operates only between immediate neighbors, as would be expected if saline water spreads primarily through seepage across shared boundaries.

6.5 Alternative Mechanisms

Our interpretation is that a neighbor's pond contaminates adjacent land with saline water, lowering the expected returns to rice and inducing conversion. We consider two alternative explanations for the conversion response.

6.5.1 Multiple Plots Converting at the Same Time Because of Shared Ownership

If adjacent plots belong to the same operator, an apparent neighbor effect could instead reflect a single decision applied across several of the operator's own fields. This is unlikely in our setting. Among surveyed shrimp farmers who cultivate more than one plot, the median distance between their own plots is 153 meters, compared with 19 meters between a plot and its neighbors. This is consistent with the fragmented land markets documented in southwestern Bangladesh (Haque, 2025), in which a single household's holdings are typically scattered rather than contiguous. Moreover, shared ownership would predict conversions concentrated in the same period as the neighbor's conversion, whereas the effects in our event study persist and remain large in the years after the neighboring pond appears.

6.5.2 Information Spillovers Between Plot Owners

Farmers may also learn about the profitability of shrimp from neighbors who have converted and follow them. Such learning likely plays some role: 18.2% of converters report that someone in their village recommended shrimp farming, compared with 1.5% who cite advice from an NGO or the government (Appendix Table **??**). However, three features of our results suggest that learning does not drive the effects we estimate. First, the effect appears immediately, in the year of the neighbor's conversion, before a neighbor's returns could be observed. Learning from a neighbor's experience would instead produce a response that builds over subsequent seasons. Second, information spillovers would not be expected to decline sharply with physical distance, since farmers report discussing shrimp farming more with landholders who do not border their plots than with those who do ($p = 0.03$). By contrast, the effect we estimate is concentrated among plots that share a boundary with the new pond (Appendix Figure B11), as a physical spillover would predict. Third, adopters do not appear to hold more optimistic beliefs about returns, as they would if conversion spread through information about profitability: at the time of the survey, shrimp-farming households expect rice and shrimp prices similar to those of non-shrimp households(Appendix Figure B2).

6.6 Heterogeneity

The effect of a neighbor's conversion varies with a plot's characteristics in ways that bear on both the model and the mechanism. We re-estimate the main specification separately for plots above and below the median distance to the nearest river and above and below the median plot area in the estimation sample. To test whether the effects differ between groups, we compute standard errors for the difference that allow for spatial correlation between plots in different groups, since plots in each pair of groups sit side by side.

Distance to rivers. The response to a neighbor's conversion is larger among plots close to a river. Splitting the sample at the median distance to the nearest river (1,140 meters), a neighbor's conversion raises the probability that a plot is a pond by 19.3 percentage points (SE 1.5) among plots near a river and by 14.7 percentage points (SE 1.4) among plots farther away (Appendix Figure B9). The difference of 4.7 percentage points is statistically significant at the 5 percent level (SE 2.0, $p = 0.019$). This pattern is consistent with the model. Proximity to a river lowers the cost of bringing saline water onto a plot, placing nearby plots closer to the margin between rice and aquaculture, where a given shift in relative returns reallocates more land. It is also consistent with the salinity channel: soil and groundwater near tidal rivers may be saltier at baseline, so rice on these plots is closer to the threshold beyond which yields decline (Maas and Hoffman, 1977), and additional seepage from a neighboring pond is correspondingly more damaging. One caveat is that plots far from rivers show positive pre-period coefficients. Thus, part of their response may reflect a return to aquaculture on land previously used for shrimp rather than first-time adoption.

Plot size. Smaller plots respond somewhat more strongly than larger ones. Splitting at the median plot area (3,273 m^2), the post-period effect is 19.6 percentage points (SE 1.5) for smaller plots and 17.1 percentage points (SE 1.2) for larger plots (Appendix Figure B8). The difference of 2.5 percentage points is marginally significant (SE 1.5, $p = 0.097$). Its direction is informative because the model predicts the opposite: in the survey land-use logit, larger plots are more suited to aquaculture, which places them closer to the margin where a given shift in relative returns moves more land. A boundary externality offers a natural explanation for the reversal. Because a plot's perimeter grows less than proportionally with its area, seepage across a shared border affects a larger fraction of a small plot than of a large one. The same geometry explains why larger plots are about twice as likely to be exposed during the panel (3,875 against 1,950 treated plots): a longer boundary has more neighbors

that might convert. The model's δ is common to all plots and abstracts from this geometry, which would instead predict that the effect rises with the share of a plot's perimeter that borders the converting neighbor.

7 Estimation and Counterfactuals

7.1 Estimation

The model in Section 4 is characterized by four sets of parameters, which we estimate by combining the household survey with the satellite panel. Table 2 summarizes the source of identification for each.

The dispersion parameter θ governs how heterogeneous plots are in their comparative advantage between rice and shrimp. Because productivity draws are Fréchet, θ is identified from the dispersion of log output value across surveyed plots. The slopes of the relative suitability index, γ, capture how plot size and distance to the river shift the relative returns to shrimp, and are estimated from a land-use logit on the survey sample. The survey cannot identify the level of γ, because marginal sampling selected plots on both sides of the rice–shrimp boundary and so excluded unexposed ponds. We therefore anchor the intercept to the satellite population. We choose it so that, in the simulated equilibrium, the share of plots without a pond neighbor that are in aquaculture matches the 2.9 percent observed in the satellite panel. Because fundamentals-driven ponds tend to induce their neighbors to convert, this observed share understates the rate implied by fundamentals alone, which the calibration puts at 5.4 percent. The intercept absorbs the conversion cost c. Finally, we recover δ, the net effect of a pond neighbor on the relative return to shrimp, by matching the model's implied change in conversion probability to the event-study estimate in Section 6 ($\Delta P = 0.180$). We estimate $\theta = 1.95$ and $\delta = 0.91$.

An important feature of δ, as discussed above, is that it is a net effect. A pond neighbor may lower the return to rice through salinity damage, raise the return to shrimp through shared access to saline water and infrastructure, or both. Land-use choices depend only on the difference, so choice data alone cannot decompose δ into these components. As we show below, the decomposition does not matter for the positive predictions of the model, but it determines the sign of the optimal policy.

7.2 Exercise 1: Cascades

We first use the estimated model to quantify how much aquaculture is driven by the externality rather than by fundamentals. We simulate a stylized landscape of

Table 2. Model Parameters and Sources of Identification

Parameter	Estimated from	What it captures
θ	Survey: spread of log output	Dispersion of comparative advantage
γ (slopes)	Survey: land-use logit	How plot size and distance to the river shift relative suitability
γ (level)	Satellite: share of plots without a pond neighbor in aquaculture (2.9%), matched in the simulated equilibrium	Baseline adoption; absorbs the conversion cost c
δ	Satellite panel: event-study ΔP	Net effect of a pond neighbor

plots on a grid, draw productivities from the estimated distribution, and solve for the equilibrium allocation by iterating plots' best responses to their neighbors' land use, starting from the allocation implied by fundamentals alone. Because the externality only lowers the threshold for conversion, every plot that converts on fundamentals converts in any equilibrium, so this procedure selects the equilibrium with the least aquaculture. This is also the allocation reached when land use starts in rice and plots respond to their neighbors' choices season by season.




Figure 8. Shrimp Cascade on a Stylized Economy

Figure 8 simulates land use on a stylized 60×60 grid of plots, each with four neighbors and with plot size and distance to the river drawn from the observed distributions. Each plot draws its productivity in rice and shrimp from the estimated Fréchet distribution, with dispersion θ and covariate effects γ estimated from the

survey, and the level set so that, in the simulated equilibrium, the share of plots without a pond neighbor that are ponds matches the 2.9 percent observed in the satellite panel. The left panel shows the allocation if every plot chose on fundamentals alone, ignoring its neighbors: 5.4% of plots become ponds. The middle panel adds the spillover: a converted neighbor shifts a plot's relative return to shrimp by δ, calibrated so the model reproduces the event-study effect of a neighbor's conversion, and choices are iterated to a fixed point. Aquaculture rises to 11.2%, and the orange plots are ponds that exist only because a neighbor converted first; they grow outward from the fundamentals-driven ponds in blue. The right panel repeats the exercise over 40 productivity draws and shows, for each plot, how often it is cascade-driven when it is a pond (grey plots are ponds in fewer than 15% of draws). The exercise separates aquaculture that farmers would choose on their own land's merits from aquaculture that spreads through the externality, and shows that the cascade roughly doubles conversion, with 52% of all ponds owing their existence to a neighbor.

7.3 Exercise 2: Optimal Policy

We next ask how a planner maximizing aggregate farm profit would intervene. The answer depends on a decomposition that land-use choices cannot reveal. The net effect of a pond neighbor combines two components,

$$\delta = \delta_R + \delta_S, \tag{9}$$

where δ_R is the damage that a neighboring pond imposes on rice and δ_S is the benefit it confers on a neighboring pond. Both components push an exposed plot toward aquaculture, so they are observationally equivalent in choice data, but they have opposite welfare implications. Damage to rice is a negative externality that calls for a tax on conversion, while a benefit to neighboring ponds is a positive externality that calls for a subsidy. The optimal policy is a tax when rice damage accounts for a large enough share of δ. In our calibration, this threshold corresponds to a rice yield loss of about 7% from a pond neighbor. At the other extreme, attributing all of δ to rice damage would imply a 60% yield loss.

We discipline the decomposition with two independent sources of evidence. Even at our satellite-based lower bound on the damage, the market over-converts, though the case for a tax rests on that damage exceeding about 7 percent of rice yields, which land-use choices alone cannot reveal. Farmers' own assessments, a 48 percent loss, place the economy well inside the tax region. Taking these beliefs as expert knowledge of local conditions, the optimal tax reduces the aquaculture share from 11.4 percent

to 0.1 percent and raises aggregate farm profit by about 11 percent, recovering 96 percent of the gain available to an unconstrained planner.

We view the satellite estimate as a lower bound on the true damage for two reasons. First, NDVI measures greenness rather than yield, and salinity may reduce grain yield more than vegetative growth. Second, the estimate is obtained from plots that do not convert, which are likely those least affected. The divergence between the two sources nonetheless shows that the welfare case for intervention, and even its sign, rests on the magnitude of damage to rice, which choice data alone cannot reveal.

8 Conclusion

This paper studies whether a private adaptation to climate change imposes costs on those who do not adopt it, and whether those costs accelerate adoption. In southwestern Bangladesh, where farmers can respond to rising salinity by converting rice paddies into saltwater shrimp ponds, we find evidence that they do. Rice plots bordering shrimp ponds are saltier and less productive than comparable plots, and farmers expect a neighbor's conversion to reduce their own rice yields substantially. Using a new plot-level panel of land use constructed from satellite imagery, we show that a neighbor's conversion lowers vegetation indices on adjacent agricultural plots and raises the probability that those plots convert by 18.0 percentage points. The effect appears immediately, is concentrated among plots that share direct borders with a new pond, and is difficult to reconcile with shared ownership or social learning. A model of land use with a boundary externality, estimated from the survey and the event study, implies that this response generates a substantial cascade: 51.5 percent of ponds in the simulated landscape exist only because a neighbor converted.

These results suggest that adaptation to climate change can be self-reinforcing in ways that are not captured by the usual focus on barriers to adoption. In many settings, frictions such as missing credit markets, risk, and limited information slow the take-up of new agricultural technologies. In ours, an externality on non-adopters works in the opposite direction, so that adoption spreads faster than comparative advantage alone would imply. Whether this is a problem depends on the size of the damage to rice. If the damage is as large as farmers believe, the market converts too much land to shrimp, and a tax on conversion would raise aggregate farm profit by about 10 percent. If the damage is closer to our satellite-based lower bound, the externality is dominated by benefits to neighboring ponds, and the case for intervention reverses. Because land-use choices alone cannot distinguish these cases, credible mea-

surement of the damage that one farmer's adaptation imposes on another is central to the design of policy.

More broadly, the setting illustrates a feature of adaptation that matters beyond coastal Bangladesh. In settings where adaptation to climate change will be decentralized, that is, carried out by households and firms responding to their own conditions rather than coordinated by the state, externalities can drive a wedge between private adaptation and the choice of the social planner. In these settings, evaluating adaptation by the gains to adopters alone will overstate its benefits, and the spread of a technology will be a poor guide to its social value. When rapid take-up can reflect the cost of remaining a non-user rather than the value of the technology itself, policy that treats fast adoption as evidence of success risks encouraging more of it than is socially desirable.

References

Adamopoulos, Tasso, Diego Restuccia, Julieta Caunedo, Kevin Donovan, Mushfiq Mobarak, Jacob Moscona, Ishan Nath, and Karthik Sastry, "Agricultural Policies for Growth," 2026.

Aker, Jenny C and B Kelsey Jack, "Harvesting the rain: The adoption of environmental technologies in the Sahel," *Review of Economics and Statistics*, 2025, *107* (5), 1197–1214.

Ayers, John C, Gregory George, David Fry, Laura Benneyworth, Carol Wilson, Leslie Wallace Auerbach, Kushal Roy, M R Karim, Farjana Akter, and Steven Goodbred, "Surface water chemistry and tidal channel time series in southwest Bangladesh," 2017. Supplement to: Ayers, JC et al. (2017): Salinization and arsenic contamination of surface water in southwest Bangladesh. Geochemical Transactions, 18, 1-23, https://doi.org/10.1186/s12932-017-0042-3.

Bangladesh Bureau of Statistics, "Agriculture Census 2019: Structure of Agricultural Holdings and Livestock & Fisheries," National Series Volume 1, Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic of Bangladesh, Dhaka December 2022.

Bari, Ehsanul, Md Arif Chowdhury, Md Ismail Hossain, and Mohammad Mahfuzur Rahman, "Using remote sensing data to study anthropogenic land degradation in Khulna Division, Bangladesh for SDG indicator 15.3. 1," *Heliyon*, 2024, *10* (19).

Beaman, Lori, Ariel BenYishay, Jeremy Magruder, and Ahmed Mushfiq Mobarak, "Can network theory-based targeting increase technology adoption?," *American Economic Review*, 2021, *111* (6), 1918–1943.

Blakeslee, David, Ram Fishman, and Veena Srinivasan, "Way down in the hole: Adaptation to long-term water loss in rural India," *American Economic Review*, 2020, *110* (1), 200–224.

Braaten, Robert O. and Mark Flaherty, "Salt balances of inland shrimp ponds in Thailand: implications for land and water salinization," *Environmental Conservation*, 2001, *28* (4), 357–367.

Brammer, Hugh, "Floods in Bangladesh: geographical background to the 1987 and 1988 floods," *Geographical journal*, 1990, pp. 12–22.

Burke, Marshall, Solomon M Hsiang, and Edward Miguel, "Global non-linear effect of temperature on economic production," *Nature*, 2015, *527* (7577), 235–239.

Carleton, Tamma, Amir Jina, Michael Delgado, Michael Greenstone, Trevor Houser, Solomon Hsiang, Andrew Hultgren, Robert E Kopp, Kelly E McCusker, Ishan Nath, James Rising, Ashwin Rode, Hee Kwon Seo, Arvid Viaene, Jiacan Yuan, and Alice Tianbo Zhang, "Valuing the Global Mortality Consequences of Climate Change Accounting for Adaptation Costs and Benefits," *The Quarterly Journal of Economics*, 11 2022, *137* (4), 2037–2105.

_ , **Esther Duflo, B. Kelsey Jack, and Guglielmo Zappalà**, "Chapter 4 - Adaptation to climate change," in Lint Barrage and Solomon Hsiang, eds., *Lint Barrage and Solomon Hsiang, eds.*, Vol. 1 of *Handbook of the Economics of Climate Change*, North-Holland, 2024, pp. 143–248.

Casaburi, Lorenzo and Jack Willis, "Time versus State in Insurance: Experimental Evidence from Contract Farming in Kenya," *American Economic Review*, 2018, *108* (12), 3778–3813.

Chen, Joyce and Valerie Mueller, "Coastal climate change, soil salinity and human migration in Bangladesh," *Nature Climate Change*, 2018, *8* (11), 981–985.

_ , _ , **Fabien Durand, Erika Lisco, Qing Zhong, V Raju Sherin, and AKM Saiful Islam**, "Salinization of the Bangladesh Delta worsens economic precarity," *Population and Environment*, 2022, *44* (3), 226–247.

Conley, Timothy G, "GMM estimation with cross sectional dependence," *Journal of econometrics*, 1999, *92* (1), 1–45.

_ **and Christopher R Udry**, "Learning about a new technology: Pineapple in Ghana," *American economic review*, 2010, *100* (1), 35–69.

Costinot, Arnaud, Dave Donaldson, and Cory Smith, "Evolving comparative advantage and the impact of climate change in agricultural markets: Evidence from 1.7 million fields around the world," *Journal of Political Economy*, 2016, *124* (1), 205–248.

Dasgupta, Susmita, Md Moqbul Hossain, Mainul Huq, and David Wheeler, "Climate change and soil salinity: The case of coastal Bangladesh," *Ambio*, 2015, *44* (8), 815–826.

de Chaisemartin, Clément and Xavier D'Haultfœuille, "Two-way fixed effects estimators with heterogeneous treatment effects," *American Economic Review*, 2020, *110* (9), 2964–2996.

Debsharma, Sanjoy K, M Akhlasur Rahman, Mahmuda Khatun, Ribed F Disha, Nusrat Jahan, Md Ruhul Quddus, Hasina Khatun, Sharifa S Dipti, Md Ibrahim, KM Iftekharuddaula et al., "Developing climate-resilient rice varieties (BRRI dhan97 and BRRI dhan99) suitable for salt-stress environments in Bangladesh," *Plos one*, 2024, *19* (1), e0294573.

Dell, Melissa, Benjamin F. Jones, and Benjamin A. Olken, "Temperature Shocks and Economic Growth: Evidence from the Last Half Century," *American Economic Journal: Macroeconomics*, July 2012, *4* (3), 66–95.

Department of Fisheries, "Yearbook of Fisheries Statistics of Bangladesh, 2024–25," Technical Report, Fisheries Resources Survey System (FRSS), Department of Fisheries, Ministry of Fisheries and Livestock, Government of the People's Republic of Bangladesh, Dhaka December 2025.

Duflo, Esther, Michael Kremer, and Jonathan Robinson, "Nudging farmers to use fertilizer: Theory and experimental evidence from Kenya," *American economic review*, 2011, *101* (6), 2350–2390.

Emerick, Kyle, Alain De Janvry, Elisabeth Sadoulet, and Manzoor H Dar, "Technological innovations, downside risk, and the modernization of agriculture," *American Economic Review*, 2016, *106* (6), 1537–1561.

Esha, Afsana Afrin, C Isabella Bovolo, Hanna A Ruszczyk, and Andrew Baldwin, ""There Is No Governance": Drinking Water Governance in Coastal Areas of Bangladesh," *Water*, 2026, *18* (7), 861.

Farrokhi, Farid and Heitor S Pellegrina, "Trade, technology, and agricultural productivity," *Journal of Political Economy*, 2023, *131* (9), 2509–2555.

Foster, Andrew D and Mark R Rosenzweig, "Learning by doing and learning from others: Human capital and technical change in agriculture," *Journal of political Economy*, 1995, *103* (6), 1176–1209.

Goodman-Bacon, Andrew, "Difference-in-differences with variation in treatment timing," *Journal of Econometrics*, 2021, *225* (2), 254–277.

Grattan, Stephen R, Linghe Zeng, Michael C Shannon, and Stacy R Roberts, "Rice is more sensitive to salinity than previously thought," *California agriculture*, 2002, *56* (6).

Griliches, Zvi, "Hybrid Corn: An Exploration in the Economics of Technological Change," *Econometrica*, 1957, *25* (4), 501–522.

Guimbeau, Amanda, Xinde James Ji, Zi Long, and Nidhiya Menon, "Ocean salinity, early-life health, and adaptation," *Journal of environmental economics and management*, 2024, *125*, 102954.

Hanna, Rema, Sendhil Mullainathan, and Joshua Schwartzstein, "Learning through noticing: Theory and evidence from a field experiment," *The Quarterly Journal of Economics*, 2014, *129* (3), 1311–1353.

Haque, Mohammad Islamul, "The role of land market in achieving the scale for adaptation to climate change: Evidence from Bangladesh," *Available at SSRN 5245815*, 2025.

IPCC, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge, UK and New York, NY, USA: Cambridge University Press, 2021.

_ , *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge, UK and New York, NY, USA: Cambridge University Press, 2022.

Islam, M. S., M. A. Islam, M. C. Rahman, A. Chowdhury, M. S. Rahaman, L. Deb, S. M. M. H. Noman, and S. A. Jui, "Farm Level Adoption and Evaluation of Modern Rice Cultivation in Bangladesh," Annual Research Review Workshop 2022–2023, Agricultural Economics Division, Bangladesh Rice Research Institute, Gazipur 2024. Varietal adoption data from the Department of Agricultural Extension, 2022–23.

Islam, M Serajul, MM Haque, MG Rabbani, and S Sharmin, "Marketing of shrimp in Bangladesh-A value chain analysis," *Journal of the Bangladesh Agricultural University*, 2014, *12* (2), 359–368.

Jamal, Md Roushon, Paul Kristiansen, Md Jahangir Kabir, and Lisa Lobry de Bruyn, "Risks and adaptation dynamics in shrimp and prawn-based farming systems in southwest coastal Bangladesh," *Aquaculture*, 2023, *562*, 738819.

Jones, Maria, Florence Kondylis, John Loeser, and Jeremy Magruder, "Factor market failures and the adoption of irrigation in Rwanda," *American Economic Review*, 2022, *112* (7), 2316–2352.

Kabir, Razin Iqbal, Sudha Narayanan, Ben Belton, Ricardo Hernandez, and Mohammad Mahfujul Haque, *Shrimp value chains in Bangladesh: A scoping study of possible research interventions*, Intl Food Policy Res Inst, 2023.

Kala, Namrata, Clare Balboni, and Eddy Zou, "Climate Adaptation," *VoxDevLit*, May 2026, *7* (2).

Karlan, Dean, Robert Osei, Isaac Osei-Akoto, and Christopher Udry, "Agricultural Decisions after Relaxing Credit and Risk Constraints," *The Quarterly Journal of Economics*, 2014, *129* (2), 597–652.

Lavreniuk, Mykola, Nataliia Kussul, Andrii Shelestov, Bohdan Yailymov, Yevhenii Salii, Volodymyr Kuzin, and Zoltan Szantoi, "Delineate Anything: Resolution-Agnostic Field Boundary Delineation on Satellite Imagery," *arXiv preprint arXiv:2504.02534*, 2025.

Maas, Eugene V and Glenn J Hoffman, "Crop salt tolerance—current assessment," *Journal of the irrigation and drainage division*, 1977, *103* (2), 115–134.

Magruder, Jeremy R, "An assessment of experimental evidence on agricultural technology adoption in developing countries," *Annual Review of Resource Economics*, 2018, *10* (1), 299–316.

McGuirk, Eoin F and Nathan Nunn, "Transhumant pastoralism, climate change, and conflict in Africa," *Review of Economic Studies*, 2025, *92* (1), 404–441.

Ministry of Environment, Forest and Climate Change, Government of the People's Republic of Bangladesh, "Third National Communication of Bangladesh to the United Nations Framework Convention on Climate Change," Technical Report, Ministry of Environment, Forest and Climate Change, Government of the People's Republic of Bangladesh, Dhaka, Bangladesh June 2018.

Mobarak, Ahmed Mushfiq and Mark R. Rosenzweig, "Informal Risk Sharing, Index Insurance, and Risk Taking in Developing Countries," *American Economic Review*, 2013, *103* (3), 375–380.

Mondal, S, MM Rahman, D Saha, RK Adhikary, and M Belal Hossain, "Present status of good aquaculture practices (GAP) in shrimp farms of southwestern coastal area, Bangladesh," *Middle-East Journal of Scientific Research*, 2013, *14* (6), 873–878.

Morshed, Md. Manjur, Md. Sariful Islam, Heman Das Lohano, and Priya Shyamsundar, "Production externalities of shrimp aquaculture on paddy farming in coastal Bangladesh," *Agricultural Water Management*, 2020, *238*, 106213.

Nurullah, SM, Md Roushon Jamal, and Md Rakibul Islam, "Bangladesh's dying shrimp industry crying for policy action," 2024.

OpenStreetMap, "OpenStreetMap," https://www.openstreetmap.org 2026. Data retrieved September 2026.

Patel, Dev, "Environmental Beliefs and Adaptation to Climate Change," *Unpublished manuscript*, 2025.

Paul, Priya Lal Chandra, Afsana Jahan, Palash Kumar Kundu, Debjit Roy, Richard W Bell, Md Belal Hossain, Rakiba Shultana, Mohammad Rezoan Bin Hafiz Pranto, Tanjina Islam, Sharon E Benes et al., "Rice growth and yield responses to saline water irrigation are related to Na+/K+ ratio in plants," *PLoS One*, 2024, *19* (11), e0312372.

Rahi, Md Lifat, Khairun Naher Azad, Maliha Tabassum, Hasna Hena Irin, Kazi Sabbir Hossain, Dania Aziz, Azam Moshtaghi, and David A Hurwood, "Effects of salinity on physiological, biochemical and gene expression parameters of black tiger shrimp (Penaeus monodon): Potential for farming in low-salinity environments," *Biology*, 2021, *10* (12), 1220.

Rahman, Ahmed Z, Mohammad Shamsudduha, Md Izazul Haq, Md Hanif, Md Sanaul Islam, Syed S Islam, Amarendranath Biswas, and Richard G Taylor, "Soil and river water salinity dynamics in coastal Bangladesh," *Scientific Reports*, 2025, *16* (1), 1044.

Roy, Andrew Donald, "Some thoughts on the distribution of earnings," *Oxford economic papers*, 1951, *3* (2), 135–146.

Ryan, Nicholas and Anant Sudarshan, "Rationing the commons," *Journal of Political Economy*, 2022, *130* (1), 210–257.

Soil Resource Development Institute, "Saline Soils of Bangladesh," Technical Report, Soil Resource Development Institute (SRDI), SRMAF Project, Ministry of Agriculture, Dhaka, Bangladesh November 2010.

Sotelo, Sebastian, "Domestic trade frictions and agriculture," *Journal of Political Economy*, 2020, *128* (7), 2690–2738.

Stein, Shaked, Eyal Shalev, Orit Sivan, and Yoseph Yechieli, "Challenges and approaches for management of seawater intrusion in coastal aquifers," *Hydrogeology Journal*, 2023, *31* (1), 19–22.

Sun, Liyang and Sarah Abraham, "Estimating dynamic treatment effects in event studies with heterogeneous treatment effects," *Journal of Econometrics*, 2021, *225* (2), 175–199.

Suri, Tavneet, "Selection and comparative advantage in technology adoption," *Econometrica*, 2011, *79* (1), 159–209.

_ and Christopher Udry, "Agricultural technology in Africa," *Journal of Economic Perspectives*, 2022, *36* (1), 33–56.

UNESCO, IWG, "The practical salinity scale1978 and the international equation of state of seawater 1980," *Tenth Report of the Joint Panel on Oceanographic Tables and Standards (JPOTS)*, 1981, *25*.

World Bank Group, "Climate Risk Profile: Bangladesh," Climate Risk Profile, The World Bank Group, Washington, DC 2024.

Xu, Hanqiu, "Modification of normalised difference water index (NDWI) to enhance open water features in remotely sensed imagery," *International journal of remote sensing*, 2006, *27* (14), 3025–3033.

Yu, Xuan, Beiyuan Xu, Rongjiang Yao, Junhong Wei, Tongbi Tu, and Zi Chen, "Temporal dynamics of soil salinization due to vertical and lateral saltwater intrusion at an onshore aquaculture farm," *Agricultural Water Management*, 2024, *306*, 109179.

A Appendix 1 - Additional Tables

Appendix Table B1. Delineation Coverage by Union

Union	Upazila	Area (km^2)	Area covered (km^2)	Share covered (%)	Plots
Anulia	Assasuni	28.82	10.54	36.6	1,202
Assasuni	Assasuni	34.86	13.64	39.1	1,528
Baradal	Assasuni	36.15	21.16	58.5	2,786
Budhhata	Assasuni	33.69	9.47	28.1	1,461
Durgapur	Assasuni	24.24	11.47	47.3	1,712
Kadakati	Assasuni	28.24	17.99	63.7	2,869
Khajra	Assasuni	37.29	18.80	50.4	1,934
Kulla	Assasuni	32.61	8.41	25.8	1,121
Sreeula	Assasuni	43.66	28.54	65.4	3,892
Atlia	Dumuria	42.01	13.52	32.2	1,506
Dumuria	Dumuria	18.68	1.34	7.2	238
Kharnia	Dumuria	21.21	2.21	10.4	509
Maguraghona	Dumuria	21.18	4.51	21.3	834
Magurkhali	Dumuria	41.66	22.75	54.6	3,359
Raghunathpur	Dumuria	34.88	12.45	35.7	1,852
Rudaghara	Dumuria	28.66	9.50	33.1	1,736
Sahas	Dumuria	25.07	7.90	31.5	1,406
Sobhana	Dumuria	45.45	7.48	16.5	918
Champaphul	Kaliganj	29.87	15.69	52.5	1,912
Krishnanagar	Kaliganj	26.22	10.10	38.5	1,800
Bidyanandakati	Keshabpur	32.76	5.69	17.4	1,051
Gaurighona	Keshabpur	25.82	4.83	18.7	1,070
Majidpur	Keshabpur	24.39	3.03	12.4	746
Mangalkot	Keshabpur	23.91	3.00	12.6	455
Sagardari	Keshabpur	28.79	3.71	12.9	1,068
Amadi	Koyra	41.87	19.95	47.6	2,744
Bagali	Koyra	48.32	23.05	47.7	2,157
Maheshwaripur	Koyra	52.03	14.35	27.6	771
Chandkhali	Paikgachha	45.67	23.52	51.5	3,657
Deluti	Paikgachha	46.09	24.72	53.6	1,433
Gadaipur	Paikgachha	20.39	8.95	43.9	860
Garuikhali	Paikgachha	42.86	28.10	65.6	1,352
Haridhali	Paikgachha	19.74	3.28	16.6	508
Kapilmuni	Paikgachha	38.01	16.92	44.5	2,233
Laskar	Paikgachha	42.16	24.92	59.1	2,516
Lata	Paikgachha	48.24	24.67	51.1	2,004
Paikgachha Paurashava	Paikgachha	4.81	0.73	15.2	56
Raruli	Paikgachha	25.36	7.74	30.5	922
Sholadana	Paikgachha	49.32	27.75	56.3	1,867
Dhulihar	Satkhira Sadar	37.95	16.68	43.9	2,078
Kashimari	Shyamnagar	34.40	16.38	47.6	1,378
Islamkati	Tala	22.18	3.38	15.2	819
Jalalpur	Tala	24.18	3.39	14.0	640
Khalilnagar	Tala	32.55	9.24	28.4	1,799
Khalishkhali	Tala	37.98	13.45	35.4	1,773
Khesra	Tala	48.02	10.83	22.6	1,066
Kumira	Tala	19.42	2.94	15.1	858
Magura	Tala	26.10	3.36	12.9	926
Sarulia	Tala	27.17	5.18	19.1	1,420
Tala	Tala	24.77	4.05	16.4	794
Tentulia	Tala	26.09	4.90	18.8	786
Total		1,655.82	610.15	36.8	76,382


Notes: The table reports the area of each union falling inside a delineated plot, and the share of the union covered. Union boundaries are from the Bangladesh Bureau of Statistics administrative boundaries. Plots are delineated from basemap imagery using Delineate Anything model described in Appendix C. Covered area is computed by clipping plots to union boundaries and merging overlapping polygons, so no land is counted twice. Plots are assigned to unions by the location of their centroid. The largest 1% of plots are included in this table, but excluded prior to analysis.

Appendix Table B2. Reasons for Shrimp Farming

	Respondents	Share (%)	Sole reason (%)
Profits are better in shrimp farming	263	79.9	39.8
Rice yields went down	123	37.4	10.0
Someone in the village recommended it	60	18.2	4.3
NGO or government advised it	5	1.5	0.0
Other	40	12.2	3.3
Shrimp-farming households	394		
Answered why they converted	329		
Gave more than one reason	140	42.6	


Notes: The table shows respondents answers to the question of why they first adopted shrimp farming. Multiple answers are permitted.

B Appendix 2 - Additional Figures




Appendix Figure B1. Sample Location

Notes: The Unions sampled in the household survey of rice and shrimp farmers across the Satkhira and Khulna Districts are shown in dark red. The satellite analysis covers unions with centroids that are southwest of the northeastern most survey-sampled union and northeast of the southwestern most survey-sampled union.




Appendix Figure B2. Respondent Beliefs About Prices

Notes: Empirical cumulative distribution functions of respondents' expected output prices, by household shrimp status. A household is classified as shrimp farming if at least one of its plots is a shrimp pond, and as non-shrimp farming otherwise. Rice is Boro-season rice; shrimp is bagda (tiger prawn). Only positive price beliefs are included. Beliefs are heaped at round numbers, which produces the steps in each curve. Each panel reports the difference in mean beliefs (shrimp minus non-shrimp farming) with a 95% confidence interval based on standard errors clustered by household.




Appendix Figure B3. Salinity Test Location Diagram

Note: The diagram above shows an example of three plots, the leftmost of which is flooded with salt water for shrimp cultivation and the remaining two are used for agriculture. The red dots illustrate the approximate locations of the salinity readings.




Appendix Figure B4. Rice Health Scale

Notes: The photograph is reproduced from Grattan et al. (2002) and shows rice plants grown at salinity levels increasing from 0.3 dS/m (left) to 12.0 dS/m (right), with plant growth visibly stunted at higher salinity. Enumerators showed this image to respondents when eliciting subjective rice health in Figure 3, so that a respondent's rating could be anchored to a common visual reference.




Appendix Figure B5. Modified Normalized Difference Water Indices Distribution

Notes: Distribution of the plot-level Modified Normalized Difference Water Index (MNDWI; Xu 2006) from Sentinel-2 imagery in the Rabi season (November–February), for plots with at least three valid pixels. The dashed line is that year's classification threshold, chosen by Otsu's method to maximize the between-class variance of the index; plots above it are classified as aquaculture, and the panel title reports how many.




Appendix Figure B6. Share of Adjacent Pond Pairs with Near-Identical Water-Index Readings

Note: Same-season converter pairs are a treated plot that switches from rice to aquaculture in the same Rabi season as its first pond neighbor, paired with that neighbor; the benchmark is a random sample of other adjacent pairs in which both plots are ponds at some point in the panel. Bars show the share of each group whose two plots are ponds in the same years (overlap of at least 90%) and, in the years when both are ponds, have a mean absolute MNDWI difference and a correlation of MNDWI readings within the stated thresholds; only pairs that are ponds together for at least four years are included.




Appendix Figure B7. Robustness: Event Study with Manual Pond Threshold

Note: The figure compares Sun and Abraham (2021) event-study estimates of own aquaculture status under the main classification, which uses a year-specific Otsu threshold, with the same specification after reclassifying every plot-year using a fixed threshold of MNDWI > 0.241, chosen to minimize the balanced error rate against survey-reported land use. Both specifications include plot and hexagon $\times$ year fixed effects and river-distance $\times$ year and plot-size $\times$ year controls; bars show 95% confidence intervals using Conley standard errors with a 2 km cutoff, and the legend reports each post-period ATT.




Appendix Figure B8. Heterogeneity by plot size

Note: The figure overlays Sun and Abraham (2021) event-study estimates of own aquaculture status for plots below and above the median plot area in the estimation sample (3,273 $\mathbf{m^2}$), each estimated separately with the main specification. Both include plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls; bars show 95% confidence intervals using Conley standard errors with a 2 km cutoff, and the legend reports each group's post-period ATT.




Appendix Figure B9. Heterogeneity by distance to river

Note: The figure overlays Sun and Abraham (2021) event-study estimates of own aquaculture status for plots below and above the median distance to the nearest river, each estimated separately with the main specification. Distance is measured from the plot boundary to the nearest OpenStreetMap river segment. Both include plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls; bars show 95% confidence intervals using Conley standard errors with a 2 km cutoff, and the legend reports each group's post-period ATT.




Appendix Figure B10. Robustness: Instrumental Variables Approach

Note: The top row reports the post-period ATT from the main Sun and Abraham (2021) event study of own aquaculture status. The bottom row reports a 2SLS estimate in which an indicator for having had an adjacent pond neighbor is instrumented by the share of a plot's second-degree neighbors (neighbors of its neighbors that do not border it directly) that were ponds in the previous year. Both specifications include plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls; bars show 95% confidence intervals using Conley standard errors with a 2 km cutoff, and the first-stage F-statistic is reported for the IV.



Appendix Figure B11. Treatment Effects by Plot Rings

Notes: Each panel is a separate Sun and Abraham (2021) event study of own aquaculture status. Treatment is the first year a pond appears in that distance ring, measured edge to edge and excluding adjacent plots for the outer rings, and plot-years are censored once a closer ring is exposed. All specifications include plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls. Shaded bands show 95% (dark) and 99% (light) confidence intervals using Conley standard errors with a 2 km cutoff.



Appendix Figure B12. Robustness: Excluding Potentially Split Ponds

Note: The figure reports the post-period ATT from the main Sun and Abraham (2021) event study of own aquaculture status, and the same estimate after excluding treated plots whose same-season pair meets the strictest and strict similarity criteria shown in Appendix Figure B6. All specifications include plot and hexagon × year fixed effects and river-distance × year and plot-size × year controls; bars show 95% confidence intervals using Conley standard errors with a 2 km cutoff.

C Appendix 3 - Delineate Anything Field-Boundary Algorithm

This appendix describes the procedure we use to convert raw satellite imagery into the individual plot polygons that serve as the units of analysis in the main text. We use *Delineate Anything*, an off-the-shelf, pre-trained machine-learning model for field-boundary delineation (Lavreniuk et al., 2025).

The goal is to take an aerial image of farmland and output a separate, closed polygon for each individual field. The engine is a neural network built on the YOLOv11 segmentation architecture, a widely used family of models designed to locate multiple objects in an image quickly and simultaneously (Lavreniuk et al., 2025). Rather than classifying loose pixels, the model is asked to identify each field as a distinct object and return that object's own filled outline (its "mask"). Because each field is detected as a whole entity, a small imperfection along one edge no longer merges it with its neighbor, which is the property we need for accurate plot-level measurement. Delineate Anything was trained on the Field Boundary Instance Segmentation dataset (FBIS-22M), which contains roughly 673,000 satellite image patches and about 22.9 million hand-delineated field outlines. These training images span a wide range of spatial resolutions (from 0.25 to 10 meters per pixel), multiple satellite sensors, and several countries and seasons. Having learned from this deliberately heterogeneous set of examples, the model is "resolution-agnostic," that is, it recognizes the general visual signature of a field—a roughly convex patch of uniform texture bounded by abrupt changes in color or tone—rather than memorizing the look of one specific sensor. This is what allows us to apply it to our imagery without retraining.

We apply the pre-trained model directly to our images without any further training or hand-labeling of Bangladeshi fields. The model requires only the three visible color bands (red, green, and blue) as input, which our Planet basemap imagery provides. This is an advantage for transparency and replicability: our plot boundaries are the output of a fixed, publicly documented model applied to public imagery, with no discretionary labeling choices on our part.

Because satellite imagery tiles overlap, the same field is sometimes detected more than once in adjacent tiles. A merging step reconciles these duplicates: when two detected outlines overlap by more than a set fraction of their area, they are treated as the same field and combined into one. This uses the standard notion of *intersection-over-union*—the ratio of the area two shapes share to the total area they jointly cover—as the measure of overlap, with a high threshold so that only near-identical detections are merged. After merging, the model's pixel masks are converted from filled raster blobs into clean vector polygons with explicit vertices, the format used by geographic information systems. Two size filters remove implausible detections: outlines smaller than a minimum ground area, and small spurious holes inside a field, are discarded.

Three caveats are worth noting for interpretation. First, the model can occasionally merge two truly distinct fields or split one field in two; because our analysis is at

the plot level, such errors introduce measurement noise into plot boundaries. Second, fields at the outer edge of a quad can be clipped where two quads meet, which our cross-quad merging step mitigates but does not perfectly eliminate. Third, the boundaries are fixed from a single basemap and treated as static over the panel; genuine changes in field boundaries over time (for example, the subdivision or consolidation of plots) are not captured, which may lead to measurement error.

Appendix Table B3. Delineate Anything Parameters

Parameter	Value	Default	Description
Detection			
Minimum confidence	0.005	0.005	Model confidence required to keep a detection
Minimum detection size	256 px	512 px	Smallest detected object retained, in pixels
Super-resolution	Auto	Auto	Optional upscaling before inference
Tiling and merging			
Tile step	0.5	0.5	Overlap between adjacent tiles (50%)
Merge IoU threshold	0.8	0.8	Overlap above which detections are merged
Tile composition IoU	0.8	0.8	Overlap above which detections across tiles are merged
Filtering			
Minimum field area	200 m^2	2,500 m^2	Smallest field retained
Minimum hole area	200 m^2	2,500 m^2	Smallest interior hole retained
Input and output			
Band order	[3, 2, 1]	[3, 2, 1]	Order in which image bands are passed to the model
Polygon simplification	Off	On	Simplification of output polygon vertices

Notes: Parameters used to run Delineate Anything (Lavreniuk et al., 2025) on Planet basemap quads. "Default" refers to the value in the model's sample configuration file. We lower the minimum detection size and minimum field area relative to the defaults because fields in southwestern Bangladesh are small; 200 m^2 corresponds to 0.02 hectares (about five decimals). IoU denotes intersection-over-union. Polygon simplification is disabled for computational stability. Computational settings (batch size of 16 and half-precision inference) do not affect the output and are omitted.

D Appendix 4 - Salinity Sensor Protocol

Section D.1 applies to the main plot, which was selected as a marginal shrimp or marginal rice plot and used to construct the sample as shown in Figure B3. Section D.2 applies to one randomly selected additional plot from each household's plot roster for additional salinity testing. Enumerators were provided with handheld electrical conductivity salinity meters with a range 0∼10.00ms/cm, resolution 0.01ms/cm and Accuracy ±2%F.S. Each meter was calibrated prior to the survey using the standard calibration solution.

D.1 Main Plots

After completing the interview with the respondent, please return to the main plot and follow the steps below for aquaculture/agriculture boundaries. For agriculture/agriculture boundaries, begin at step 3 and repeat steps 3 and 4 twice instead of beginning with step 1. Rinse the sensor with bottled water and dry with a clean cloth after each reading.

1. Measure the salinity of the water at the boundary of the aquaculture plot. Submerge the entire probe of the salinity sensor. Photograph the sensor display and record the reading on the tablet.
2. Take a second reading at the same boundary, photograph the display, and record the reading on the tablet.
3. Return to the border between the aquaculture and agricultural plots and walk five paces into the agricultural plot. Insert the sensor into the soil, photograph the display, and record the reading on the tablet. Wipe the sensor clean.
4. Move five paces to the left, parallel to the border, and repeat step 4, for a total of three soil readings.
5. Record an estimate of the width of the central agricultural plot.
6. Record whether it was raining at any time during the interview.

D.2 Other Plots from the Plot Roster

For each additional agricultural plot listed in the plot roster, the enumerator follows the steps below. Rinse the sensor with bottled water and dry with a clean cloth after each reading.

1. Walk five paces into the agricultural plot. Insert the sensor into the soil, photograph the display, and record the reading on the tablet.
2. Move five paces to the left and repeat step 1 for a total of two readings.