

This example solves the distributed minimization problem

$$\begin{aligned} \min J(q, u) &= \frac{1}{2} \|u - u^d\|^2 + \frac{\alpha}{2} \|q\|^2 \\ \text{s.t. } (\nabla u, \nabla \phi) &= (f(q), \phi) \quad \forall \phi \in H_0^1(\Omega; \mathbb{R}^2) \end{aligned}$$

on the domain $\Omega = [0, 1]^2$, and the data is chosen as follows:

$$\begin{aligned} f(q) &= q_0 \begin{pmatrix} 2\pi^2 \sin(\pi x) \sin(\pi y) \\ 0 \end{pmatrix} \\ &+ q_1 \begin{pmatrix} 5\pi^2 \sin(\pi x) \sin(2\pi y) \\ 0 \end{pmatrix} \\ &+ q_2 \begin{pmatrix} 0 \\ 8\pi^2 \sin(2\pi x) \sin(2\pi y) \end{pmatrix} \end{aligned}$$

with $\alpha = 0$ and u^d such that the solution is given by:

$$\begin{aligned} \bar{q} &= (1; 0.5; 1) \\ \bar{u} &= \begin{pmatrix} \sin(\pi x)(\sin(\pi y) + 0.5 \sin(2\pi y)) \\ \sin(2\pi x) \sin(2\pi y) \end{pmatrix} \end{aligned}$$