


# Gli1 → Kdm6b/Jmjd3 eraser of H3K27me3 at *Ccnd1* — detailed wiring + equations (v44.1)

mitogen-gated ERASER (Gli1 → KDM6B/JMJD3 demethylase) vs cycle-gated WRITER (EZH2/PRC2) — the per-division race that sets the mark



## ① Gli1 axis (mitogen input, with slow autoregulatory memory)

$$\begin{aligned} \text{Gli\_rep} &\rightleftharpoons \text{Gli\_act} && \text{(Smo-gated activator/repressor switch)} \\ d[\text{Gli1\_mRNA}]/dt &= V_{\text{max}} \cdot \text{Gli\_act}^2 / (K_{\text{GA}}^2 + \text{Gli\_act}^2) \cdot (1 - \text{Gli\_rep}^2 / (K_{\text{GR}}^2 + \text{Gli\_rep}^2)) \cdot \gamma_{\text{Ptch}} \\ &\quad + k_{\text{Gli1\_auto}} \cdot \text{Gli1\_epi} - k_{\text{mdeg}} \cdot \text{Gli1\_mRNA} \\ d[\text{Gli1\_epi}]/dt &= k_{\text{epi\_on}} \cdot (1 - \text{Ptch}) \cdot \text{Smo}^4 / (K_{\text{auto}}^4 + \text{Smo}^4) \cdot (1 - \text{Gli1\_epi}) - k_{\text{epi\_off}} \cdot \text{Gli1\_epi} \\ d[\text{Gli1}]/dt &= k_{\text{Gli1\_tl}} \cdot \text{Gli1\_mRNA} - k_{\text{Gli1\_deg}} \cdot \text{Gli1} \end{aligned}$$

## ② Eraser (KDM6B/JMJD3, implicit)

KDM6B is NOT a state variable.  
Quasi-static:  $[\text{KDM6B}] \propto \text{Gli1}$ ,  
folded into  $k_{\text{jmd3\_gli}}$ .

$$\begin{aligned} \text{erase flux} &= \\ k_{\text{jmd3\_gli}} \cdot \text{Gli1} \cdot \text{Mk} \end{aligned}$$

## ③ H3K27me3 (Mk) balance at the *Ccnd1* domain

$$\begin{aligned} d[\text{Mk}]/dt &= \underbrace{\text{PRC2} \cdot (1 - \text{Mk})}_{\text{(EZH2/PRC2, cycle)}} - \underbrace{\text{del\_mk} \cdot \text{Mk}}_{\text{(Gli1/KDM6B, mitogen)}} - \underbrace{k_{\text{jmd3\_gli}} \cdot \text{Gli1} \cdot \text{Mk}}_{\text{ERASE}} \\ \text{discrete: at Dna} &> 0.05 \rightarrow \text{Mk} \leftarrow \frac{1}{2} \cdot \text{Mk} && \text{(replicative dilution, once per S)} \\ \text{PRC2} &= \text{EZH2} \cdot (1 - \text{EZH2i}) \cdot (a_0 + a_{\text{rw}} \cdot \text{Mk}) \cdot (1 - g \cdot \text{Cd\_mRNA}^p / (K_{\text{tx}}^p + \text{Cd\_mRNA}^p)) \end{aligned}$$

## ④ Baked parameters (v44.1) + net effect

$$\begin{aligned} k_{\text{jmd3\_gli}} &0.0478 & \text{del\_mk} &0.0015 \\ a_0 &3.78\text{e-}4 & a_{\text{rw}} &4.30\text{e-}3 & (a_{\text{rw}}/a_0 &11.4) \\ g_{\text{prc2}} &0.0424 & K_{\text{prc2}} &3.5\text{e-}3 & n &3.39 \\ f_0_{\text{prc2}} &0.0505 & K_{\text{tx}} &5.12 & p_{\text{tx}} &3.27 \\ \text{Gli: } V_{\text{max}} &1.27 & \text{auto} &0.012 & \text{on/off} &0.02/8\text{e-}4 \end{aligned}$$

**MB Gli1 ≈ 6.9× GNP → erase ≈ 6.9× stronger → ChIP *Ccnd1* mark MB/GNP ≈ 0.55 (MB < GNP)**

Net over divisions: *Mk* accumulates when  $\text{WRITE} > (\text{turnover} + \text{ERASE} + \frac{1}{2}\text{-dilution})$  each cycle; MB's high *Gli1* tilts the race toward erasure despite higher *EZH2*.