

v44 model — experimental conditions: input parameters & where they sit in the proliferation parameter space


A Input parameters per condition

	SHH	Ptch1	f MYCN	p16	HHi	EZH2i	HU	CDK4/6i	yc?
GNP +SHH	0.5	1	1	0	.	.	.	.	●
GNP +HHi	0.5	1	1	0	✓	.	.	.	○
GNP +EZH2i	0.5	1	1	0	.	✓	.	.	●
GNP serum-starve	0.5	1	1	0	.	.	.	.	○
Ptch+/- (P7)	0.5	0.5	1.38	1	.	.	.	.	●
MB	0.5	0.1	2.86	3.3	.	.	.	.	●
MB +HHi	0.5	0.1	2.86	3.3	✓	.	.	.	○
MB +HHi +EZH2i	0.5	0.1	2.86	3.3	✓	✓	.	.	●
MB +CDK4/6i	0.5	0.1	2.86	3.3	.	.	.	✓	○
MB +HU	0.5	0.1	2.86	3.3	.	.	✓	.	●

serum-starve = k_Cd_transl→0; CDK4/6i = context input

drug switches

B Parameter space: CyclinD1 vs p16 commitment threshold



tumour CyclinD1/p27 heterogeneity straddles the boundary → fractional response (MB+HHi 22% → +EZH2i 69% cycling)