

Expressed
n=15575
Unexpressed
n=4987
Granule neuron lineages
n=12704
Cell cycle
n=997
Alternative lineage
n=1874
excluded
n=4987
Dynamically expressed
n=7334
unchanged
n=5370
excluded
n=7858
P14 to P56
n=2675
P0 to P7
n=1750
E15
n=2784
incompatible max−min
n=125
unchanged
n=5370
excluded
n=7858
P14 to P56
n=2050
P0 to P7
n=1434
E15
n=2294
p−val inadequate
n=1431
unchanged
n=5370
excluded
n=7983
P14 to P56
n=1877
P0 to P7
n=1434
E15
n=2294
increase before P0
n=173
unchanged
n=6801
excluded
n=7983
>2 log2FC
n=787
1−2 log2FC
n=1090
>2 log2FC
n=440
1−2 log2FC
n=994
>2 log2FC
n=783
1−2 log2FC
n=1511
excluded
n=14957
differentiation genes
n=787
proliferation genes
n=440
embryonic genes
n=783
Step 1:
Remove genes
unexpressed during
the time course
Step 2:
Remove genes
contaminated
from other cell lines
Taylor scRNAseq
Step 3:
Remove genes that
change less
then 2 fold
Step 4:
Split changing genes
based on when they
arrive at max
Step 5:
Remove genes with
adjusted p−val > 0.001
in differential expression
from RNAseq or NeuroD1−TRAP
Step 6:
Remove differentiation
genes that
increase most
prior to P0
Step 7:
Seperate
by fold change
Step 8:
Final groups