

ASTR 421

Stellar Observations and Theory

Lecture 09

Line Profiles: II

Prof. James Davenport (UW)

1



This area reserved for webcam overlay

Today

- Details of line broadening mechanisms
- Ca II HK
- Revisiting other line profiles (P Cygni)



This area reserved for webcam overlay


Overview of Line Broadening

Many things can impact “doppler broadening”, anything w/ gas velocities

- Doppler broadening (thermal motions)
- Natural broadening (Heisenberg uncertainty)
- Pressure broadening (a few kinds)

Gaussian profile


Lorentzian profile



The Voigt Profile


- A *convolution* of Gaussian & Lorentzian profiles
- This the classic line profile equation we use in most cases
 - Core: mostly Gaussian shape
 - Wings: mostly Lorentz shape
- **Useful for constraining physics behind both thermal & pressure broadening in a line!**

$$V(x; \sigma, \gamma) = \int_{-\infty}^{\infty} G(x'; \sigma) L(x - x'; \gamma) dx'$$



This area reserved for webcam overlay

Natural broadening





- Arises due to Heisenberg uncertainty principle
 $\Delta E \Delta t \approx \hbar$
- Time electron spends in excited state is finite... the photon *does* get emitted!
- Plugging in the classic $E = hc/\lambda$ for photon...
- BOB gives estimate of the line width for natural broadening of:
$$FWHM = \frac{\lambda^2}{\pi c \Delta t} \quad (\text{eqn 9.61})$$
- Example for $H\alpha$: $FWHM \sim 2 \times 10^{-5} \text{ nm (!)}$
 - **TYPICALLY VERY SMALL**



Natural broadening

- Arises due to Heisenberg uncertainty principle
 $\Delta E \Delta t \approx \hbar$
- Another way to think about this is the *location* of the energy levels cannot be precisely determined at any given moment.




Gray, Fig 11.1



This area reserved for webcam overlay

Thermal (Doppler) broadening

- Arises due to atoms *moving around* in the medium
- Each transition (absorption or emission) is blue/red shifted due to simple doppler velocity shift
- **The line width is therefore determined by the velocity distribution**



This area reserved for webcam overlay

Thermal (Doppler) broadening


- Thankfully we have a good model for velocity distribution of an ideal gas... our friend once again, the Maxwell-Boltzmann distribution, which classically gives

$$v_{peak} = \sqrt{2kT/m}$$

- A handy reminder for Doppler shift: $\frac{\Delta\lambda}{\lambda} = \frac{v}{c}$
(in non-relativistic limit)

- This yields good approximation of:

$$\Delta\lambda \approx \frac{2\lambda}{c} \sqrt{\frac{2kT}{m}}$$





This area reserved for webcam overlay

Thermal (Doppler) broadening

$$\Delta\lambda \approx \frac{2\lambda}{c} \sqrt{\frac{2kT}{m}}$$



- For $H\alpha$ on (above?) the Sun, we expect just the *temperature* to cause $\Delta\lambda \approx 0.04\text{nm}$
 - MUCH LARGER than Natural Broadening...
- Other velocity motions can cause Doppler broadening i.e. not explicitly *thermal* behavior
- e.g. microturbulence, especially due to convective motions of gas



This area reserved for webcam overlay

Pressure broadening

- Collisional interaction between atoms (of the same type or different), ions, e-... even molecules in some cases.
- These interactions can blur the locations of the energy levels
- Again, due to Heisenberg, this results in line broadening



This area reserved for webcam overlay

Pressure broadening

- Collisions more frequent when pressure (and density) is high in the gas
- If we use our intuition from natural broadening, $FWHM = \frac{\lambda^2}{\pi c \Delta t}$ where Δt is now the typical time between collisions
- Assume a typical velocity from the thermal motion (i.e. Doppler broadening)
- These combine to give a useful estimate:



$$\Delta\lambda \approx \frac{\lambda^2 n \sigma}{\pi c} \sqrt{\frac{2kT}{m}}$$



Pressure broadening

$$\Delta\lambda \approx \frac{\lambda^2 n\sigma}{\pi c} \sqrt{\frac{2kT}{m}}$$

- Line width due to temperature, but also density of material!
- Thus Pressure broadening is the indicator of surface gravity (mean density)
- For dwarf stars, this is often small (e.g. for $H\alpha$), but can be VERY large for high surface gravity!



This area reserved for webcam overlay

Pressure broadening

$$\Delta\lambda \approx \frac{\lambda^2 n\sigma}{\pi c} \sqrt{\frac{2kT}{m}}$$

- Various physics creating these collisions:
 - **Linear Stark Effect** (mostly for H lines, impacting Protons & e-)
 - **Quadratic Stark** (most lines in hot stars, impacting ions & e-)
 - **van der Waals** (most lines in cool stars, impacting neutral H)
- Can be VERY broad, esp. for strong dipoles (e.g. H)


STARK EFFECT




This area reserved for webcam overlay

Fraunhofer lines

- Cataloged in solar spectrum by Fraunhofer in 1814
- Seen in many other stars
- Kirchhoff & Bunsen noticed these == emission line from burning! (1859)
Lines = chemical “fingerprints”!



Ca II (or Ca+) “HK”




Whoa...




This area reserved for webcam overlay

Ca II (or Ca+) “HK”

Important tracer of chromospheric B activity




Duncan+1991, Baliunas+1998




This area reserved for webcam overlay

Ca II (or Ca+) “HK”

- **Wings** formed at photosphere surface, driven by scattering (super broad Lorentzian profile)




Duncan+1991, Baliunas+1998




This area reserved for webcam overlay

Ca II (or Ca+) “HK”

- **Peaks** due to increased temperature in chromosphere (like $H\alpha$)
- **Dip** due to scattering, increases opacity again right before the Transition Region



Duncan+1991, Baliunas+1998





This area reserved for webcam overlay

Ca II (or Ca+) “HK”

- Interestingly, the dip can be offset from line center, due to motions of gas (Doppler shift) in the upper chromosphere

Björger+2018




Duncan+1991, Baliunas+1998




This area reserved for webcam overlay

Ca II (or Ca+) “HK”

- The core is the most “interesting” part historically
- Strongly dependent on **B** heating of the chromosphere!




Duncan+1991, Baliunas+1998



This area reserved for webcam overlay

Wilson-Bappu Effect

- Empirical connection between width of Ca II line core and a star's *luminosity*




Gomez & Wallerstein (2011)




This area reserved for webcam overlay

Wilson-Bappu Effect

- Empirical connection between width of Ca II line core and a star's *luminosity*
- Since Ca II lines can be very broad, can be easier way to estimate luminosity (and thus distance) to a star...
- Ca II line definition varies from author to author works best for dwarfs with chromospheres
- Sensitive to both gravity effects AND chromospheric activity...




Ca II HK



Duncan+1991, Baliunas+1998


- Strongly dependent on **B** heating of the chromosphere!
- This means chromospheric heating changes if the total **B** strength changes.
- Ca II HK (and $H\alpha$) the most widely used indicators of total surface **B** field
e.g. see [Hall \(2008\)](#)
- Activity cycles... a rich history!



This area reserved for webcam overlay



Ca II HK

- Activity cycles... a rich history!




Duncan+1991, Baliunas+1998

Mount Wilson Observatory - HK Project



This area reserved for webcam overlay

Ca II HK



Visible light
(broadband)



Ca II K filter



This area reserved for webcam overlay



Emission lines in Wolf Rayet stars?

- Line profiles are REALLY interesting
- Very broad, very Gaussian, but not totally symmetric...



Emission lines from outflows



Red-shifted line: *motion away*



This area reserved for webcam overlay

Emission lines from outflows



Broadened line: *motion all directions*



This area reserved for webcam overlay

Emission lines from outflows




What happens when some of the outflow gets in the way of the star?



This area reserved for webcam overlay

Emission lines from outflows



P Cygni line profile:
expanding shell of material



This area reserved for webcam overlay


Emission lines in Wolf Rayet stars?

- Line profiles are REALLY interesting
- Very broad, very Gaussian, but not totally symmetric...



Inverse P Cygni Profiles

- If a P Cygni profile is a classical indicator of outflows... an *inverse* P Cygni = *infall*
- We DO see this... here's a really strange (neat) example of a star that shows **both**
- An indicator of *accretion!*



This area reserved for webcam overlay

Inverse P Cygni Profiles


- If a P Cygni profile is a classical indicator of outflows... an *inverse* P Cygni = *infall*
- We DO see this... here's a really strange (neat) example of a star that shows **both**

- An indicator of *accretion!*


- Example of H₂O lines around protostars

- “*Waterfalls around Protostars*”, amazing paper title....

Mottram+2013



33



This area reserved for webcam overlay

Next time...

- Extracting stellar parameters from spectral lines
- Also point to a really good, short (<20 min) primer on line broadening by Aaron Parsons (Berkeley): <https://www.youtube.com/watch?v=wzhnF66ZomE>



This area reserved for webcam overlay



This area reserved for webcam overlay

ASTR 421

Stellar Observations and Theory

Lecture 10

Line Profiles: III

Prof. James Davenport (UW)

36



This area reserved for webcam overlay

Today

- Actual measurements of lines (FWHM, EW)
- How we get “stellar parameters” out of spectra



This area reserved for webcam overlay

Things that lines can teach us

Line analysis

- Stellar atmospheres are parameterized by:
 - Effective temperature
 - Gravity
 - Metallicity
 - Macro/microturbulence
- The depth & shape of absorption lines depend on all of these:
 - Equivalent width ratios of the same species depend upon temperature & density
 - Pressure broadening depends upon density and temperature
 - Doppler broadening depends on temperature and turbulence
 - Line ratios between species depend upon metallicity & temperature
- We would like to go from spectral properties to stellar properties



Inverse problem


- But, the connection is complicated: requires 3D magneto-hydrodynamic simulations with radiation transfer!
- How do we go from stellar spectra to stellar properties?
- Challenges:
 - Difficult to include all lines
 - Many free parameters to adjust
 - 3D models are expensive
- Semi-empirical approach:
 - Parameterize the 1-D temperature profile
 - Fit for each absorption line separately
 - Using LTE & assumed abundances: solve for line profiles & limb-darkening (if available).
 - Infer the temperature, surface gravity, metallicity & turbulence.



Where do lines form?

- First, some terminology:

- Spectrum $\equiv (\tau = 2/3)$
- Continuum (blackbody)
- Wings (both sides)
- Core (center)
- λ_{lab} (note: vac or air wavelengths)
- **FWHM**



This area reserved for webcam overlay

Equiv width



This area reserved for webcam overlay

Curve of growth



This area reserved for webcam overlay

Estimating abundances?



This area reserved for webcam overlay



This area reserved for webcam overlay