

Scientific/Technical/Management

1 Introduction

Among the key observational properties of main sequence stars in our galaxy, age is the most difficult to determine. Traditionally, fitting isochrones to cluster stars was one of the only precise methods for measuring ages, and was impossible for the majority of isolated field stars. Methods such as asteroseismology and measuring Li abundances require time intensive observations for each target and are not capable of producing the large quantity of ages needed for exoplanet and galactic population studies. To improve our understanding of star and planet formation and evolution, as well as the history of the Milky Way, we must be able to constrain the ages of low-mass stars like the Sun in the galactic field.

Fortunately nature has provided a power means to determine ages for main sequence stars via their rotation. Angular momentum is carried away through magnetically driven stellar winds, which slows the star's rotation over cosmic time. This rotation-based "clock" is known as *gyrochronology*. Cool spots on the star's surface rotate in-to and out-of view, creating small amplitude ($\sim 1\%$) quasi-periodic changes in the stellar brightness. While rotation periods have previously been laboriously measured from starspot-induced flux modulations for hundreds of stars, space-based photometric surveys have opened the door to homogeneous ensemble measures of stellar rotation, and therefore age. **With the precise, long-duration light curves available from the *Kepler*/K2 mission, we can determine the rotation periods and ages for nearly 100,000 main sequence field stars.**

The *Kepler* mission broke new ground by producing rotation periods for tens-of-thousands of field stars within a single ~ 110 sq deg field of view, and discovered a surprising bimodal distribution of rotation periods. Two competing explanations have arisen for this mysterious feature: a bimodal age distribution for nearby stars, or a new subtlety in the angular-momentum-loss mechanism. Detailed calibrations of gyrochronology models with the *Kepler* rotation sample also revealed the need for samples of stars with a wider range of ages and compositions. Fortunately the ongoing *Kepler* extended mission, K2, has currently produced light curves from 14 additional fields throughout the Galaxy.

To enable studies of stellar ages from rotation periods with K2, we propose to:

1. Measure accurate rotation periods for every available K2 target, using new statistical methods we have developed to cope with significant instrumental systematics in the K2 data. This value-added dataset will improve the *Kepler* data legacy for field stars, and provide a critical training set for the TESS mission.
2. Produce updated gyrochronology relations based on a wider range of field star ages, and additional open clusters within the K2 fields.
3. Determine the origin of the mysterious rotation period bimodality discovered with *Kepler* by tracing the rotation period distribution in each K2 field, and out to further distances utilizing public Gaia data.
4. Measure the star formation history within each K2 field using a new Bayesian age-dating system.

2 Scientific Motivation

in new era of both precision exoplanet studies and large surveys of the Milky Way, our ability to better understand stars and their basic properties is key to these larger projects (star formation history of MWY, ages of planet hosts, environment of young-earth like systems, etc).

2.1 Rotation Periods from *Kepler* and K2

rotation periods previously were very scarce, had to do with doppler broadening, very rarely could do with photometry.

now with *Kepler* we have seen a true revolution, going from hundreds of stars with measured periods from a wide variety of instruments and approaches, to tens of thousands of stars with homogeneous data and analysis. this opens door to understanding a fundamental property of stars

2.2 Age-Dating Field Stars with Rotation

Skumanich (1972) Barnes (2007) the promise of gyrochronology is key, has ability to get ages with accuracy that rivals any other method (10%). with *Kepler*/K2 is a mass production job, but with K2 new challenges in the data are lurking




Figure 1: figure 6 from Angus et al. (2015), showing new gyrochronology relation calibrated using *Kepler* asteroseismic targets

2.3 A Mysterious Period Bimodality

Then this weird thing appeared, which is an open mystery - a bimodal period distribution - whoa (McQuillan et al., 2013). this was then extended to higher masses by Davenport (2017). Explanations are either a break from single spin-down law (possibly from different initial periods, or some new physics maybe due to chemical abundances) or represents an age bimodality of local stars.

Since this effect is seen only within 300pc so far (and only within Kepler data) due to sample properties, cannot be sure what spatial or compositional dependence this has, need a sample that spans a wider spatial area and more ages of stars




Figure 2: Left – Figure 9 from McQuillan et al. (2013); the discovery of a bimodality in rotation periods from *Kepler* M dwarfs (middle panel). Right – Figure 3 from Davenport (2017), showing all *Kepler* rotation periods from McQuillan et al. (2014) (black dots), and main sequence stars with Gaia DR1 distances. The bimodality discovered for M dwarfs extends to G and K stars, and straddles a 600 Myr “gyrochrone” (red line).

3 Proposed Research

3.1 Measuring Rotation from K2 Using the Systematics-Insensitive Periodogram

we propose the first systematic study of stellar rotation periods from the K2 data.

this will include \sim XXXXX stars from Campaigns 0-16. This includes data for multiple stellar clusters that were taken as part of the Guest Observer program.

data will be reduced using SOME FLAVOR of K2 pipeline.

the Angus et al. (2016b) rotation period pipeline will then be applied to every light curve. we will generate and save every periodogram for potential follow-up analysis of differential rotation and multi-period systems

spurious signals from transits and flares will be filtered out (some smoothing or high-freq removed in the SIP?)

for stars with good period recovery, fourier fits will identify eclipsing binaries and pulsating systems (or something like this)




Figure 3: figure 6 from Angus et al. (2016b), showing the processing of the light curve and resulting S/IPeriodogram. more simple methods give erroneous periods for this object of either 3 days via ACF, or 59 days via normal Lomb-Scargle methods.

3.2 Exploring the Period Bimodality

by combining our sample with the distances provided in the forthcoming Gaia DR2 catalog, we will be able to extend the methodology of Davenport (2017) to filter our subgiants and binary stars for our entire sample.

we then can make 3d exploration of period distribution, to see if the bimodality exists for all field stars near the sun.

the big goal here is to decide which explanation is correct!

3.3 Mapping Ages in each Field

our catalog of rotation periods from K2, combined with existing catalogs from Kepler, we will have the largest set of rotation periods ever to use for population analysis. using new technique being developed by Ruth (Chronometer) we will have improved age estimate for every star based on single rotation values. then we combine this to get age distribution within each K2 field.




Figure 4: Left – Figure 2 from Davenport (2017), showing the absolute Gaia magnitude versus temperature for *Kepler* stars with known rotation periods in the Gaia DR1 catalog. Sub-giant stars can be separated from main sequence targets using isochrone models (black solid & dashed lines). Right – Figure 4 from Davenport (2017), showing the rotation period distribution relative to a 600 Myr “gyrochrone” before (green line) and after (black line) filtering our sub-giants. The bimodal distribution is apparent, and fit with a 2-Gaussian model (blue line).

Demonstration figure?

discussion of improved gyrochronology calibration?

this is partially an extension of the period bimodality work (if it’s an age effect), but to create a detailed map of the ages of these field stars at a range of galactic latitudes in the 17 pencil-beams available from K2. combining with the periods from *Kepler*, will be even better. ultimately we’d like to compare to the age distributions in simulations of these fields from TRILEGAL, and from other age indicators (asteroseismology, flare ages, etc)

4 Team Qualifications

PI Davenport has used *Kepler* to conduct the largest survey to-date of stellar activity from flares, as well as multiple investigations of starspots and their evolution with time using *Kepler* data. From these studies, Davenport has developed an age model for flare activity that will be directly comparable to the ages and starspot amplitudes derived from this study. He has also recently discovered the rotation period bimodality first noted with *Kepler* M dwarfs by McQuillan et al. (2013) extends to G and K dwarf stars (Davenport, 2017). Davenport previously collaborated on NASA ADP grant NNX09AC77G to characterize NIR variability using the 2MASS Calibration Scan Point Source Working Database (Davenport et al., 2012, 2015). He has mentored numerous students on projects using *Kepler* data,

resulting in student-led publications such as the flare activity of a unique M dwarf binary system GJ 1245AB (Lurie et al., 2015), and exploring the poorly understood origins of wide binary stars through stellar rotation (R. Clarke in prep). He will manage the overall project, and lead the investigation and publication of the period bimodality exploration.

Co-I Angus is an expert in the extraction of periodic signals from *Kepler* data using Gaussian Processes (Angus et al., 2016a) and other cutting-edge statistical techniques. She is the author of tools for generating Systematics-Insensitive Periodograms for both *Kepler* and K2 data (Angus et al., 2016b), as well as new gyrochronology calibrations using *Kepler* asteroseismic targets (Angus et al., 2015). She will lead the effort to measure and publish rotation periods for all K2 sources, and lead the graduate student in measuring ages for field stars.

other Co-Is?

5 Relevance to NASA Programs

history of MWY, age of planet systems, TESS

6 Plan of Work

Year 1: process all rotation period data

Year 2: write papers

References

- Angus, R., Aigrain, S., & Foreman-Mackey, D. 2016a, IAU Focus Meeting, 29, 191
- Angus, R., Aigrain, S., Foreman-Mackey, D., & McQuillan, A. 2015, MNRAS, 450, 1787
- Angus, R., Foreman-Mackey, D., & Johnson, J. A. 2016b, ApJ, 818, 109
- Barnes, S. A. 2007, ApJ, 669, 1167
- Davenport, J. R. A. 2017, ApJ, 835, 16
- Davenport, J. R. A., Becker, A. C., Kowalski, A. F., et al. 2012, ApJ, 748, 58
- Davenport, J. R. A., Ruan, J. J., Becker, A. C., Macleod, C. L., & Cutri, R. M. 2015, ApJ, 803, 2
- Lurie, J. C., Davenport, J. R. A., Hawley, S. L., et al. 2015, ApJ, 800, 95
- McQuillan, A., Aigrain, S., & Mazeh, T. 2013, MNRAS, 432, 1203
- McQuillan, A., Mazeh, T., & Aigrain, S. 2014, ApJS, 211, 24
- Skumanich, A. 1972, ApJ, 171, 565