

Boulder Flood Impacts Pre-flood and Post-flood - 2013

JOHN RAESLY*

University of Colorado - Boulder

john.raesly@gmail.com

3/29/2015

Abstract

The 2013 flood in Boulder, called the 100 year flood, was a natural disaster for the century in Boulder. The floods' power changed much of the ecosystem that surrounds Boulder, CO. The purpose of this project is to examine the 2013 flood in Boulder, CO and its impacts on the river morphology of the creeks and rivers surrounding Boulder, CO and the change in land cover that occurred. Direct correlations between the rivers and the amount of rainfall in those areas feeding into the rivers are of particular interest. This study will investigate how river morphology has changed from pre to post flood? Do areas designated as floodplains affect the river change? What is the correlation between rainfall amounts and river density change? What is the land cover compared to water cover for pre versus post flood? QGIS will be utilized to perform file conversions and other usable formats for Grass GIS. Then Grass GIS will be used to integrate an analysis method for the raster data and dissolve it as necessary. Previous techniques from other flood studies that have shown to be effective will be analyzed. Sources will include the Boulder Open GIS, which has some from the NOAA Disaster Data to analyze similar conditions (rainfall amount). This study will focus on Boulder County in Colorado.

I. INTRODUCTION

Understanding the river morphology for the Boulder 2013 Flood is important for assessing the floodplains that are at the top concern if such an event were to happen again. The 2013 flood in Boulder was a natural disaster that could be cause for concern. The flood changed the stream channel within Boulder County, CO. The flood caused a total of 3 fatalities during its 8 day rain period. 306 people are still reported as missing (as of September 19th 2013). The flood had affected an area estimated to be 4,500 square miles — roughly the size of Jamaica. 12,118 people were under mandatory evacuation orders and 1,000 people had to be airlifted to safety from remote locations, the largest airlift rescue operation in USA since Katrina. 1,502 homes destroyed, 17,494 homes damaged (Colorado

Office of Emergency Management estimates) and 30 (state-maintained) bridges destroyed, 20 (state-maintained) bridges damaged (according to Colorado Department of Transportation). [1] This study focuses specifically on Boulder County. In this study, the 2013 flood in Boulder, CO is examined and its impacts on the river morphology of the creeks and rivers surrounding Boulder, CO and the change in land cover that occurred. Direct correlations between the rivers the amount of rainfall in those areas feeding into the rivers are of interest. In particular, this study will investigate: How river morphology has changed from pre to post flood?; Do areas designated as floodplains affect the river change?; What is the correlation between rainfall amounts and river density change?; What is the land cover compared to water cover for pre versus post flood?

*A thank you or further information

II. DATA

The data for this study was gathered from the Boulder Open GIS (Geographic Information Systems) on the Boulder County government website. This data is free to use and be distributed. Some of this data is sourced from a make up of NOAA's Disaster Data. NOAA (National Oceanic and Atmospheric Administration) is an American scientific agency within the United States Department of Commerce focused on the conditions of the oceans and the atmosphere. [2] The rainfall amounts on the Boulder Open GIS website were a compilation of all the days added into one dataset.

III. METHODS

Several measures were developed to assess the stream channel - pre flood against the stream channel - post flood shapefiles.

- Clip the shapefiles
- Find the difference in the shapefiles
- Branch shapefiles and find flow patterns
- Assign rainfall minimum and maximum attribute for Boulder County
- Assign general rainfall minimum and maximum to separated areas
- Correlate the rainfall amounts against the post-flood stream channel

I. Clip the Shapefiles

QGIS was used for most operations during this research. Using the built-in function, Clip, was able to show the similarities in both of the shapefiles. It created a new shape based on the area of the input layer that is overlapped by the clipping layer. The attributes of the chosen layer only were copied to the new feature. With the new information of which parts stayed the same, it was then useful to examine the portion that did not change. It was then discovered that only a small portion of the stream channel did not change. In figure 1 (image file not created yet), you can view the similar stream channel across Boulder County.

II. Find the Difference in Shapefiles

The difference function in QGIS was then used to determine the difference in the two shapefiles being examined. Both of the shapefiles had the same attributes which made it simpler to compare and contrast. The difference function created a new feature based on the area of the input layer that was not overlapped by the clipping layer.

III. Assign Rainfall Attribute for Boulder County

The purpose of this part was to assign a general legend for each area that fell into similar rainfall minimum and maximum ranges. The numbers can be viewed in figure 1. It was based upon finding a general rule for each set of values for the designated areas.

IV. Correlate Rainfall and Post-flood River Morphology

The correlation was created by implementing Spatial Autocorrelation within QGIS. It was also easily viewed by overlapping the rainfall data with the post-flood stream channel shapefile. Which can be viewed at figure 4.

IV. RESULTS

Only 48.4% of the stayed consistent with the pre-flood stream channel during this disaster. The mass of the findings concluded that most of the now stream channel was newly created. The post-flood stream channel is mostly represented by the rainfall that happened during the flood. Where there was less rain, most of those channels have disappeared and dispersed into the areas where there was the most rainfall. There was very little stream channel in Boulder, CO pre-flood, but now there are many new streams that have been created because of this flood and it mostly amounts to the mass rainfall that was recorded in the Southwestern part of the county.




Figure 1: Rainfall Distribution

V. DISCUSSION

The entirety of this disaster created many new stream channels throughout Boulder County. Whereas most of the stream channel occurred in the Northwestern part of the county it is now distributed evenly throughout the county with more of it leaning towards the eastern border. Large amounts of rainfall can be extremely effective in rerouting a whole river system. This causes many unknown side effects. There could be a loss of habitat in the area that for so long had relied on the river system, but the newly created streams could thrive and welcome all new sorts of habitat. This does mess up the ecological balance that nature depends on. It is like taking the wolves out of Yellowstone.

VI. LIMITATIONS

Since there was not a lot of time allotted for this study, some limitations have come about. The nature of the vectors provided by Boulder Open GIS did not include any information about the river density which was a subject of interest. It would have been beneficial to look at multiple other sources for data to see

if there were any conflicting results. As most weather companies have their own rain catchers', some of the resulting rainfall data could have changed from a day to day basis. As there are now many reasons to attribute to habitat loss, ie global warming, the main focus should be upon the species that whole heartedly relied upon the previous river system. Another focus of study should be if there are new disasters that were previously averted because of the river morphology. One of the disasters could be the snow melt and the floodplains. Does the new stream channel cause concern for residents situated around these floodplains that were previously unaffected?

REFERENCES

- [1] "Colorado Flood Facts and Figures — FloodList." FloodList. September 19, 2013. Accessed May 3, 2015. <http://floodlist.com/america/usa/colorado-flood-facts-figures>.
- [2] "National Oceanic and Atmospheric Administration." Wikipedia. Accessed May 3, 2015. http://en.wikipedia.org/wiki/National_Oceanic_and_Atmospheric_Administration.




Figure 2: Rainfall Distribution over Boulder County




Figure 3: Pre-Flood Stream Channel and Rainfall Distribution




Figure 4: *Post-Flood Stream Channel and Rainfall Distribution*