

Squirrel Medication Dosage Chart (listed alphabetically)

(Abbreviations: **PO** – Orally, **Sub-Q** – Subcutaneously, **IM** – Intramuscularly, **SID** – 1X/day, **BID** – 2X/day, **TID** – 3X/day, **QID** – 4x/day)

Use	Antibiotic and Antiparasitic	Antiparasitic		
Dose	15 mg / kg	7 mg / kg		
Form	50 mg / ml	25 mg / ml		
Route	PO	PO		
Frequency	BID can use up to 21 days	SID 3 - 5 days		
Weight / grams	ALBON*	BAYCOX		
10	0.00	0.00		
20	0.01	0.01		
30	0.01	0.01		
40	0.01	0.01		
50	0.02	0.01		
60	0.02	0.02		
70	0.02	0.02		
80	0.02	0.02		
90	0.03	0.03		
100	0.03	0.03		
200	0.06	0.06		
300	0.09	0.08		
400	0.12	0.11		
500	0.15	0.14		
600	0.18	0.17		
700	0.21	0.20		
800	0.24	0.22		
900	0.27	0.25		
1000	0.30	0.28		

*Note: Use BeneBac with all antibiotics

ALBON: (Sulfadimethoxine) An oral antibiotic which works on bacterial infections, especially associated with coccidiosis. Low dosage, long lasting, rapidly absorbed long-acting sulfonamide.

Form: Oral suspension (5%). Tastes really good, everyone likes it.

Life: Expiration on bottle or container. Protect from light.

Evaluation: 2-3 days. Has long treatment period (up to 21 days)

Used for: It is also used in the treatment of respiratory tract infections, Genito-urinary, protozoal enteric(inflammation of intestines) and soft tissue infections.

BAYCOX: Treatment of preclinical coccidiosis.

Form: Liquid

Life: Expiration on bottle or container. REFRIGERATE.

Evaluation: 3-5 days

Use: Treatment of Coccidiosis. Give for 3-5 days. Works quickly.

Use	Antibiotic	Antibiotic	Antibiotic	Antibiotic
Dose	5 mg / kg	10 mg / kg	5 mg / kg	10 mg / kg
Form	2.27 mg / ml	2.27 mg / ml	22.7 mg / ml	22.7 mg / ml
Route	PO, Sub Q, IM	PO, Sub Q, IM	PO, Sub Q, IM	PO, Sub Q, IM
Frequency	BID** dose can be doubled and given once a day 5-7+ days	BID** dose can be doubled and given once a day 5-7+ days	BID** dose can be doubled and given once a day 5-7+ days	BID** dose can be doubled and given once a day 5-7+ days
Weight / grams	BAYTRIL* 1:10 DILUTION LOW DOSE under 200 grams	BAYTRIL* 1:10 DILUTION HIGH DOSE under 200 grams	BAYTRIL* FULL STRENGTH LOW DOSE over 200 grams	BAYTRIL* FULL STRENGTH HIGH DOSE over 200 grams
10	0.02	0.04	For squirrels under 200 grams use 1:10 Baytril or move decimal  point to the left for full strength (e.g. for 100 gram squirrel use .022 low dose or .044 high dose. It helps to pull .1cc of water, then the small amount of Baytril	
20	0.04	0.09		
30	0.07	0.13		
40	0.09	0.18		
50	0.11	0.22		
60	0.13	0.26		
70	0.15	0.31		
80	0.18	0.35		
90	0.20	0.40		
100	0.22	0.44		
200	For squirrels over 200 grams use full strength Baytril 		0.04	0.09
300			0.07	0.13
400			0.09	0.18
500			0.11	0.22
600			0.13	0.26
700			0.15	0.31
800			0.18	0.35
900			0.20	0.40
1000			0.22	0.44

*Note: Use BeneBac with all antibiotics

BAYTRIL: Injectable or oral antibiotic which works on aerobic gram negative and many gram positive bacterial infections. Not effective against anaerobes or Streptococci.

Form: Liquid (very bad taste). **May be given twice a day, or dosage doubled and given once a day. Have available in full strength or 1:10 dosage depending on size of squirrel.

Life: Expiration on bottle. Protect from direct sunlight

Evaluation Period: 5-7 days

Used for: Dermal infections, wounds and abscesses, intestinal, respiratory infections and some urinary. Best choice for animal bites. Distributes well thru body. Effective against Chlamydia. Drug of choice for rabbits because doesn't disturb flora of intestines. Not to be used for an extended period of time on animals in rapid growth stage due to cartilage abnormalities. May work better when combined with Amoxicillin (eg.Clavamox) for trauma.

Use	Antiparasitic	Antibiotic	Antibiotic	Antibiotic
Dose		20 mg / kg	5.5 mg / kg	11 mg / kg
Form		50 mg / ml	25 mg / ml	25 mg / ml
Route	PO	PO	PO	PO
Frequency		BID 5-7+ days	BID 5-7+ days	BID 5-7+ days
Weight / grams	CAPSTAR	CLAVAMOX*	CLINDAMYCIN* LOW DOSAGE	CLINDAMYCIN* HIGH DOSAGE
10	.25 pill for older squirrels (fleas are generally controlled with kitten flea powder sprinkled on cloth and placed in basket away from direct contact with squirrel)	0.00	0.00	0.00
20		0.01	0.00	0.01
30		0.01	0.01	0.01
40		0.02	0.01	0.02
50		0.02	0.01	0.02
60		0.02	0.01	0.03
70		0.03	0.02	0.03
80		0.03	0.02	0.04
90		0.04	0.02	0.04
100		0.04	0.02	0.04
200		0.08	0.04	0.09
300		0.12	0.07	0.13
400		0.16	0.09	0.18
500		0.20	0.11	0.22
600		0.24	0.13	0.26
700		0.28	0.15	0.31
800		0.32	0.18	0.35
900		0.36	0.20	0.40
1000		0.40	0.22	0.44

*Note: Use BeneBac with all antibiotics

CAPSTAR: Oral treatment for maggots and fleas. Very safe for young animals, this product is safe in a wide dosage range. Works very quickly.

CLAVAMOX: An oral antibiotic that is a combination of Amoxicillin and Clavulanate Potassium. With the addition of Clavulanate, the Amoxicillin becomes much more active against a broad spectrum of bactericidal activity.

Form: Powder and Pill. Reconstitute powder with water. Refrigerate after reconstituting.

Life: 30 days from reconstitution (this is for generic "Clavamox" being used, name brand Clavamox is only good for 10 days after reconstitution).

Evaluation Period: 5 days

Used for: Urinary tract infections, skin, soft tissue infections and canine periodontal disease. Useful against cat bite Pasteurella. Stable with gastric juices. Make sure animal stays hydrated. May work well with Baytril for trauma.

CLINDAMYCIN: (Clindacure) Oral antibiotic, broad spectrum against many anaerobes. Do not use on rabbits or ruminants.

Form: Liquid, administered orally.

Life: Expiration date on bottle.

Evaluation: 5-7 days

Used For: General purpose antibiotic. Can be used for wound abscesses, osteomyelitis, and toxoplasmosis. Food decreases rate of absorption. May create stomach problems.

Use	Anti-inflammatory	Antifungal/parasitic	Antifungal/parasitic	
Dose	2 mg / kg			
Form	2 mg / ml	250 mg/5cc H ₂ O		
Route	IM	PO	PO	
Frequency	BID	BID	1X/week for 2-3 weeks	
Weight / grams	DEX	FLAGYL	IVERMECTIN	
10	0.01	0.01	IVERMECTIN: Broad spectrum anti-parasite useful for ecto-parasites and other skin disorder parasites. Form: Liquid. VERY STRONG. <u>Ask for help if unfamiliar.</u> Safest Route: Place drop behind one ear, 1X/week for 2-3 weeks, depending on progress, or feed nut with drop on or injected into it. Life: See bottle Evaluation: With skin, it may take time to see. Used for: Mange or other skin disorders. (Diluted can be used to wash area with maggots or thru skin absorption.)	
20	0.02	0.02		
30	0.03	0.02		
40	0.04	0.03		
50	0.05	0.04		
60	0.06	0.05		
70	0.07	0.06		
80	0.08	0.06		
90	0.09	0.07		
100	0.10	0.08		
200	0.20	0.16		
300	0.30	0.24		
400	0.40	0.32		
500	0.5	0.40		
600	0.6	0.48		
700	0.7	0.56		
800	0.8	0.64		
900	0.9	0.72		
1000	1.0	0.80		

DEXAMETHASONE: Synthetic steroid hormone anti-inflammatory. Goal is to use as much as required for as short amount of time as possible. Should not be mixed with Metacam (24 hour wait). If multiple doses of Dex are given, wait 4-5 days before giving Metacam

Form: Liquid injectable in strength 2mg/ml. **ONLY GIVE IM (HARD ON INTESTINES)**

Life: Expiration on bottle. Should be protected from light and heat

Evaluation: After initial dose wait 24 hours. Begin to taper frequency after positive response.

Used for: Head trauma, CNS or spinal trauma, shock (weak, depressed, pale mucous membranes, cold clammy skin, rapid heartbeat), and endotoxemia (results from septicemia, local infections). Not used as an Anti-fungal. May see polydipsia (thirst), polyphagia (over eating) and polyuria (excess urine).

FLAGYL: Metronidazole. Antibacterial/Antifungal bactericidal (kills bacterial as it tries to grow).

Form: Pills made to order. Protect from light or heat. REFRIGERATE once water is added.

Used for: Tricomoniasis or giardiasis in raptors, pigeons and doves. Do not use on severely debilitated, pregnant or nursing animals.

Use	Anti-inflammatory	Anti-inflammatory	Antibiotic (respiratory issues)	Antifungal
Dose	0.1 mg / kg	0.3 mg / kg	150 mg / kg	100,000 IU / kg
Form	0.15 mg / ml	0.15 mg / ml	300 mg / ml	100,000 IU / ml
Route	PO(give with food)	PO(give with food)	PO	PO
Frequency	SID	SID	Usually 1 double dose only(*see below)	QID
Weight / grams	METACAM 1:10 DILUTION LOW DOSE	METACAM 1:10 DILUTION HIGH DOSE	NUFLOR*	NYSTATIN
10	0.01	0.02	0.005	0.01
20	0.01	0.04	0.010	0.02
30	0.02	0.06	0.015	0.03
40	0.03	0.08	0.020	0.04
50	0.03	0.10	0.025	0.05
60	0.04	0.12	0.030	0.06
70	0.05	0.14	0.035	0.07
80	0.05	0.16	0.040	0.08
90	0.06	0.18	0.045	0.09
100	0.07	0.20	0.05	0.10
200	0.13	0.40	0.10	0.20
300	0.20	0.60	0.15	0.30
400	0.27	0.80	0.20	0.40
500	0.33	1.00	0.25	0.50
600	0.40	1.20	0.30	0.60
700	0.47	1.40	0.35	0.70
800	0.53	1.60	0.40	0.80
900	0.60	1.80	0.45	0.90
1000	0.67	2.00	0.50	1.00

*Note: Use BeneBac with all antibiotics

METACAM: NSAID, an anti-inflammatory analgesic, non-steroid available in oral form only. Should not be mixed with Dex; wait 24 hours after a single dose of Dex. If multiple doses of Dex are given, wait 4-5 days before using Metacam. Give with food.

Form: Oral suspension. Have full strength and 1:10 dosages, cloudy appearance is normal

Life: Expiration on bottle

Evaluation: 5-7 days

NUFLOR: An oral antibiotic for respiratory distress, especially pneumonia caused by chilling or aspiration. Very strong.

Form: Liquid. *Give only once in a double dose, and not again for 48-72 hours. Second dose not generally needed. If second dose is given, it should be only a single dose.

Life: Expiration on bottle.

Evaluation: 2-3 days

Used for: Respiratory infections.

NYSTATIN: An oral anti-fungal antibiotic (Candida) for yeast and yeast-like fungi.

Form: Liquid, store in tight, light resistant containers. Shake well before using. Store at room temperature. Deteriorates if exposed to heat, light, air or moisture.

Life: See bottle

Evaluation:

Used for: Use orally in treatment of oral or gastrointestinal CANDIDA infections. Also crop mycosis (infection caused by fungi and mycotic diarrhea).

Use	Antiparasitic	Antiparasitic	Antibiotic	
Dose	20 mg/kg	.0028/g	30 mg / kg	
Form	100 mg/ml		240 mg / 5 ml	
Route	PO	PO	PO	
Frequency	SID 3 days	1 dose, 2 nd dose in 2 weeks	BID	
Weight / grams	PANACUR	STRONGID	TMS*	
10	0.00	0.03	0.01	
20	0.00	0.06	0.01	
30	0.01	0.08	0.02	
40	0.01	0.11	0.03	
50	0.01	0.14	0.03	
60	0.01	0.17	0.04	
70	0.01	0.20	0.04	
80	0.02	0.22	0.05	
90	0.02	0.25	0.06	
100	0.02	0.28	0.06	
200	0.04	0.56	0.13	
300	0.06	0.84	0.19	
400	0.08	1.12	0.25	
500	0.10	1.40	0.31	
600	0.12	1.68	0.38	
700	0.14	1.96	0.44	
800	0.16	2.24	0.50	
900	0.18	2.52	0.56	
1000	0.20	2.80	0.63	

*Note: Use BeneBac with all antibiotics

PANACUR: Used for the removal of ascarids, hookworms and stomach worms in mammals.

Form: Liquid

Life: Expiration on bottle

Evaluation: 3 days

Use: Treatment of internal parasites, especially ones located in the lower GI Tract. More coverage than Strongid.

STRONGID: (Pyrantel Pamoate) Used for the removal of ascarids, hookworms and stomach worms in mammals. Similar to Panacur but doesn't cover as many parasites.

Form: Liquid

Evaluation:

Use: Treatment of internal parasites, especially ones located in the lower GI Tract.

TMS: (Trimethoprim and Sulfamethoxazole) A synthetic antibacterial product.

Form: Oral Suspension

Life: Expiration on bottle

Evaluation: 7-11 days

Used for: Treatment of urinary tract infections, acute otitis media (middle/inner ear infection), chronic bronchitis and pneumonia. Equivalent to Septra.