

# A simplified dynamic bioenergetic model for coral-*Symbiodinium* symbioses and coral bleaching as an alternate stable state

Ross Cunning<sup>a,\*</sup>, Erik B. Muller<sup>b</sup>, Ruth D. Gates<sup>a</sup>, Roger M. Nisbet<sup>b</sup>

<sup>a</sup>*Hawaii Institute of Marine Biology, Kaneohe, HI 96744, USA*

<sup>b</sup>*Department of Ecology, Evolution, and Marine Biology, Santa Barbara, CA 93106, USA*

---

## Abstract

Coral reef ecosystems owe their ecological success—and vulnerability to climate change—to the symbiotic metabolism of corals and *Symbiodinium* spp. The urgency to understand and predict the stability and breakdown of these symbioses (i.e., coral ‘bleaching’) demands the development and application of theoretical tools; however, the complexity of some approaches may limit their potential application and utility. Here, we develop a simplified dynamic bioenergetic model of coral-*Symbiodinium* symbioses that demonstrates realistic steady-state patterns in coral growth and symbiont abundance across gradients of light, nutrients, and feeding. Furthermore, through a mechanistic treatment of photo-oxidative stress, the model displays dynamic behaviors of bleaching and recovery, and suggests that these phenomena represent transitions between alternate stable states. A suite of complex responses to multiple, interacting environmental factors reproduced by the model suggests it unifyingly captures many important attributes of the system; meanwhile, its modular framework and open source R code are designed to facilitate expansion and problem-specific applications. We see significant potential applications for this modeling framework in generating testable hypotheses and predicting integrated, mechanistic responses of corals to environmental change.

---

## Introduction

The nutritional exchange between corals and *Symbiodinium* directly underlies the capacity of corals to build coral reef ecosystems, worth trillions of US Dollars annually (Costanza, Groot, and Sutton 2014). However, the complex symbiotic metabolism of corals is vulnerable to disruption by numerous anthropogenic environmental perturbations, jeopardizing their future persistence. In order to understand and predict responses of corals to complex changes in their environment, a mechanistic understanding of how multiple interacting factors drive the individual and emergent physiology of both symbiotic partners is necessary. Such a task is well suited for theoretical modeling frameworks such as Dynamic Energy Budget (DEB) theory (Kooijman 2010), although the complexity of such theory makes these efforts inaccessible to many biologists (Jager, Martin, and Zimmer 2013). In order to bridge this gap, we present here a simplified dynamic bioenergetic model for coral-*Symbiodinium* symbioses that aims to mechanistically integrate the impacts of complex environmental change on the physiological performance of reef corals, including responses to environmental stress.

In reef coral symbioses, intracellular *Symbiodinium* translocate photosynthetically fixed carbon to support coral metabolism, while the animal host provides access to inorganic nutrients and carbon dioxide (Muscattine and Porter 1977). Previous application of DEB theory to this syntrophic system (Muller et al. 2009) demonstrated a stable symbiotic relationship and qualitatively realistic growth and biomass ratios across gradients of ambient irradiance, nutrients, and food. This model assumed that 1) *Symbiodinium* has priority access to fixed carbon through photosynthesis, 2) the coral animal has priority access to dissolved nitrogen through contact with seawater, and 3) each partner shares with the other only what it cannot use for its own growth. In its simplest form, this

---

\*Corresponding Author

Email address: ross.cunning@gmail.com (Ross Cunning)

principle of sharing the surplus is sufficient to describe the dynamics of diverse syntrophic organs and organisms (e.g., trees, duckweeds, corals), suggesting the mechanism is mathematically and evolutionarily robust (Nisbet et al., in prep.).

While the formal DEB model of Muller et al. (2009) represents the most significant theoretical contribution in coral symbiosis research to date, we aim to strengthen the role of theory and broaden its potential application in three primary ways:

1. *Develop a module of photooxidative stress.* Of primary interest to coral biologists and ecologists is symbiosis dysfunction under environmental stress, resulting in coral “bleaching”—the loss of algal symbionts from the association (Jokiel and Coles 1977). Photooxidative stress in *Symbiodinium* is considered a primary trigger of bleaching in response to high temperature and/or light (Weis 2008), and prolonged or severe bleaching can result in mortality, though corals sometimes recover their symbionts. Bleaching susceptibility, severity, and recovery may be influenced by interacting factors such as heterotrophy and nutrient availability (Wooldridge 2014b). To simulate these events and interactions, we develop a photooxidative stress module linking overreduction of the photosynthetic light reactions to downstream impacts of photoinhibition and photodamage.
2. *Reduce theoretical and mathematical complexity.* Following the logic of Jager, Martin, and Zimmer (2013), we exclude certain features of formal DEB theory in order to capture behaviors of interest with the simplest possible formulation. Here, we present a model without reserves, maturity, or reproduction (see Kooijman 2010). This formulation restricts the model’s scope to the bioenergetics of adult corals (i.e., reproduction, larval stages, and metamorphosis are not considered), but greatly reduces theoretical complexity and parameter numbers, which is advantageous given the relative paucity of data for corals. However, our primary motivation for reducing complexity was to increase accessibility and applicability for biologists and ecologists without requiring significant expertise in DEB theory.
3. *Provide well-documented, open-access code.* In order to facilitate the continued development and application of theoretical modeling tools for coral symbioses, we provide open access to the model in the form of detailed, commented code written in the R language (R Core Team 2014). With an accessible and modular framework, we envision this model as a resource for further development by the scientific community to include additional complexity and problem-specific components. We chose the R language, which is freely available and in common use by biologists and ecologists, to reach the widest possible audience with this work.

With these as our primary motivations, we describe a simplified approach to dynamic bioenergetic modeling of coral-*Symbiodinium* symbioses that dynamically integrates the influence of external irradiance, nutrients, and prey availability on coral growth and symbiosis dynamics (i.e., symbiont:host biomass ratios), allowing for the possibility of coral bleaching in the event of photooxidative stress. In the following sections, we describe and provide rationale for the model structure, and demonstrate a range of steady state and dynamic behaviors that are consistent with observed phenomena.

## Model description

In this dynamical system, both partners acquire and use carbon and nitrogen to construct biomass: the symbiont fixes carbon through photosynthesis, and receives nitrogen shared by the host, while the host acquires nitrogen from the environment and receives carbon shared by the symbiont. A graphical representation of the model is presented in Fig. 1, and each model flux and parameter is defined in Tables 1 and 2, respectively. We use C-moles as the unit of biomass for consistency with the rigorous mass balance of DEB theory: 1 C-mole is equivalent to the amount of biomass containing 1 mole of carbon atoms. Host biomass ( $H$ ), symbiont biomass ( $S$ ), and prey biomass ( $X$ ) have fixed, but different, molar N:C ratios (Table 2). Biomass is produced from carbon and nitrogen by synthesizing units (SU), which are mathematical specifications of the formation of a product from two substrates; we use the parallel complementary formulation of Kooijman (2010) to specify these fluxes. The two state variables of this system are symbiont biomass and coral biomass; because resources are acquired proportionally to surface area, and surface area is proportional to volume (i.e., corals are “V1-morphs” in DEB

terminology (Kooijman 2010)), biomass increases exponentially during growth (indeed, corals grow exponentially (Bak 1976)).

Environmental stress is implemented in the form of photooxidative stress, which is thought to be the primary trigger of coral bleaching (Lesser 1997; Weis 2008; Wooldridge 2009). To simulate these events, we model the absorption and quenching of light energy by photochemistry and non-photochemical quenching, and the phenomenological responses that occur when these capacities are overwhelmed (i.e., photoinhibition, photodamage, and symbiont loss). While bleaching in response to high light alone has been observed experimentally (Schutter et al. 2011; Downs et al. 2013), mass coral bleaching events occur in response to high temperature (Hoegh-Guldberg 1999); thus, it is important to justify our consideration of light as the primary stressor. In reality, light and temperature interact synergistically (Coles and Jokiel 1978; Jones et al. 1998), and in fact, any stressor that disrupts the quenching of light energy may lead to bleaching (Wooldridge 2010; Baker and Cuning 2015). This is because the proximate cause of photo-oxidative stress is excess excitation energy, but the upstream events that lead to this situation may be diverse: indeed, elevated temperature may inhibit Rubisco functioning (Jones et al. 1998) and the repair of the D1 protein in photosystem II (Warner, Fitt, and Schmidt 1999), which reduces the capacity of photochemical quenching and leads to an excess of light energy. In this way, elevated temperature serves to reduce the threshold above which light stresses the system (Hoegh-Guldberg 1999); importantly, light is still the proximate stressor. Therefore, we omitted temperature from the model to maintain a desired level of simplicity, while still allowing photooxidative stress and bleaching to be simulated with biological realism in response to light.

Below we describe the formulation of each flux in the model.

Table 1. Model fluxes (mass-specific)

Symbol	Description	Units	Eq. no.
$j_X$	Prey uptake rate	molX CmolH <sup>-1</sup> d <sup>-1</sup>	3
$j_N$	Nitrogen uptake rate	molN CmolH <sup>-1</sup> d <sup>-1</sup>	4
$j_{HG}$	Host biomass formation rate	CmolH CmolH <sup>-1</sup> d <sup>-1</sup>	5
$r_{NH}$	Recycled nitrogen from host turnover	molN CmolH <sup>-1</sup> d <sup>-1</sup>	6
$\rho_N$	Nitrogen shared with the symbiont	molN CmolH <sup>-1</sup> d <sup>-1</sup>	7
$j_{eC}$	Excess carbon used to activate host CCMs	molC CmolH <sup>-1</sup> d <sup>-1</sup>	8
$j_{CO_2}$	CO <sub>2</sub> input to photosynthesis	molCO <sub>2</sub> CmolH <sup>-1</sup> d <sup>-1</sup>	9
$j_L$	Light absorption rate	molph CmolS <sup>-1</sup> d <sup>-1</sup>	11
$r_{CH}$	Recycled CO <sub>2</sub> from host turnover	molCO <sub>2</sub> CmolH <sup>-1</sup> d <sup>-1</sup>	12
$r_{CS}$	Recycled CO <sub>2</sub> from symbiont turnover	molCO <sub>2</sub> CmolS <sup>-1</sup> d <sup>-1</sup>	13
$j_{CP}$	Photosynthesis rate	molC CmolS <sup>-1</sup> d <sup>-1</sup>	14
$j_{eL}$	Light energy in excess of photochemistry	molph CmolS <sup>-1</sup> d <sup>-1</sup>	15
$j_{NPQ}$	Total capacity of NPQ	molph CmolS <sup>-1</sup> d <sup>-1</sup>	16
$c_{ROS}$	ROS production proportional to baseline	–	17
$r_{NS}$	Recycled nitrogen from symbiont turnover	molN CmolS <sup>-1</sup> d <sup>-1</sup>	18
$j_{SG}$	Symbiont biomass formation rate	CmolS CmolS <sup>-1</sup> d <sup>-1</sup>	19
$\rho_C$	Fixed carbon shared with host	molC CmolS <sup>-1</sup> d <sup>-1</sup>	20
$j_{ST}$	Symbiont turnover rate	CmolS CmolS <sup>-1</sup> d <sup>-1</sup>	21

Table 2. Model parameters. Derivations of each parameter value along with supporting references are provided in the Supplementary Information.

Symbol	Description	Units	Value
$n_{NH}$	N:C molar ratio in host biomass	–	0.18
$n_{NS}$	N:C molar ratio in symbiont biomass	–	0.13
$n_{NX}$	N:C molar ratio in prey biomass	–	0.2
$j_{HT}^0$	Specific turnover rate of host biomass	CmolH CmolH <sup>-1</sup> d <sup>-1</sup>	0.03
$j_{ST}^0$	Specific turnover rate of symbiont biomass	CmolS CmolS <sup>-1</sup> d <sup>-1</sup>	0.03

Symbol	Description	Units	Value
$\sigma_{NH}$	Proportion host N-turnover recycled to host	–	0.9
$\sigma_{CH}$	Proportion host C-turnover recycled to photosynthesis	–	0.1
$\sigma_{NS}$	Proportion symbiont N-turnover recycled to symbiont	–	0.9
$\sigma_{CS}$	Proportion symbiont C-turnover recycled to photosynthesis	–	0.9
$j_{Xm}$	Maximum specific feeding rate of host	molX CmolH <sup>-1</sup> d <sup>-1</sup>	0.13
$K_X$	Half-saturation constant for prey uptake by host	molX L <sup>-1</sup>	$1.95 \times 10^{-5}$
$j_{Nm}$	Maximum specific DIN uptake rate by host	molN CmolH <sup>-1</sup> d <sup>-1</sup>	0.035
$K_N$	Half-saturation constant for DIN uptake by host	molN L <sup>-1</sup>	$1.5 \times 10^{-6}$
$k_{CO_2}$	Efficacy of CO <sub>2</sub> delivery by host CCMs	molCO <sub>2</sub> molC <sup>-1</sup>	10
$j_{HGm}$	Maximum specific growth rate of host	CmolH CmolH <sup>-1</sup> d <sup>-1</sup>	1
$y_{CL}$	Quantum yield of photosynthesis	molC mol ph <sup>-1</sup>	0.1
$\bar{a}^*$	Effective light-absorbing cross-section of symbiont	m <sup>2</sup> CmolS <sup>-1</sup>	1.34
$k_{NPQ}$	NPQ capacity of symbiont	mol ph CmolS <sup>-1</sup> d <sup>-1</sup>	112
$k_{ROS}$	Excess photon energy that doubles ROS production, relative to baseline levels	mol ph CmolS <sup>-1</sup> d <sup>-1</sup>	80
$j_{CPm}$	Maximum specific photosynthesis rate of symbiont	molC CmolS <sup>-1</sup> d <sup>-1</sup>	2.8
$j_{SGm}$	Maximum specific growth rate of symbiont	CmolS CmolS <sup>-1</sup> d <sup>-1</sup>	0.25
$b$	Scaling parameter for strength of bleaching response	–	5

### State equations

The balance equations for symbiont ( $S$ ) and host ( $H$ ) biomass are expressed as:

$$\frac{dS}{Sdt} = j_{SG} - j_{ST} \quad (1)$$

$$\frac{dH}{Hdt} = j_{HG} - j_{HT}^0 \quad (2)$$

These are expressed per unit of symbiont and host biomass, respectively. The specific biomass growth and turnover rates that define these balance equations are produced by combinations of the individual model fluxes (see Table 1 for definitions and units), which are each expressed as mass-specific rates (e.g., per C-mole of symbiont or host biomass per day). When necessary, conversions between symbiont-mass-specific and host-mass-specific rates are accomplished by multiplying or dividing by the symbiont:host biomass ratio.

### Coral animal fluxes

The coral animal acquires both carbon and nitrogen from feeding on prey from the environment. Prey acquisition is specified by Michaelis-Menten kinetics (i.e., a Holling type II function) with a maximum feeding rate  $j_{Xm}$  and half-saturation constant  $K_X$ :

$$j_X = \frac{j_{Xm} \cdot X}{X + K_X} \quad (3)$$

Additionally, the coral animal acquires nitrogen dissolved in the surrounding seawater, which is assumed to represent ammonium, the primary form utilized by corals (Yellowlees, Rees, and Leggat 2008). This gives the host (rather than the symbiont) priority in nitrogen utilization; this capacity is supported by experimental evidence (Wang and Douglas 1998) and is consistent with the physical arrangement of the partners, where the host is in

direct contact with the external environment. The uptake of nitrogen from the environment is thus specified by Michaelis-Menten kinetics using a maximum uptake rate  $j_{Nm}$  and half-saturation constant  $K_N$ :

$$j_N = \frac{j_{Nm} \cdot N}{N + K_N} \quad (4)$$

Coral biomass formation is then specified by a parallel complementary SU that combines carbon and nitrogen, according to:

$$j_{HG} = \left( \frac{1}{j_{HGm}} + \frac{1}{\rho_C \frac{S}{H} + j_X} + \frac{1}{(j_N + n_{NX}j_X + r_{NH})/n_{NH}} - \frac{1}{\rho_C \frac{S}{H} + j_X + (j_N + n_{NX}j_X + r_{NH})/n_{NH}} \right)^{-1} \quad (5)$$

where  $\rho_C$  is fixed carbon shared by the symbiont (see Eq. 20), and  $r_{NH}$  is recycled nitrogen liberated by host biomass turnover:

$$r_{NH} = \sigma_{NH} n_{NH} j_{HT}^0 \quad (6)$$

The amount of nitrogen input to the coral biomass SU in excess of what is actually consumed in biomass formation (i.e., surplus nitrogen, or the rejection flux<sup>1</sup> of the SU) is then made available to the symbiont:

$$\rho_N = (j_N + n_{NX}j_X + r_{NH} - n_{NH}j_{HG})_+ \quad (7)$$

Due to the inherent inefficiency of the parallel complementary SU formulation, there is always some nitrogen shared with the symbiont even when coral biomass formation is strongly nitrogen-limited. Likewise, there is always a non-zero rejection flux of excess carbon from the coral biomass SU. Likewise, the carbon rejected from this SU reflects the amount of excess fixed carbon available to the host that is not used in biomass formation:

$$j_{eC} = (j_X + \rho_C \frac{S}{H} - j_{HG})_+ \quad (8)$$

This flux,  $j_{eC}$ , is assumed to be available to the host as a respiratory substrate to support various energetically-demanding processes; of particular importance is the host's active carbon concentrating mechanisms (CCMs) that supply  $\text{CO}_2$  for symbiont photosynthesis (Wooldridge 2013; Hopkinson, Tansik, and Fitt 2015). We therefore specify  $j_{\text{CO}_2}$  as the host-mediated delivery of  $\text{CO}_2$  to photosynthesis that encompasses potentially diverse carbon-concentrating mechanisms (CCMs), including active transport of bicarbonate, carbonic anhydrase-catalyzed conversion of bicarbonate to  $\text{CO}_2$  to promote diffusion toward the symbiont (Tansik, Fitt, and Hopkinson 2015), and acidification of the symbiosome to increase localized  $\text{CO}_2$  concentrations around the symbiont (Barott et al. 2014). Since these active CCMs require energetic input by the host, we define  $j_{\text{CO}_2}$  as proportional to  $j_{eC}$ , assuming that some of this carbon is respired to energize the CCMs. This formulation means that the symbiont indirectly ensures its own  $\text{CO}_2$  supply by providing fixed carbon (=energy) to the host (Wooldridge 2013), which establishes a positive feedback in the model that leads to the existence of multiple stable states, which we discuss below in the context of coral bleaching. The parameter  $k_{\text{CO}_2}$  scales the efficacy of host CCMs, which enables the comparison of different rates of  $\text{CO}_2$  delivery that may characterize different coral species (Wooldridge 2014a). The active input of  $\text{CO}_2$  to the photosynthesis SU is therefore specified as:

$$j_{\text{CO}_2} = k_{\text{CO}_2} j_{eC} \quad (9)$$

---

<sup>1</sup>Rejection fluxes must always be positive, and hence are specified with the notation  $(x)_+$ , which means  $\max(x, 0)$ .

## Symbiodinium fluxes

The symbiont produces fixed carbon through photosynthesis, a process represented here by a single SU with two substrates: light (photons) and inorganic carbon ( $\text{CO}_2$ ). The amount of light absorbed by the symbiont depends on the scalar irradiance at the site of light absorption, which is modified substantially relative to external downwelling irradiance owing to multiple scattering by the coral skeleton and self-shading by surrounding symbionts (Enríquez, Méndez, and Iglesias-Prieto 2005; Marcelino et al. 2013). We used data from Marcelino et al. (2013) to empirically derive an amplification factor,  $A$ , indicating the ratio of internal scalar irradiance to external downwelling irradiance as a function of symbiont density (S:H biomass), which is specified as:

$$A = 1.26 + 1.39 \cdot \exp\left(-6.48 \cdot \frac{S}{H}\right) \quad (10)$$

This amplification factor is then multiplied by the external downwelling irradiance  $L$  and the effective light-absorbing surface area of symbiont biomass  $\bar{a}^*$  to specify the total light absorption:

$$j_L = A \cdot L \cdot \bar{a}^* \quad (11)$$

$\text{CO}_2$  arrives at the photosynthesis SU from multiple sources: in addition to the  $\text{CO}_2$  actively supplied by the host through its CCMs ( $j_{\text{CO}_2}$ ; Eq. 9), a fixed proportion of metabolic  $\text{CO}_2$  generated by host biomass turnover is passively available to the photosynthesis SU, according to:

$$r_{CH} = \sigma_{CH} j_{HT}^0 \quad (12)$$

along with a fixed proportion of  $\text{CO}_2$  generated by symbiont biomass turnover:

$$r_{CS} = \sigma_{CS} j_{ST}^0 \quad (13)$$

Fixed carbon is then produced by the photosynthesis SU according to:

$$j_{CP} = \left( \frac{1}{j_{CPm}} + \frac{1}{y_{CL} j_L} + \frac{1}{(j_{\text{CO}_2} + r_{CH}) \frac{H}{S} + r_{CS}} - \frac{1}{y_{CL} j_L + (j_{\text{CO}_2} + r_{CH}) \frac{H}{S} + r_{CS}} \right)^{-1} \cdot c_{ROS}^{-1} \quad (14)$$

where  $j_{CPm}$  is the maximum specific rate of photosynthesis, and  $c_{ROS}$  is the relative rate of reactive oxygen species production (see Eq. 17). Dividing the photosynthetic rate by  $c_{ROS}$  causes a decline in response to photooxidative stress at high light levels, and the emergent outcome of this SU formulation demonstrates a classic photoinhibition response (Fig. S2).

Light energy absorbed in excess of what is used to fix carbon is specified by the SU rejection flux, according to:

$$j_{eL} = (j_L - j_{CP}/y_{CL})_+ \quad (15)$$

This excess light energy must be quenched by alternative pathways in order to prevent photooxidative damage (Powles 1984). *Symbiodinium* utilize a variety of pathways for non-photochemical quenching (NPQ; Roth 2014), which we collect in a total NPQ capacity specified as a parameter of the symbiont ( $k_{NPQ}$ ). The NPQ flux  $j_{NPQ}$  is then specified as a single-substrate SU formula with a maximum of  $k_{NPQ}$ :

$$j_{NPQ} = \left( \frac{1}{k_{NPQ}} + \frac{1}{j_{eL}} \right)^{-1} \quad (16)$$

If light energy further exceeds the capacity of both photochemistry and NPQ, then reactive oxygen species (ROS) are produced. We represent this as a relative quantity  $c_{ROS}$ , which takes a value of 1 when all light energy

is quenched by photochemistry and NPQ, and increases as the amount of excess excitation energy increases, specified as:

$$c_{ROS} = 1 + \frac{(j_{eL} - j_{NPQ})_+}{k_{ROS}} \quad (17)$$

where  $k_{ROS}$  is a parameter of the symbiont that determines the rate of ROS production (specifically, the amount of excess excitation energy that doubles ROS production relative to baseline levels). Importantly,  $c_{ROS}$  is specified here not as a function of absolute external light, but rather the amount of excess light after accounting for quenching by carbon fixation and NPQ. A direct consequence of this formulation is that CO<sub>2</sub>-limitation of photosynthesis can lead to ROS production, an important mechanism (Wooldridge 2009) that was not captured by previous representations of photooxidative stress (Eynaud, Nisbet, and Muller 2011).

Carbon fixed by photosynthesis ( $j_{CP}$ ; Eq. 14) is then combined with nitrogen shared by the host ( $\rho_N$ ; Eq. 7) and nitrogen recycled from symbiont biomass turnover

$$r_{NS} = \sigma_{NS} n_{NS} j_{ST}^0 \quad (18)$$

to build new symbiont biomass, following the SU equation:

$$j_{SG} = \left( \frac{1}{j_{SGm}} + \frac{1}{j_{CP}} + \frac{1}{(\rho_N \frac{H}{S} + r_{NS})/n_{NH}} - \frac{1}{j_{CP} + (\rho_N \frac{H}{S} + r_{NS})/n_{NH}} \right)^{-1} \quad (19)$$

The rejection flux of carbon from this SU represents the amount of fixed carbon produced by photosynthesis in excess of what can be used to produce symbiont biomass; this surplus,  $\rho_C$ , is translocated to the coral host:

$$\rho_C = (j_{CP} - j_{SG})_+ \quad (20)$$

The rejection flux of nitrogen from the symbiont biomass SU is lost to the environment.

Symbiont biomass turnover includes a component of constant turnover specified by the parameter  $j_{ST}^0$ , representing fixed maintenance costs, plus a component that scales with the magnitude of ROS production.

$$j_{ST} = j_{ST}^0 (1 + b(c_{ROS} - 1)) \quad (21)$$

This second component of symbiont biomass loss represents both photodamage and/or symbiont expulsion (i.e., bleaching), both of which occur in response to high levels of ROS production. The parameter  $b$  is included to scale biomass loss due to bleaching in response to ROS. (Note that recycling of symbiont biomass turnover ( $r_{NS}$  and  $r_{CS}$ ) only occurs based on the maintenance component of turnover (i.e.,  $j_{ST}^0$ ), and not the photodamage/bleaching component, as this loss represents biomass that is damaged or expelled from the host.)

### *Numerical analysis*

A time-stepping Euler method was used to solve the state equations since the production and rejection fluxes of the SUs are implicitly defined due to feedbacks present in the model. Specifically, the rejection fluxes of carbon and nitrogen from the symbiont and host biomass SUs act as reciprocal input fluxes to the other SU. In addition, the photosynthesis SU receives CO<sub>2</sub> proportional to the carbon rejection flux from the host biomass SU, and the rejection flux of excitation energy from the photosynthesis SU acts to reduce its own production (i.e., photoinhibition). A vector of time steps is used for each simulation, along which dynamic environmental forcing functions (irradiance, DIN, and prey abundance). These vectors, along with initial values of symbiont and host biomass, then serve as input to the time-stepping function, which solves for the current system state using values of the previous system state where necessary. A time step of 0.1 days was used for all simulations, which were performed using R code that is available in the data repository accompanying this

article: [github.com/jrcunning/Rcoral](https://github.com/jrcunning/Rcoral). For steady state estimations, simulations were run until the change in both growth and S:H biomass was  $< 5e-7$  per time step.

To aid in visualizing model results, we calculated values to indicate the degree to which product formation at an SU was limited by availability of either of its two substrates:

$$\log \left( \frac{\min(j_{S1}, j_{Pm})}{\min(j_{S2}, j_{Pm})} \right) \quad (22)$$

where  $j_{S1}$  and  $j_{S2}$  are the specific input fluxes of the two substrates and  $j_{Pm}$  is the maximum specific product formation rate, in units of  $\text{Cmol Cmol}^{-1} \text{d}^{-1}$ .

## Steady state behavior

In a constant environment, the system ultimately reaches a steady state of exponential growth or decline. However, under some conditions, either of these outcomes may occur depending on initial values of symbiont and host biomass, indicating the presence of alternate stable states. The mechanism that produces these alternate stable states is the positive feedback between carbon-limitation of the host and  $\text{CO}_2$ -limitation of photosynthesis: if symbiont biomass is initially very low (i.e., a “bleached” coral), the system cannot escape this positive feedback and cannot grow. However, if symbiont biomass is initially high (i.e., a “healthy” coral), then the system remains in a nitrogen-limited state with positive growth. For practical purposes, this section of the manuscript considers only positive growth steady states under constant environments; the subsequent “dynamic behavior” section explores how changing environments may cause the system to switch between alternate stable states, which we interpret in the context of coral bleaching.

To analyze steady state behavior, we ran the model to steady state across gradients of external irradiance and nutrients (Fig. 2), which revealed patterns consistent with observed phenomena in corals. Predicted growth rates are low at low light and DIN ( $\sim 0.01 \text{ d}^{-1}$ ), and begin increasing as both of these factors increase (Fig. 2A). Low light limits photosynthetic rates, resulting in less fixed carbon shared with the host and an associated increase in the symbiont to host biomass ratio (Fig. 2B). In agreement with this trend are many observations of negative correlation between irradiance and symbiont density (Stimson 1997; Brown et al. 1999; Fitt et al. 2000; Titlyanov et al. 2001). As higher light alleviates light-limitation of photosynthesis, host growth becomes less carbon-limited.

Similarly, increasing DIN alleviates nitrogen-limitation (Fig. 2A). Increased growth at higher DIN is predicted by the DEB model of Muller et al. (2009), and has also been observed experimentally (Muller-Parker et al. 1994; Tanaka et al. 2007; Tanaka et al. 2013). However, DIN elevation beyond a certain point (e.g.,  $\sim 3\text{-}4 \text{ }\mu\text{M}$  in these simulations) has little effect on growth as carbon becomes limiting. Very high nutrient levels may even reduce growth (Shantz, Lemoine, and Burkepile 2015), although these impacts are not likely to occur within the range of concentrations considered here ( $< 4 \text{ }\mu\text{M}$ ) (Ferrier-Pagès et al. 2000). In addition to increasing growth, DIN also increases the symbiont to host biomass ratio (Fig. 2B), a phenomenon also observed in reef corals (Marubini and Davies 1996). At low DIN and intermediate light, more typical of coral reef environments, symbiont to host biomass ratios are around  $\sim 0.06\text{-}0.21$ , which is consistent with values reported in the literature (Muscattine, R McCloskey, and E Marian 1981; Edmunds et al. 2011; Hawkins et al. 2016).

The maximum predicted growth rates of  $\sim 0.1 \text{ d}^{-1}$ , occurring between  $\sim 10\text{-}25 \text{ mol photons m}^{-2} \text{ s}^{-1}$  light and  $\sim 4 \text{ }\mu\text{M}$  DIN (Fig. 2A), are comparable to the rate of  $0.07 \text{ d}^{-1}$  measured by Tanaka et al. (2007) under similar N-enriched conditions. Predicted growth rates under conditions more typical of reef environments ( $< 0.5 \text{ }\mu\text{M}$  DIN) are  $\sim 0.01\text{-}0.03 \text{ d}^{-1}$ . Observed specific growth rates, which are typically based on skeletal mass, fall near or below  $0.01 \text{ d}^{-1}$  (Osinga et al. 2011; Osinga et al. 2012), though values as high as  $0.025 \text{ d}^{-1}$  have been reported (Schutter et al. 2010). It is not surprising that predicted growth rates are slightly higher than most observations, as the model does not account for external ecological factors that may reduce growth, such as competition, predation, and bioerosion. Furthermore, the model does not explicitly account for skeletal formation, which may not always correlate with biomass growth, potentially due to differential substrate-limitation (Anthony 2002). Nevertheless, when growth of coral tissue biomass has been directly measured over short periods, values are consistent with



model predictions (e.g.,  $0.07 \text{ d}^{-1}$  in Tanaka et al. (2007)), and similar rates ( $0.04 \text{ d}^{-1}$ ) have been measured in *Aiptasia diaphana*, a non-calcifying symbiotic anemone (Armoza-Zvuloni et al. 2014).

As irradiance continues to increase above  $\sim 25 \text{ mol photons m}^{-2} \text{ d}^{-1}$ , growth rates decline until positive growth ceases above  $\sim 40 \text{ } \mu\text{mol photons m}^{-2} \text{ d}^{-1}$  (Fig. 2A). The mechanism underlying this decline is the increase in light energy beyond the capacities of photosynthesis and non-photochemical quenching: excess excitation energy generates reactive oxygen species (ROS) (Weis 2008; Roth 2014), which, in this model, have the phenomenological consequences of reducing the photosynthetic rate (representing photoinhibition) and increasing symbiont biomass loss (representing photodamage and/or symbiont expulsion) (see Eynaud, Nisbet, and Muller 2011). Together, these impacts reduce the symbiont to host biomass ratio (Fig. 2B), as occurs during coral bleaching. This reduction in symbionts consequently reduces the flux of fixed carbon to the host, resulting in increasing carbon-limitation (Fig. 2B) and eventual cessation of growth (Fig. 2A).

The incorporation of photooxidative stress in the model sets an upper limit to the amount of light at which a stable symbiotic interaction can be maintained, but even below this threshold of breakdown, negative effects of high light reduce steady state growth and symbiont:host biomass (Fig. 2). This gradual decline is consistent with experimental results showing that high light levels decrease growth (Schutter et al. 2011), and field studies documenting optimum growth rates at intermediate depths (Baker and Weber 1975; Huston 1985). By incorporating these impacts of light stress, the model predicts greater, and more realistic, variation in state variables across light gradients than was predicted by the model of Muller et al. (2009), which did not include photoinhibition and photodamage. It is important to recognize that the boundary set by photooxidative stress on a stable symbiosis under steady state conditions (Fig. 2) may be temporarily crossed by a dynamic system, which may experience a period of symbiont loss (bleaching) and reduced growth, after which a return to benign conditions may restore symbiont biomass and positive growth. To explore this further and illustrate the behavior of the model in more detail, we evaluate a number of dynamic simulations below (see “Dynamic behavior”).

## Sensitivity analysis

The values used for each parameter in the model (Table 2) are derived from relevant literature; these derivations are described in the Supplementary Information. Here we evaluate the sensitivity of the model to changes in these parameter values, which also serves to demonstrate the behavior of the dynamical system. We measured fractional change in steady state values in response to fractional changes in parameter values, relative to their default values, under environmental conditions typical of coral reefs.

Overall, relative changes in the steady state of the system are less than the equivalent relative change in parameter value, indicating that the system maintains internal stability. However, changes in certain parameter values have more significant impacts: increasing  $j_{Nm}$  or decreasing  $K_N$  both dramatically increase host growth (Fig. 3), demonstrating the strong nitrogen-limitation that characterizes these symbioses. The parameter  $\bar{a}^*$  has a strong impact on S:H biomass ratios (Fig. 3) since this parameter determines the amount of light absorbed by symbionts, with lower values increasing light-limitation. Increasing the maximum growth and turnover rates have the expected effects of increasing and decreasing growth, respectively. Parameters relating to photooxidative stress and bleaching have little impact under these environmental conditions (Fig. 3), but have larger impacts under higher light (e.g., Fig. S6). Sensitivity analyses conducted under different combinations of external light and nutrients are presented in Figs. S3-S7.

## Dynamic behavior

The dynamic behavior of the model demonstrates its power to integrate multiple environmental forcings simultaneously. Here we present several scenarios that demonstrate the model’s ability to reproduce complex phenomena that have been observed in corals.

### *Seasonal variability*

Symbiont densities and coral growth rates are known to vary seasonally, representing an integrated response to changes in a suite of environmental factors. Light in particular is a strong driver of these trends (Stimson 1997; Brown et al. 1999; Fagoonee et al. 1999; Fitt et al. 2000), with high light associated with lower symbiont abundance and reduced tissue biomass. The role of light in driving seasonal changes in symbiont density was demonstrated nicely by Stimson (1997), who also found that experimental nutrient-enrichment amplified the light-driven seasonal oscillation. Using the levels of light and nutrients from this study as inputs, the model reproduces this observed interaction among environmental factors (Fig. 4), and also provides the mechanism: under high light in summer, photooxidative stress decreases symbiont biomass and increases carbon-limitation of host growth regardless of nutrient status; when light is reduced in winter, growth and symbiont biomass become constrained instead by nitrogen, and therefore increase more when DIN is enriched (Fig. S8).

The prediction of higher growth when light is reduced indicates that growth is not limited by low light in winter, but is actually reduced by excess light in summer, consistent with the experimental findings of Schutter et al. (2011). Seasonal summertime reductions in tissue biomass have also been well-documented in the field (Fitt et al. 2000), along with reductions in net photosynthetic capacity (Muller-Parker 1987). Importantly, while light alone (at these levels) may drive seasonal dynamics in the ways discussed, temperature fluctuations may attenuate or even reverse the effect of light as cooler winters depress metabolism; thus, the relative magnitude of fluctuation in temperature, light, and other factors may produce wide variability in the direction and magnitude of seasonal changes in growth and symbiont abundance, depending on location and microhabitat. Nevertheless, the seasonal variability predicted here (Fig. 4) is consistent with experimental and field observations for corals, and demonstrates the power of this model to predict dynamic behavior that mechanistically integrates multiple environmental drivers.

### *Coral bleaching*

Coral bleaching is the stress-induced loss of symbiotic algae from coral tissues, which can occur in response to a variety of environmental stressors. In most cases, coral bleaching is thought to begin with photooxidative stress in symbiont photosynthesis (Lesser 1997), which triggers a cascade of events leading to symbiont expulsion (Weis 2008). As symbionts are expelled, the host receives less fixed carbon, which may then compromise its ability to activate CCMs that deliver  $\text{CO}_2$  to photosynthesis (Wooldridge 2013). Increasing  $\text{CO}_2$ -limitation for remaining symbionts, along with an amplified internal light environment due to reduced self-shading (Enríquez, Méndez, and Iglesias-Prieto 2005), may further exacerbate photooxidative stress and accelerate symbiont expulsion, driving the coral into a bleached state.

While these positive feedbacks have been discussed previously in the literature, this is the first attempt to implement and explore their properties within a dynamical model. Interestingly, these feedbacks lead to alternate stable states. The ‘healthy’ state is characterized by nitrogen-limitation of both symbiont and host: under these conditions, the symbiont translocates sufficient carbon to support host growth and CCMs, which ensures that photosynthesis does not become  $\text{CO}_2$ -limited. However, if carbon translocation is disrupted (and light is sufficiently high), then the system is driven into the ‘bleached’ state by photooxidative stress and positively reinforcing carbon- and  $\text{CO}_2$ -limitation. In this context, coral bleaching can be understood as a transition from one stable state to another, and bleaching thresholds are sets of environmental conditions that push a healthy-state coral onto a trajectory leading to a bleached state.

We are highly interested in the conditions under which the system switches from a healthy to a bleached state, and can use this model as a tool to explore this dynamic behavior. Most straightforwardly, this switch occurs when increasing external irradiance (Fig. 5A) causes sufficient ROS production (Fig. 5B) and photoinhibition (Fig. 5C) such that the positive feedbacks between host carbon-limitation (Fig. 5D) and  $\text{CO}_2$ -limitation of photosynthesis (Fig. 5E) are rapidly engaged, leading to even greater photooxidative stress and a rapid decline in S:H biomass (Fig. 5F), characteristic of coral bleaching. However, the positive feedbacks involved in bleaching are not engaged only in response to high external irradiance alone; in fact, they depend on the relative balance of light energy absorption and quenching, which in turn depends on the availability of  $\text{CO}_2$  for photosynthesis. While previous models framed photooxidative stress as a fixed response to absolute external irradiance (Eynaud, Nisbet, and Muller 2011), our implementation considers the dynamic balance of multiple energy sinks in the causation of

stress, which is more consistent with current understanding of symbiosis dysfunction (Wooldridge 2013), and establishes a critical role of host CCMs in providing CO<sub>2</sub> for photosynthesis (Tansik, Fitt, and Hopkinson 2015; Hopkinson, Tansik, and Fitt 2015).

The importance of host CCM activity establishes significant interactive roles for other factors in influencing coral bleaching responses. For example, simulations of high light stress (Fig. 6) demonstrate that bleaching can be attenuated by heterotrophic feeding, a phenomenon which has been observed experimentally (Borell et al. 2008). The mechanism underlying this prediction is that feeding by the host increases host CCM activity, which delays the onset of CO<sub>2</sub>-limitation of photosynthesis and reduces bleaching severity. On the other hand, elevated nutrients exacerbate bleaching (Fig. 6), since higher symbiont densities are more susceptible to CO<sub>2</sub>-limitation (Wooldridge 2009). Several experimental (Cunning and Baker 2013; Vega Thurber et al. 2014) and correlational studies (Wooldridge and Done 2009) are consistent with this mechanistic link between high nutrients and bleaching.

Since bleaching represents an alternative stable state, it follows that a bleached coral cannot recover unless the external condition that induced bleaching is alleviated. In natural settings, this typically occurs by seasonal declines in temperature and light. However, due to hysteresis, a coral cannot recover along the same trajectory it followed during bleaching; indeed, the stressor must be alleviated well *below* the threshold that initially caused bleaching in order for the system to recover (Fig. 7). This is because under the same external conditions, a bleached coral (relative to a healthy one) is characterized by greater light amplification and weaker CCM activity, which serve to maintain the bleached state. The magnitude of hysteresis will depend on external nutrients and prey availability, as well as model parameters, yet the fundamental concept demonstrates a core behavior of the system.

The hysteresis demonstrated in Fig. 7 reveals another interesting behavior of the system: during recovery from bleaching, the S:H biomass ratio reaches a higher value that ‘overshoots’ the ratio maintained in the healthy stable state. In fact, such a phenomenon has been observed in both experimental (Cunning, Silverstein, and Baker 2015) and field studies (Kemp et al. 2014) documenting recovery from bleaching, and has been interpreted as a potential ‘disequilibrium in host-symbiont regulation’ (Kemp et al. 2014). The model explains this ‘overshoot’ as an approach to a bifurcation point in the dynamical system: during recovery from the ‘bleached’ state, host growth is more strongly carbon-limited than symbiont growth (since the symbiont has priority access to carbon), so S:H biomass increases. S:H biomass will continue to increase beyond the alternative steady state levels as long as the symbiont remains carbon-limited; not until nitrogen becomes more strongly limiting to symbiont growth than carbon (hence forcing the symbiont to share carbon with the host) will the system flip back to the ‘healthy’ state, characterized by high carbon fixation rates and overall nitrogen-limitation. Of course, the magnitude of this ‘overshoot’ depends on external nutrient concentrations and the rate of change in the stressor during recovery, though simulations predict similar magnitudes to what was observed by Kemp et al. (2014) and Cunning, Silverstein, and Baker (2015) (Fig. S9).

## Conclusions

This simplified dynamic bioenergetic model of coral-*Symbiodinium* symbioses mechanistically reproduces patterns in steady-state coral growth and symbiont abundance commonly observed in corals, including higher symbiont abundance with higher nutrients and feeding, lower symbiont abundance with increasing light, and optimal growth at intermediate light levels. Moreover, the model reproduces complex dynamic behaviors including seasonal changes in symbiont density at different nutrient levels, rapid bleaching above a threshold of high light, mitigation of bleaching by heterotrophic feeding, exacerbation of bleaching by elevated nutrients, and an overshoot of symbiont density during recovery from bleaching. These examples demonstrate the model’s ability to integrate multiple environmental forcing functions to reproduce complex responses; meanwhile, the diversity of these phenomena suggest the model has captured many of the important features of the system in a unifying mechanistic framework. This model also provides a new conceptual framework for considering coral bleaching as a transition to an alternate stable state, which has important implications for understanding the performance and maintenance of symbiotic interactions. In this context, the ‘healthy’ stable state represents a scenario in which the host is ‘in control’ of the system, with nitrogen-limitation stabilizing the symbiont to host biomass ratio and maintaining positive growth. Conversely, carbon-limitation represents a loss of host ‘control’, wherein

positive feedbacks result in a loss of capacity for photosynthetic carbon fixation and positive growth. Importantly, these ‘control’ mechanisms arise entirely passively through feedbacks present in the system.

In addition to the suite of specific examples presented here, the model can be used to explore many different dynamic environmental scenarios, and represents a tool that biologists and ecologists can use to generate hypotheses and make predictions in both experimental and natural settings. Moreover, parameter values can be modified to correspond to different genetic or functional types of coral hosts and *Symbiodinium* partners in order to evaluate variability in system responses. Thus, the diversity of potential applications for this model is high, and we envision this work as a foundation for continued development, which may include more detailed treatments of specific modules (e.g., DIC processing), and the incorporation of more external forcing capabilities (e.g., external DIC). Importantly, this effort is best undertaken by the scientific community at large, including those with theoretical and empirical backgrounds, which is why the code for this model has been developed openly using the R language. Ultimately, the continued refinement of these tools is fundamental in elucidating the mechanisms of symbiosis function and dysfunction, and in predicting coral responses to environmental change.

## References

- Anthony, K. 2002. “Comparative analysis of energy allocation to tissue and skeletal growth in corals.” *Limnology and Oceanography* 47 (5): 1417–29. <https://www.scopus.com/inward/record.uri?partnerID=HzOxMe3b&scp=0036733455&origin=inward>.
- Armoza-Zvuloni, Rachel, Esti Kramarsky-Winter, Yossi Loya, Ami Schlesinger, and Hanna Rosenfeld. 2014. “Trioecy, a unique breeding strategy in the sea anemone *Aiptasia diaphana* and its association with sex steroids.” *Biology of Reproduction* 90 (6). Society for the Study of Reproduction: 122–22. doi:10.1095/biolreprod.113.114116.
- Bak, R. 1976. “The growth of coral colonies and the importance of crustose coralline algae and burrowing sponges in relation with carbonate accumulation.” *Netherlands Journal of Sea Research* 10 (3): 285–337. doi:10.1016/0077-7579(76)90009-0.
- Baker, Andrew C, and Ross Cunning. 2015. “Coral ‘Bleaching’ as a Generalized Stress Response to Environmental Disturbance.” In *Diseases of Coral*, 396–409. Hoboken, NJ: John Wiley & Sons, Inc. doi:10.1002/9781118828502.ch30.
- Baker, P, and Jon N Weber. 1975. “Coral growth rate: Variation with depth.” *Earth and Planetary Science Letters* 27 (1): 57–61. doi:10.1016/0012-821X(75)90160-0.
- Barott, Katie L, Alexander A Venn, Sidney O Perez, Sylvie Tambutte, and Martin Tresguerres. 2014. “Coral host cells acidify symbiotic algal microenvironment to promote photosynthesis.” *Proceedings Of The National Academy Of Sciences Of The United States Of America*, December. National Acad Sciences, 201413483. doi:10.1073/pnas.1413483112.
- Borell, Esther M, Ade Yuliantri, Kai Bischof, and Claudio Richter. 2008. “The effect of heterotrophy on photosynthesis and tissue composition of two scleractinian corals under elevated temperature.” *Journal of Experimental Marine Biology and Ecology* 364: 116–23.
- Brown, Barbara E, R P Dunne, I Ambarsari, Martin Le Tissier, and U Satapoomin. 1999. “Seasonal fluctuations in environmental factors and variations in symbiotic algae and chlorophyll pigments in four Indo-Pacific coral species.” *Marine Ecology Progress Series* 191: 53–69.
- Coles, SL, and Paul L Jokiel. 1978. “Synergistic effects of temperature, salinity and light on the hermatypic coral *Montipora verrucosa*.” *Marine Biology* 49: 187–95. <http://www.springerlink.com/index/LU582837924G7674.pdf>.
- Costanza, R, R de Groot, and P Sutton. 2014. “Changes in the global value of ecosystem services.” *Global Environmental Change* 26: 152–58. doi:10.1016/j.gloenvcha.2014.04.002.
- Cunning, Ross, and Andrew C Baker. 2013. “Excess algal symbionts increase the susceptibility of reef corals to bleaching.” *Nature Climate Change* 3 (March): 259–62. doi:10.1038/nclimate1711.
- Cunning, Ross, Rachel N Silverstein, and Andrew C Baker. 2015. “Investigating the causes and consequences of

- symbiont shuffling in a multi-partner reef coral symbiosis under environmental change.” *Proceedings of the Royal Society B*, 1–9. doi:[10.1098/rspb.2014.1725](https://doi.org/10.1098/rspb.2014.1725){\&}domain=pdf{\&}date\_stamp.
- Downs, C A, Kathleen E McDougall, Cheryl M Woodley, John E Fauth, Robert H Richmond, Ariel Kushmaro, Stuart W Gibb, Yossi Loya, Gary K Ostrander, and Esti Kramarsky-Winter. 2013. “Heat-Stress and Light-Stress Induce Different Cellular Pathologies in the Symbiotic Dinoflagellate during Coral Bleaching.” *PLoS ONE* 8 (12): e77173. doi:[10.1371/journal.pone.0077173](https://doi.org/10.1371/journal.pone.0077173).
- Edmunds, Peter J, Hollie M Putnam, Roger M Nisbet, and Erik B Muller. 2011. “Benchmarks in organism performance and their use in comparative analyses.” *Oecologia* 167 (2): 379–90. doi:[10.1007/s00442-011-2004-2](https://doi.org/10.1007/s00442-011-2004-2).
- Enríquez, Susana, Eugenio R Méndez, and Roberto Iglesias-Prieto. 2005. “Multiple scattering on coral skeletons enhances light absorption by symbiotic algae.” *Limnology and Oceanography* 50 (4): 1025–32.
- Eynaud, Yoan, Roger M Nisbet, and Erik B Muller. 2011. “Impact of excess and harmful radiation on energy budgets in scleractinian corals.” *Ecological Modelling* 222 (7). Elsevier: 1315–22. <http://www.sciencedirect.com/science/article/pii/S0304380011000263>.
- Fagoonee, I, H B Wilson, M P Hassell, and J R Turner. 1999. “The Dynamics of Zooxanthellae Populations: A Long-Term Study in the Field.” *Science* 283 (5403): 843–45. doi:[10.1126/science.283.5403.843](https://doi.org/10.1126/science.283.5403.843).
- Ferrier-Pagès, Christine, Jean-Pierre Gattuso, S Dallot, and Jean Jaubert. 2000. “Effect of nutrient enrichment on growth and photosynthesis of the zooxanthellate coral *Stylophora* . . .” *Coral Reefs*. <http://www.springerlink.com/index/3DRC3K1Q57TPC0UT.pdf>.
- Fitt, William K, F K McFarland, Mark E Warner, and Geoff C Chilcoat. 2000. “Seasonal patterns of tissue biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral bleaching.” *Limnology and Oceanography* 45 (3): 677–85.
- Hawkins, Thomas D, Julia C G Hagemeyer, Kenneth D Hoadley, Adam G Marsh, and Mark E Warner. 2016. “Partitioning of Respiration in an Animal-Algal Symbiosis: Implications for Different Aerobic Capacity between *Symbiodinium* spp.” *Frontiers in Physiology* 7 (April). Frontiers: 125. doi:[10.3389/fphys.2016.00128](https://doi.org/10.3389/fphys.2016.00128).
- Hoegh-Guldberg, Ove. 1999. “Climate change, coral bleaching and the future of the world’s coral reefs.” *Marine and Freshwater Research* 50 (November): 839–66.
- Hopkinson, Brian M, Anna L Tansik, and William K Fitt. 2015. “Internal carbonic anhydrase activity in the tissue of scleractinian corals is sufficient to support proposed roles in photosynthesis and calcification.” *The Journal of Experimental Biology*, April. The Company of Biologists Ltd, jeb.118182. doi:[10.1242/jeb.118182](https://doi.org/10.1242/jeb.118182).
- Huston, M. 1985. “Variation in coral growth rates with depth at Discovery Bay, Jamaica.” *Coral Reefs* 4 (1): 19–25. doi:[10.1007/BF00302200](https://doi.org/10.1007/BF00302200).
- Jager, Tjalling, Benjamin T Martin, and Elke I Zimmer. 2013. “DEBkiss or the quest for the simplest generic model of animal life history.” *Journal of Theoretical Biology* 328: 9–18. doi:[10.1016/j.jtbi.2013.03.011](https://doi.org/10.1016/j.jtbi.2013.03.011).
- Jokiel, Paul L, and SL Coles. 1977. “Effects of temperature on the mortality and growth of Hawaiian reef corals.” *Marine Biology* 43 (3): 201–8.
- Jones, Ross J, Ove Hoegh-Guldberg, Anthony W D Larkum, and Ulrich Schreiber. 1998. “Temperature-induced bleaching of corals begins with impairment of the CO<sub>2</sub> fixation mechanism in zooxanthellae.” *Plant, Cell and Environment* 21 (12): 1219–30.
- Kemp, Dustin W, X Hernandez-Pech, Roberto Iglesias-Prieto, William K Fitt, and Gregory W Schmidt. 2014. “Community dynamics and physiology of *Symbiodinium* spp. before, during, and after a coral bleaching event.” *Limnology and Oceanography*. [http://www.aslo.org/lo/pdf/vol\\_59/issue\\_3/0788.pdf](http://www.aslo.org/lo/pdf/vol_59/issue_3/0788.pdf).
- Kooijman, SALM. 2010. *Dynamic Energy Budget Theory for Metabolic Organization*. 3rd ed. Cambridge University Press.
- Lesser, Michael P. 1997. “Oxidative stress causes coral bleaching during exposure to elevated temperatures.” *Coral Reefs* 16 (July): 187–92.
- Marcelino, Luisa A, Mark W Westneat, Valentina Stoyneva, Jillian Henss, Jeremy D Rogers, Andrew Radosevich,

- Vladimir Turzhitsky, et al. 2013. "Modulation of Light-Enhancement to Symbiotic Algae by Light-Scattering in Corals and Evolutionary Trends in Bleaching." *PLoS ONE* 8 (4): e61492. doi:[10.1371/journal.pone.0061492.s008](https://doi.org/10.1371/journal.pone.0061492.s008).
- Marubini, F, and PS Davies. 1996. "Nitrate increases zooxanthellae population density and reduces skeletogenesis in corals." *Marine Biology* 127 (2): 319–28. <http://www.scopus.com/inward/record.url?partnerID=yv4JPVwI&eid=2-s2.0-0030301761&md5=0971551a6865f30046294b9e17fb4081>.
- Muller, Erik B, Sebastiaan A L M Kooijman, Peter J Edmunds, Francis J Doyle, and Roger M Nisbet. 2009. "Dynamic energy budgets in syntrophic symbiotic relationships between heterotrophic hosts and photoautotrophic symbionts." *Journal of Theoretical Biology* 259 (1): 44–57. doi:[10.1016/j.jtbi.2009.03.004](https://doi.org/10.1016/j.jtbi.2009.03.004).
- Muller-Parker, Gisèle. 1987. "Seasonal variation in light-shade adaptation of natural populations of the symbiotic sea anemone *Aiptasiapulchella* (Carlgren, 1943) in Hawaii." *Journal of Experimental Marine Biology and Ecology* 112 (2): 165–83. doi:[10.1016/0022-0981\(87\)90115-8](https://doi.org/10.1016/0022-0981(87)90115-8).
- Muller-Parker, Gisèle, Lawrence R McCloskey, Ove Hoegh-Guldberg, and PJ McAuley. 1994. "Effect of ammonium enrichment on animal and algal biomass of the coral *Pocillopora damicornis*." *Pacific Science* 48 (3). <http://scholarspace.manoa.hawaii.edu/handle/10125/2236>.
- Muscantine, Leonard, and James W Porter. 1977. "Reef corals: mutualistic symbioses adapted to nutrient-poor environments." *Bioscience* 27 (7): 454–60.
- Muscantine, Leonard, L R McCloskey, and R E Marian. 1981. "Estimating the daily contribution of carbon from zooxanthellae to coral animal respiration." *Limnology and Oceanography* 26 (4): 601–11. doi:[10.4319/lo.1981.26.4.0601](https://doi.org/10.4319/lo.1981.26.4.0601).
- Osinga, Ronald, Miriam Schutter, Ben Griffioen, René H Wijffels, Johan A J Verreth, Shai Shafir, Stéphane Henard, Maura Taruffi, Claudia Gili, and Silvia Lavorano. 2011. "The Biology and Economics of Coral Growth." *Marine Biotechnology* 13 (4): 658–71. doi:[10.1007/s10126-011-9382-7](https://doi.org/10.1007/s10126-011-9382-7).
- Osinga, Ronald, Miriam Schutter, Tim Wijgerde, Buki Rinkevich, Shai Shafir, Muki Shpigel, Gian Marco Luna, et al. 2012. "The CORALZOO project: a synopsis of four years of public aquarium science." *Journal of the Marine Biological Association of the UK* 92 (04). Cambridge University Press: 753–68. doi:[10.1017/S0025315411001779](https://doi.org/10.1017/S0025315411001779).
- Powles, Stephen B. 1984. "Photoinhibition of photosynthesis induced by visible light." *Annual Review of Plant Physiology* 35: 15–44.
- R Core Team. 2014. "R: A Language and Environment for Statistical Computing." Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>.
- Roth, M S. 2014. "The engine of the reef: Photobiology of the coral-algal symbiosis." *Frontiers in Microbiology*. <http://journal.frontiersin.org/Journal/10.3389/fmicb.2014.00422/pdf>.
- Schutter, M, J Crocker, A Paijmans, M Janse, R Osinga, A J Verreth, and R H Wijffels. 2010. "The effect of different flow regimes on the growth and metabolic rates of the scleractinian coral *Galaxea fascicularis*." *Coral Reefs*, April. doi:[10.1007/s00338-010-0617-2](https://doi.org/10.1007/s00338-010-0617-2).
- Schutter, Miriam, Rosa M van der Ven, Max Janse, Johan A J Verreth, René H Wijffels, and Ronald Osinga. 2011. "Light intensity, photoperiod duration, daily light flux and coral growth of *Galaxea fascicularis* in an aquarium setting: a matter of photons?" *Journal of the Marine Biological Association of the UK* 92 (04). Cambridge University Press: 703–12. doi:[10.1017/S0025315411000920](https://doi.org/10.1017/S0025315411000920).
- Shantz, Andrew A, Nathan P Lemoine, and Deron E Burkepile. 2015. "Nutrient loading alters the performance of key nutrient exchange mutualisms." *Ecology Letters* 19 (1): 20–28. doi:[10.1111/ele.12538](https://doi.org/10.1111/ele.12538).
- Stimson, J. 1997. "The annual cycle of density of zooxanthellae in the tissues of field and laboratory-held *Pocillopora damicornis* (Linnaeus)." *Journal of Experimental Marine Biology and Ecology* 214 (1-2): 35–48.
- Tanaka, Yasuaki, Akira Iguchi, Mayuri Inoue, Chiharu Mori, K Sakai, Atsushi Suzuki, Hodaka Kawahata, and Takashi Nakamura. 2013. "Marine Pollution Bulletin." *Marine Pollution Bulletin* 68 (1-2): 93–98. doi:[10.1016/j.marpolbul.2012.12.017](https://doi.org/10.1016/j.marpolbul.2012.12.017).
- Tanaka, Yasuaki, Toshihiro Miyajima, Isao Koike, Takeshi Hayashibara, and Hiroshi Ogawa. 2007. "Imbalanced

- coral growth between organic tissue and carbonate skeleton caused by nutrient enrichment.” *Limnology and Oceanography*. [http://www.aslo.org/lo/toc/vol\\_52/issue\\_3/1139.pdf](http://www.aslo.org/lo/toc/vol_52/issue_3/1139.pdf).
- Tansik, Anna L, William K Fitt, and Brian M Hopkinson. 2015. “External carbonic anhydrase in three Caribbean corals: quantification of activity and role in CO<sub>2</sub> uptake.” *Coral Reefs* 34 (3). Springer Berlin Heidelberg: 703–13. doi:[10.1007/s00338-015-1289-8](https://doi.org/10.1007/s00338-015-1289-8).
- Titlyanov, E A, T V Titlyanova, K Yamazato, and Robert van Woesik. 2001. “Photo-acclimation dynamics of the coral *Stylophora pistillata* to low and extremely low light.” *Journal of Experimental Marine Biology and Ecology* 263 (2): 211–25.
- Vega Thurber, Rebecca L, Deron E Burkepile, Corinne Fuchs, Andrew A Shantz, Ryan McMinds, and Jesse R Zaneveld. 2014. “Chronic nutrient enrichment increases prevalence and severity of coral disease and bleaching.” *Global Change Biology* 20 (2): 544–54. doi:[10.1111/gcb.12450](https://doi.org/10.1111/gcb.12450).
- Wang, J, and Angela E Douglas. 1998. “Nitrogen recycling or nitrogen conservation in an alga-invertebrate symbiosis?” *The Journal of Experimental Biology* 201: 2445–53.
- Warner, Mark E, William K Fitt, and Gregory W Schmidt. 1999. “Damage to photosystem II in symbiotic dinoflagellates: A determinant of coral bleaching.” *Proceedings Of The National Academy Of Sciences Of The United States Of America* 96 (14): 8007–12.
- Weis, Virginia M. 2008. “Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of symbiosis.” *The Journal of Experimental Biology* 211 (Pt 19): 3059–66. doi:[10.1242/jeb.009597](https://doi.org/10.1242/jeb.009597).
- Wooldridge, Scott A. 2009. “A new conceptual model for the warm-water breakdown of the coral-algae endosymbiosis.” *Marine and Freshwater Research* 60 (June): 483–96.
- . 2010. “Is the coral-algae symbiosis really ‘mutually beneficial’ for the partners?” *BioEssays*, May.
- . 2013. “Breakdown of the coral-algae symbiosis: towards formalising a linkage between warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae.” *Biogeosciences* 10: 1647–58. <http://www.biogeosciences.net/10/1647/2013/bg-10-1647-2013.html>.
- . 2014a. “Differential thermal bleaching susceptibilities amongst coral taxa: re-posing the role of the host.” *Coral Reefs* 33 (1). Springer Berlin Heidelberg: 15–27. doi:[10.1007/s00338-013-1111-4](https://doi.org/10.1007/s00338-013-1111-4).
- . 2014b. “Formalising a mechanistic linkage between heterotrophic feeding and thermal bleaching resistance.” *Coral Reefs*. Springer Berlin Heidelberg, 1–6. doi:[10.1007/s00338-014-1193-7](https://doi.org/10.1007/s00338-014-1193-7).
- Wooldridge, Scott A, and Terry J Done. 2009. “Improved water quality can ameliorate effects of climate change on corals.” *Ecological Applications* 19 (6). Ecological Society of America: 1492–9. doi:[10.1890/08-0963.1](https://doi.org/10.1890/08-0963.1).
- Yellowlees, David, T A V Rees, and William Leggat. 2008. “Metabolic interactions between algal symbionts and invertebrate hosts.” *Plant, Cell and Environment* 31: 679–94.



## List of Figures

- 1 Graphical representation of coral-algal symbiosis model. Light, CO<sub>2</sub>, prey, and DIN are acquired from the external environment proportional to the biomass of the partner indicated by the black box for uptake. Mass fluxes (see Table 1 for definitions) are represented by  $j$ 's with subscripts indicating the type of mass, and in some cases the process (e.g.,  $j_{CP}$  is the flux of carbon produced by photosynthesis), and  $\rho$ 's indicate fluxes that are shared by one partner with the other. Parallel complementary synthesizing units (SUs) are represented by large circles, and rejection fluxes from these SUs are indicated by dashed lines.  $c_{ROS}$  is a proportional rate that impacts other model fluxes by inhibition or acceleration; likewise,  $j_{eC}$  accelerates the rate of  $j_{CO_2}$ . Recycled mass fluxes from biomass turnover are not shown for clarity (but see Table 1 for definitions). . . . . 17
- 2 Steady state values of **A)** specific growth (Cmol Cmol<sup>-1</sup> d<sup>-1</sup>), **B)** the symbiont to host biomass ratio (CmolS CmolH<sup>-1</sup>), **C)** relative C- or N-limitation of host biomass formation, and **D)** relative CO<sub>2</sub>- or light-limitation of symbiont photosynthesis, across gradients of external irradiance and dissolved inorganic nitrogen. Note that typical conditions for reefs are ~1e-7 M DIN and 10-20 mol photons m<sup>-2</sup> d<sup>-1</sup>. Simulations for each combination of light and nutrients (41 points along each axis) were run for 100 days with a time step of 1 day. Negative steady state growth rates, and corresponding S:H ratios, were set to zero. . . . . 18
- 3 Sensitivity analysis. Plots show the fractional change in steady state values of growth (solid lines) and S:H biomass (dashed lines) in response to fractional changes in default parameter values (see Table 2 for default values). Parameters are grouped by which processes they are involved in. This sensitivity analysis was conducted at conditions typical for coral reef environments: low DIN (1e-7 M) and intermediate light (15 mol m<sup>-2</sup> d<sup>-2</sup>). Sensitivity analyses conducted other environmental conditions are presented in Figs S3-S7. . . . . 19
- 4 Figure 4. Light-driven seasonal dynamics of symbiont abundance and coral growth. Light input (upper panel) was designed as a sinusoidal curve with a period of one year, with maximum and minimum values of 44 and 22 mol photons m<sup>-2</sup> d<sup>-1</sup>, based on Stimson (1997). The dynamic behavior of symbiont to host biomass ratio (middle panel) and the specific growth rate of host biomass (lower panel) show seasonal oscillations that are greater in magnitude under higher nutrients (1e-5 molN L<sup>-1</sup>; dashed lines) relative to lower nutrients (1e-7 molN L<sup>-1</sup>; solid lines), consistent with the findings of Stimson (1997). Prey density was set at 1e-5 CmolX L<sup>-1</sup>. . . . . 20
- 5 Figure 5. Coral bleaching as a switch from a nitrogen- to carbon-limited alternative stable state. This transition is demonstrated in response to gradually increasing external light (A), which causes an increase in production of ROS (B) that reduces the photosynthetic rate through photoinhibition (C). Decreasing photosynthesis moves the system from overall nitrogen-limitation toward carbon-limitation (D); when this threshold is crossed, the system rapidly becomes highly carbon-limited as photosynthesis becomes CO<sub>2</sub>-limited (E) and symbiont densities rapidly decline (F) into a bleached state. (All parameters at default values; external DIN=1e-7 molN/L, prey density=0). . . . . 21
- 6 Figure 6. Bleaching with interactive factors. Simulations of high light stress (sinusoid with maximum=48 mol ph m<sup>-2</sup> d<sup>-1</sup>) under default environmental conditions (solid line; DIN=1e-7 molN/L; prey=3e-6 CmolX/L), or with elevated feeding (dashed line; prey=6e-6 CmolX/L), or elevated nutrients (dotted line; DIN=1e-6 molN/L). Initial symbiont biomass was set to the steady state for each set of starting conditions, respectively. (All parameters set to default values). . . . 22
- 7 Figure 7. Hysteresis in coral bleaching and recovery. The symbiont:host biomass ratio is plotted as a function of light from a simulation of linearly increasing, then decreasing, external light. The direction of the trajectory is indicated with arrowheads evenly spaced with respect to time. The rate of change in light was slow (0.01 mol ph d<sup>-1</sup>) such that dynamics track closely to steady state values. (All parameters at default values; DIN=1e-7 molN/L; Prey=2e-6 CmolX/L). . . . . 23







Figure 1: Graphical representation of coral-algal symbiosis model. Light, CO<sub>2</sub>, prey, and DIN are acquired from the external environment proportional to the biomass of the partner indicated by the black box for uptake. Mass fluxes (see Table 1 for definitions) are represented by  $j$ 's with subscripts indicating the type of mass, and in some cases the process (e.g.,  $j_{CP}$  is the flux of carbon produced by photosynthesis), and  $\rho$ 's indicate fluxes that are shared by one partner with the other. Parallel complementary synthesizing units (SUs) are represented by large circles, and rejection fluxes from these SUs are indicated by dashed lines.  $c_{ROS}$  is a proportional rate that impacts other model fluxes by inhibition or acceleration; likewise,  $j_{eC}$  accelerates the rate of  $j_{CO_2}$ . Recycled mass fluxes from biomass turnover are not shown for clarity (but see Table 1 for definitions).

### A. Specific growth



### B. Symbiont:host biomass




Figure 2: Steady state values of **A**) specific growth ( $\text{Cmol Cmol}^{-1} \text{d}^{-1}$ ), **B**) the symbiont to host biomass ratio ( $\text{CmolS CmolH}^{-1}$ ), **C**) relative C- or N-limitation of host biomass formation, and **D**) relative  $\text{CO}_2$ - or light-limitation of symbiont photosynthesis, across gradients of external irradiance and dissolved inorganic nitrogen. Note that typical conditions for reefs are  $\sim 1\text{e-}7$  M DIN and  $10\text{-}20$  mol photons  $\text{m}^{-2} \text{d}^{-1}$ . Simulations for each combination of light and nutrients (41 points along each axis) were run for 100 days with a time step of 1 day. Negative steady state growth rates, and corresponding S:H ratios, were set to zero.




Figure 3: Sensitivity analysis. Plots show the fractional change in steady state values of growth (solid lines) and S:H biomass (dashed lines) in response to fractional changes in default parameter values (see Table 2 for default values). Parameters are grouped by which processes they are involved in. This sensitivity analysis was conducted at conditions typical for coral reef environments: low DIN ( $1\text{e-}7\text{ M}$ ) and intermediate light ( $15\text{ mol m}^{-2}\text{ d}^{-2}$ ). Sensitivity analyses conducted other environmental conditions are presented in Figs S3-S7.




Figure 4: Figure 4. Light-driven seasonal dynamics of symbiont abundance and coral growth. Light input (upper panel) was designed as a sinusoidal curve with a period of one year, with maximum and minimum values of 44 and 22 mol photons m<sup>-2</sup> d<sup>-1</sup>, based on Stimson (1997). The dynamic behavior of symbiont to host biomass ratio (middle panel) and the specific growth rate of host biomass (lower panel) show seasonal oscillations that are greater in magnitude under higher nutrients (1e-5 molN L<sup>-1</sup>; dashed lines) relative to lower nutrients (1e-7 molN L<sup>-1</sup>; solid lines), consistent with the findings of Stimson (1997). Prey density was set at 1e-5 CmolX L<sup>-1</sup>.




Figure 5: Figure 5. Coral bleaching as a switch from a nitrogen- to carbon-limited alternative stable state. This transition is demonstrated in response to gradually increasing external light (A), which causes an increase in production of ROS (B) that reduces the photosynthetic rate through photoinhibition (C). Decreasing photosynthesis moves the system from overall nitrogen-limitation toward carbon-limitation (D); when this threshold is crossed, the system rapidly becomes highly carbon-limited as photosynthesis becomes CO<sub>2</sub>-limited (E) and symbiont densities rapidly decline (F) into a bleached state. (All parameters at default values; external DIN=1e-7 molN/L, prey density=0).




Figure 6: Figure 6. Bleaching with interactive factors. Simulations of high light stress (sinusoid with maximum= $48 \text{ mol ph m}^{-2} \text{ d}^{-1}$ ) under default environmental conditions (solid line;  $\text{DIN}=1\text{e-}7 \text{ molN/L}$ ;  $\text{prey}=3\text{e-}6 \text{ CmolX/L}$ ), or with elevated feeding (dashed line;  $\text{prey}=6\text{e-}6 \text{ CmolX/L}$ ), or elevated nutrients (dotted line;  $\text{DIN}=1\text{e-}6 \text{ molN/L}$ ). Initial symbiont biomass was set to the steady state for each set of starting conditions, respectively. (All parameters set to default values).




Figure 7: Figure 7. Hysteresis in coral bleaching and recovery. The symbiont:host biomass ratio is plotted as a function of light from a simulation of linearly increasing, then decreasing, external light. The direction of the trajectory is indicated with arrowheads evenly spaced with respect to time. The rate of change in light was slow ( $0.01 \text{ mol ph d}^{-1}$ ) such that dynamics track closely to steady state values. (All parameters at default values;  $\text{DIN}=1\text{e-}7 \text{ molN/L}$ ;  $\text{Prey}=2\text{e-}6 \text{ CmolX/L}$ ).