Prodigy Anesthesia Quick Review

AIDS/HIV

To what family of viruses does HIV belong?	Retroviridae (retrovirus) *Barash PG, Cullen BF, Stoelting RK, Cahalan MK, Stock MC, Ortega R. Clinical Anesthesia. 7th ed. Philadelphia, PA: Lippincott Williams and Wilkins; 2013: 423-424.*
What type of lymphocyte is destroyed by the AIDS virus?	T-helper lymphocytes (also called CD4 T cells) *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 485.*
Where are T-helper lymphocytes predominantly located?	98% of all T-helper lymphocytes are located within the lymph nodes. *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 485.*
What is the most common opportunistic pathogen associated with AIDS?	Pneumocystis jiroveci (formerly Pneumocystis carinii) *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1029.*
What are the odds of seroconversion from an HIV contaminated needlestick?	0.4% *Miller RD. Miller's Anesthesia. 8th ed. Philadelphia: Elsevier Saunders; 2015: 1081.*
What form of pneumonia is associated with AIDS?	Pneumocystis jiroveci (formerly known as pneumocystic carinii) pneumonia *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 1029.*
Pneumocystic carinii pneumonia usually doesn't develop until the CD4 T cell count reaches what level?	200 cell/mL *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 486.*
What are the potential respiratory complications from pneumocystis carinii pneumonia?	Breathlessness, night sweats, bacterial lung abscesses, tuberculosis, fungal infections, pneumothorax, pulmonary Kaposi's sarcoma, and respiratory failure. Pulmonary adenopathy can be so severe that it results in tracheobronchial and pulmonary vessel compression. Kaposi's sarcoma within the lungs can result in massive hemoptysis. HIV can also lead to an emphysematous destruction of the alveolar tissue. *Stoelting RK, Dierdorf SF. Anesthesia & Co-Existing Diseases. 5th ed. New York, NY: Churchill-Livingston; 2008: 489.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.

Prodigy Anesthesia Quick Review

AIDS/HIV

How does the chest xray of a patient with pneumocystis carinii pneumonia appear?	It often appears normal. A 'ground-glass' appearance may be apparent on xray, but often requires high-resolution CT to identify. Pneumothoraces may sometime be present. *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 487.*
A patient with AIDS has a positive acid-fast bacillus test. What does this signify?	The patient is positive for tuberculosis. *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 487.*
What are the potential cardiac complications of AIDS?	An echocardiogram will be abnormal in about 50% of AIDS patients. Approximately 25% will exhibit pericardial effusions. Infections from opportunistic organisms such as cryptococcus, coxsackie B virus, cytomegalovirus, toxoplasmosis, and aspergillus can cause myocarditis with ventricular dilation. *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 486.*
With relation to HIV, what does the acronym HAART stand for?	Highly active antiretroviral therapy, which represents the current drug regimen used to prevent the advancement of HIV infection into AIDS. *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 484.*
An HIV-positive patient takes a non-nucleoside reverse transcriptase inhibitor. How might this affect your anesthetic management plan?	All of the NNRTIs can prolong the half-life and effects of drugs such as diazepam, midazolam, fentanyl, and meperidine. *Hines RL, Marschall KE. Anesthesia & Co-Existing Diseases. 6th ed. Philadelphia, PA: Elsevier-Saunders; 2012: 488.*
What are the five major classes of antiretroviral drugs used in the treatment of HIV?	Nucleoside analogue reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, protease inhibitors, entry inhibitors, and integrase inhibitors. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1028.*
What is zidovudine and what is its principal side effect?	Zidovudine is a nucleoside analogue reverse transcriptase inhibitor used in the treatment of HIV. Its principal side effect is bone marrow suppression. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1028.*
An anesthetist suffers a needle-stick that was contaminated with blood from an HIV-positive patient. According to current recommendations, how soon should a postexposure prophylaxis medication regimen be started?	Postexposure prophylaxis should be started within 1-2 hours if possible.. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1030.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.