Prodigy Anesthesia Quick Review

Airway Block

What nerve provides sensation to the larynx below the vocal cords?	The recurrent laryngeal nerve provides sensation to the larynx below the vocal cords. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 310.*
What nerve provides sensory innervation to the airway between the epiglottis and the vocal cords?	The airway between the epiglottis and the vocal cords derives sensory innervation from the internal branch of the superior laryngeal nerve. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 310.*
What nerves provide sensation to the mucous membranes of the nose?	The mucous membranes of the nose are innervated by the opthalmic division of the trigeminal nerve (V1) anteriorly and the maxillary division of the trigeminal nerve (V2) posteriorly. These nerves are also known as the anterior ethmoidal and sphenopalatine nerves respectively. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 311.*
What is the only laryngeal muscle that does not receive its motor innervation from the recurrent laryngeal nerve?	The cricothyroid muscle, which is innervated by the external laryngeal nerve. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 310.*
How is a glossopharyngeal block performed?	The glossopharyngeal block is performed by having the patient open their mouth, then using a 25-gauge needle to inject 2 mL of local anesthetic bilaterally at the base of the palatoglossal arch (also called the anterior tonsillar pillar). This blocks the lingual and some of the pharyngeal branches of the glossopharyngeal nerve that provide sensation to the posterior third of the tongue and oropharynx. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1064-1065.*
How is a transtracheal block performed?	A transtracheal block is performed by injecting 4 mL of 4% lidocaine through a needle penetrating the cricothyroid membrane into the trachea. The needle is advanced until air is aspirated. At the end of expiration, the lidocaine is injected. This will usually precipitate a cough, which will spread the anesthetic throughout the airway. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 338.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.

Prodigy Anesthesia Quick Review

Airway Block

How would you anesthetize the posterior 1/3 of the tongue?	The posterior 1/3 of the tongue and tonsillar pillars can be anesthetized by blocking the lingual and pharyngeal branches of the glossopharyngeal nerve with 2 mL of local anesthetic at the base of the palatoglossal arch. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1064-1065.*
How would you anesthetize the superior laryngeal nerve?	Injection of 3 mL at the cornu on either side of the hyoid bone anesthetizes the superior laryngeal nerve, which supplies sensation to the airway below the epiglottis. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 338-339.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.