Prodigy Anesthesia Quick Review

Ankle Block

What nerves are blocked when performing an ankle block?	Deep peroneal, Saphenous, Posterior tibial, Sural, Superficial peroneal *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1015-1016.*
When performing an ankle block, how is the deep peroneal nerve anesthetized? What other nerves can be anesthetized at the same time?	By injecting 5-8ml of local anesthetic in the groove located proximally by the extensor hallucis longus tendon and extensor digitorum longus tendon. By redirecting the needle toward the lateral malleolus and injecting another 5-8 mL, the superficial peroneal nerve can be blocked. Redirecting again toward the medial malleolus and injecting another 5-8 mL, the saphenous nerve can be anesthetized. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1015-1017.*
How would you anesthetize the tibial nerve as part of an ankle block?	Injecting 5-8 mL of local anesthetic in a fanlike manner posterior to the medial malleolus. The posterior tibial and deep peroneal nerves are the only two nerves anesthetized during an ankle block that are not in the subcutaneous tissue. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1015-1017.*
How would you anesthetize the sural nerve as part of an ankle block?	The injection of 5-8 mL of local anesthetic posterior to the lateral malleolus will block the sural nerve. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1015-1017.*
The injection of local anesthetic subcutaneously from the medial malleolus of the ankle to the extensor digitorum longus tendons would anesthetize which two nerves?	The injection of local anesthetic superficially near the medial malleolus will anesthetize the saphenous nerve while the continued injection toward the extensor digitorum longus tendon will anesthetize the superficial peroneal nerve. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1015-1017.*
When doing an ankle block, the injection of local anesthetic between the flexor hallucis longus and the extensor digitorum tendons just lateral to the anterior tibial artery will anesthetize which nerve?	Injecting local anesthetic between the flexor hallucis longus and the extensor digitorum tendons will anesthetize the deep peroneal nerve. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 1015-1017.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.

Ankle Block

The saphenous nerve is the terminal branch of what nerve?	The saphenous nerve is the terminal branch of the femoral nerve and can be blocked by injecting 5 mL of anesthetic anterior to the medial malleolus. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1124.*
Which nerve is crucial to block adequately so that the patient can tolerate an ankle tourniquet?	The saphenous nerve--it innervates the medial side of the malleolus and the medial aspect of the lower leg. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1124.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.