Prodigy Anesthesia Quick Review

Management of Complications

What is the most common injury after anesthesia?	Although injuries from anesthesia are uncommon, the most frequent injury that does occur is corneal abrasion. It can occur during general anesthesia, regional anesthesia, or monitored anesthesia care and requires the anesthetist to pay close attention to the potential for eye injury at all times. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 997.*
What are the risk factors that are associated with an increased risk of postoperative blindness?	Risk factors for the development of postoperative blindness include: intraoperative hypotension, male sex, obesity, anemia from blood loss greater than 1 liters, surgery greater than 5 hours in duration, use of a Wilson frame, and decreased percent colloid administration. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 1010.*
A patient who has numbness to the forehead probably has damage to which nerve?	The supraorbital nerve *Moses KP, Banks JC, Nava PB, Petersen D. Atlas of Clinical Gross Anatomy. Philadelphia, PA: Mosby-Elsevier; 2005: 6-144.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.