# Prodigy Anesthesia Quick Review

## Hemipelvectomy

| | |
|---|---|
| What is a hemipelvectomy? | A hemipelvectomy is an amputation of the entire leg and ipsilateral pelvis.<br>*Molnar R, Emery G, Choong PF. Anaesthesia for hemipelvectomy--a series of 49 cases. Anaesth Intensive Care. 2007 Aug;35(4):536-43.* |
| What are the most common reasons for performing a hemipelvectomy? | Reasons for performing a hemipelvectomy include severe trauma, osteosarcoma, chondrosarcoma, or Ewing's sarcoma.<br>*Molnar R, Emery G, Choong PF. Anaesthesia for hemipelvectomy--a series of 49 cases. Anaesth Intensive Care. 2007 Aug;35(4):536-43.* |
| What are the major anesthetic implications for a patient undergoing a hemipelvectomy? | The primary anesthetic concerns in these patients involve the extensive trauma, blood loss, and postoperative pain associated with the procedure. A study in Australia reported that the mean crystalloid requirement was 8500 mL with an average of 7 units of blood transfused. A combination of epidural/general anesthesia is often implemented. Fluid and forced air warmers are a necessity and the patient can be expected to recover in the intensive care unit.<br>*Molnar R, Emery G, Choong PF. Anaesthesia for hemipelvectomy--a series of 49 cases. Anaesth Intensive Care. 2007 Aug;35(4):536-43.* |

*Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.*