Prodigy Anesthesia Quick Review

Hepatitis

What are the general signs and symptoms of acute viral hepatitis?	There is typically a 1-2 week period of mild symptoms such as fatigue, nausea, fever, and/or vomiting. It may or may not be accompanied by jaundice. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 709.*
What forms of viral hepatitis are associated with fecal-oral contamination of food or water?	Hepatitis A and E, both of which are self-limiting forms of viral hepatitis *Longnecker DE, Newman MF, Brown DL, Zapol WM. Anesthesiology. 2nd ed. New York: McGraw-Hill; 2012: 189-190.*
What forms of viral hepatitis can be transmitted via blood products or percutaneously through the use of shared needles?	Hepatitis B, C, and D *Longnecker DE, Newman MF, Brown DL, Zapol WM. Anesthesiology. 2nd ed. New York: McGraw-Hill; 2012: 189-190.*
What is the incidence of conversion to chronic hepatitis for hepatitis A, B, and C?	Hepatitis A does not convert into chronic hepatitis. As many as 10% of patients with hepatitis B will become chronic and up to 40% of patients with hepatitis C will develop chronic hepatitis. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 769.*
For what forms of viral hepatitis are vaccines available?	Specific vaccines are available for hepatitis A and B. Because hepatitis D only coincides with a hepatitis B infection, it is covered by vaccine as well. *Longnecker DE, Newman MF, Brown DL, Zapol WM. Anesthesiology. 2nd ed. New York: McGraw-Hill; 2012: 189.*
What is the mortality rate for a patient undergoing surgery who is also in alcohol withdrawal?	The mortality rate for these patients approaches 50%. *Butterworth JF, Mackey DC, Wasnick JD. Morgan & Mikhail's Clinical Anesthesiology. 5th ed. New York, NY: McGraw-Hill; 2013: 710.*
How do volatile anesthetics result in hepatitis?	Volatile anesthetic hepatitis is a mild, self-limiting form of hepatitis that is believed to be due to decreases in oxygen supply to the liver relative to demand. There is also some evidence that IgG antibodies form in response to liver proteins modified by the trifluoroacetyl metabolite of halothane which may contribute as a causative factor. *Barash PG, Cullen BF, Stoelting RK, Cahalan MK, Stock MC, Ortega R. Clinical Anesthesia. 7th ed. Philadelphia, PA: Lippincott Williams and Wilkins; 2013: 1302.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.

Prodigy Anesthesia Quick Review

Hepatitis

What is the most common cause of drug-induced hepatitis? What are some other drugs that can result in hepatitis?	Alcohol is the most common cause of drug-induced hepatitis. Other drug-related causes include: acetaminophen, valproic acid, isoniazid, salicylates, azothioprine, ibuprofen, erythromycin, nitrofurantoin, methotrexate, tetracycline, and sulfonamides. *Nagelhout JJ, Plaus KL. Nurse Anesthesia. 5th ed. St. Louis, MO: Elsevier Saunders Company; 2014: 769-771.*
What is the least severe form of viral hepatitis?	Hepatitis A *Longnecker DE, Newman MF, Brown DL, Zapol WM. Anesthesiology. 2nd ed. New York: McGraw-Hill; 2012: 190.*
With what form of viral hepatitis is hepatitis D associated?	Hepatitis D only occurs as a coinfection with hepatitis B. By itself, it does not produce hepatitis. The hepatitis B vaccine, therefore, protects against hepatitis D as well. *Longnecker DE, Newman MF, Brown DL, Zapol WM. Anesthesiology. 2nd ed. New York: McGraw-Hill; 2012: 189-190.*
What body fluids are known to carry and possibly transmit the hepatitis B virus?	All body fluids should be considered potential vectors in the transmission of hepatitis B, including urine, semen, saliva, and blood. *Longnecker DE, Newman MF, Brown DL, Zapol WM. Anesthesiology. 2nd ed. New York: McGraw-Hill; 2012: 190.*

Copyright 2015 Prodigy Anesthesia, Inc. Unlawful to duplicate, distribute, or sell this document.