



# Announcements

- Homework 3 due Monday night
  - Post questions to Campuswire over the weekend!
- I'll be out of the office for much of the afternoon at the SCRP talks
- Monday Reading: Ch 2.3 and Ch 2.4



Evaluate

$$\int_{\text{all space}} (r^2 + \vec{r} \cdot \vec{a} + a^2) \delta^3(\vec{r} - \vec{a}) d\tau$$

- A.  $3a^2$
- B.  $2a^2$
- C.  $a^2$
- D. 0



Evaluate

$$\int_{\text{all space}} (r^2 + \vec{r} \cdot \vec{a} + a^2) \delta^3(\vec{r} - \vec{a}) d\tau$$

- A.  $3a^2$
- B.  $2a^2$
- C.  $a^2$
- D. 0



Evaluate

$$\nabla \cdot \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}}$$

- A.  $\frac{q}{\epsilon_0}$
- B.  $\frac{q}{\epsilon_0} \delta^3(\vec{r})$
- C.  $\frac{q}{\epsilon_0} \hat{\mathbf{r}}$
- D.  $\frac{4\pi q}{\epsilon_0}$




Evaluate

$$\nabla \cdot \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}}$$

- A.  $\frac{q}{\epsilon_0}$
- B.  $\frac{q}{\epsilon_0} \delta^3(\vec{r})$
- C.  $\frac{q}{\epsilon_0} \hat{\mathbf{r}}$
- D.  $\frac{4\pi q}{\epsilon_0}$




Consider a small charged bead pierced by a long straight narrow charged wire, as seen below. Could you use Gauss's Law to solve for the electric field a short distance above the bead?



- A. Yes, just choose the correct surface.
- B. Yes, but it requires multiple surfaces.
- C. Yes, but only if the bead and wire have opposite charges.
- D. No, this doesn't have the needed symmetry.



Consider a small charged bead pierced by a long straight narrow charged wire, as seen below. Could you use Gauss's Law to solve for the electric field a short distance above the bead?



- A. Yes, just choose the correct surface.
- B. Yes, but it requires multiple surfaces.
- C. Yes, but only if the bead and wire have opposite charges.
- D. No, this doesn't have the needed symmetry.



Does superposition apply to electric potential,  $V$ ?

$$V_{tot} \stackrel{?}{=} \sum_i V_i = V_1 + V_2 + V_3 + \dots$$

- A. Yes
- B. No
- C. Sometimes





Does superposition apply to electric potential,  $V$ ?

$$V_{tot} \stackrel{?}{=} \sum_i V_i = V_1 + V_2 + V_3 + \dots$$

- A. Yes
- B. No
- C. Sometimes



The potential is zero at some point in space.  
You can conclude that:

- A. The E-field is zero at that point
- B. The E-field is zero near that point
- C. The E-field is non-zero at that point
- D. You can conclude nothing about the E-field at that point



The potential is zero at some point in space.  
You can conclude that:

- A. The E-field is zero at that point
- B. The E-field is zero near that point
- C. The E-field is non-zero at that point
- D. You can conclude nothing about the E-field at that point



The potential is constant everywhere in some region of space.  
You can conclude that:

- A. The E-field is changing at a constant rate in that space.
- B. The E-field has a constant magnitude in that space.
- C. The E-field is zero in that space.
- D. You can conclude nothing at all about the E-field in that space.



The potential is constant everywhere in some region of space.  
You can conclude that:

- A. The E-field is changing at a constant rate in that space.
- B. The E-field has a constant magnitude in that space.
- C. The E-field is zero in that space.
- D. You can conclude nothing at all about the E-field in that space.




We usually choose the reference point  $\mathcal{O}$  to be 0 at  $\vec{r} = \infty$  when calculating the potential of a point charge. ( $V = kq/r$ ) How does this potential change if we choose our reference point to be  $V(R) = 0$  when  $R$  is close to  $q$ ?

- A.  $V(r)$  is higher than before
- B.  $V(r)$  is lower than before
- C.  $V(r)$  doesn't change
- D. Impossible to say without knowing how close it gets



We usually choose the reference point  $\mathcal{O}$  to be 0 at  $\vec{r} = \infty$  when calculating the potential of a point charge. ( $V = kq/r$ ) How does this potential change if we choose our reference point to be  $V(R) = 0$  when  $R$  is close to  $q$ ?


- A.  $V(r)$  is higher than before
- B.  $V(r)$  is lower than before
- C.  $V(r)$  doesn't change
- D. Impossible to say without knowing how close it gets



Could the above be a plot of  $|\vec{\mathbf{E}}|$ ? or  $V(r)$ ? (Assuming for some physical situation)

- A. Could be  $E(r)$  or  $V(r)$
- B. Could be  $E(r)$ , but not  $V(r)$
- C. Can't be  $E(r)$ , but could be  $V(r)$
- D. Can't be either





Could the above be a plot of  $|\vec{\mathbf{E}}|$ ? or  $V(r)$ ? (Assuming for some physical situation)

- A. Could be  $E(r)$  or  $V(r)$
- B. Could be  $E(r)$ , but not  $V(r)$
- C. Can't be  $E(r)$ , but could be  $V(r)$
- D. Can't be either