

# Announcements

- Homework
  - One final homework 8 to be due next Monday
- No class on Wednesday due to SSRD.
- No class on Friday – work on your projects. I will be at the Chemistry talks.
- Next Monday will probably be another project work day (and I'll be around), but may need to have presentations. I'll let you know soon.
- If you haven't, LET ME KNOW WHO YOU ARE PLANNING TO WORK WITH, EVEN IF YOU ARE WORKING ALONE.
- Feel free to run a dataset by me if you have found one you think would work well for the final project but aren't sure.
- Presentation dates will be randomly assigned. Written portion is due in class on the last day of class.
- Polling: `rembold-class.ddns.net`

## Regression so far ...

At this point we have covered:

- Simple linear regression
  - Relationship between numerical response and a numerical or categorical predictor
- Multiple regression
  - Relationship between numerical response and multiple numerical and/or categorical predictors

What we haven't seen is what to do when the predictors are weird (nonlinear, complicated dependence structure, etc.) or when the response is weird (categorical, count data, etc.)

## Regression so far ...

At this point we have covered:

- Simple linear regression
  - Relationship between numerical response and a numerical or categorical predictor
- Multiple regression
  - Relationship between numerical response and multiple numerical and/or categorical predictors

What we haven't seen is what to do when the predictors are weird (nonlinear, complicated dependence structure, etc.) or when the response is weird (categorical, count data, etc.)

# Odds

Odds are another way of quantifying the probability of an event, commonly used in gambling (and logistic regression).

## Odds

For some event  $E$ ,

$$\text{odds}(E) = \frac{P(E)}{P(E^c)} = \frac{P(E)}{1 - P(E)}$$

Similarly, if we are told the odds of  $E$  are  $x$  to  $y$  then

$$\text{odds}(E) = \frac{x}{y} = \frac{x/(x+y)}{y/(x+y)}$$

which implies

$$P(E) = x/(x+y), \quad P(E^c) = y/(x+y)$$

## Understanding Check

Your bookie tells you that the odds of your horse winning the race are 8:1 or 8 to 1 odds. What is the probability that your horse wins?

- A) 8%
- B) 13%
- C) 81%
- D) 89%

## Understanding Check

Your bookie tells you that the odds of your horse winning the race are 8:1 or 8 to 1 odds. What is the probability that your horse wins?

- A) 8%
- B) 13%
- C) 81%
- D) 89%

## Example - Donner Party

In 1846 the Donner and Reed families left Springfield, Illinois, for California by covered wagon. In July, the Donner Party, as it became known, reached Fort Bridger, Wyoming. There its leaders decided to attempt a new and untested route to the Sacramento Valley. Having reached its full size of 87 people and 20 wagons, the party was delayed by a difficult crossing of the Wasatch Range and again in the crossing of the desert west of the Great Salt Lake. The group became stranded in the eastern Sierra Nevada mountains when the region was hit by heavy snows in late October. By the time the last survivor was rescued on April 21, 1847, 40 of the 87 members had died from famine and exposure to extreme cold.

*Source: Ramsey, F.L. and Schafer, D.W. (2002). The Statistical Sleuth: A Course in Methods of Data Analysis (2nd ed)*

## Example - Donner Party - Data

	Age	Sex	Status
1	23.00	Male	Died
2	40.00	Female	Survived
3	40.00	Male	Survived
4	30.00	Male	Died
5	28.00	Male	Died
⋮	⋮	⋮	⋮
43	23.00	Male	Survived
44	24.00	Male	Died
45	25.00	Female	Survived




# Donner Party - Exploration

Status vs. Gender:

	Male	Female
Died	20	5
Survived	10	10

Status vs. Age:



## Donner Party - Predict Survivors?

It seems clear that both age and gender have an effect on someone's survival, how do we come up with a model that will let us explore this relationship?

Even if we set Died to 0 and Survived to 1, this isn't something we can transform our way out of - we need something more.

One way to think about the problem - we can treat Survived and Died as successes and failures arising from a probability distribution where the probability of a success is given by a transformation of a linear model of the predictors.

## Generalized linear models

It turns out that this is a very general way of addressing this type of problem in regression, and the resulting models are called generalized linear models (GLMs). Logistic regression is just one example of this type of model.

# Generalized linear models

It turns out that this is a very general way of addressing this type of problem in regression, and the resulting models are called generalized linear models (GLMs). Logistic regression is just one example of this type of model.

All generalized linear models have the following three characteristics:

1. A probability distribution describing the outcome variable
2. A linear model
  - $\eta = \beta_0 + \beta_1 X_1 + \cdots + \beta_n X_n$
3. A link function that relates the linear model to the parameter of the outcome distribution
  - $g(p) = \eta$  or  $p = g^{-1}(\eta)$

# Logistic Regression

Logistic regression is a GLM used to model a binary categorical variable using numerical and categorical predictors.

We assume a binomial distribution produced the outcome variable and we therefore want to model  $p$  the probability of success for a given set of predictors.

# Logistic Regression

Logistic regression is a GLM used to model a binary categorical variable using numerical and categorical predictors.

We assume a binomial distribution produced the outcome variable and we therefore want to model  $p$  the probability of success for a given set of predictors.

To finish specifying the Logistic model we just need to establish a reasonable link function that connects  $\eta$  to  $p$ . There are a variety of options but the most commonly used is the logit function.

## Logit function

$$\text{logit}(p) = \log\left(\frac{p}{1-p}\right), \text{ for } 0 \leq p \leq 1$$

## Properties of the Logit

The logit function takes a value between 0 and 1 and maps it to a value between  $-\infty$  and  $\infty$ .

### Inverse logit (logistic) function

$$g^{-1}(x) = \frac{\exp(x)}{1 + \exp(x)} = \frac{1}{1 + \exp(-x)}$$

The inverse logit function takes a value between  $-\infty$  and  $\infty$  and maps it to a value between 0 and 1.

This formulation also has some use when it comes to interpreting the model as logit can be interpreted as the log odds of a success, more on this later.

# The logistic regression model

The three GLM criteria give us:

$$y_i \sim \text{Binom}(p_i)$$

$$\eta = \beta_0 + \beta_1 x_1 + \cdots + \beta_n x_n$$

$$\text{logit}(p) = \eta$$

From which we arrive at,

$$p_i = \frac{\exp(\beta_0 + \beta_1 x_{1,i} + \cdots + \beta_n x_{n,i})}{1 + \exp(\beta_0 + \beta_1 x_{1,i} + \cdots + \beta_n x_{n,i})}$$



## Example - Donner Party - Model

In R we fit a GLM in the same way as a linear model except using `glm` instead of `lm` and we must also specify the type of GLM to fit using the `family` argument.

```
summary(glm(Status ~ Age, data=donner, family=binomial))
```

```
## Call:
```

```
## glm(formula = Status ~ Age, family = binomial, data = donner)
```

```
##
```

```
## Coefficients:
```

```
##           Estimate Std. Error z value Pr(>|z|)
```

```
## (Intercept)  1.81852    0.99937   1.820   0.0688 .
```

```
## Age         -0.06647    0.03222  -2.063   0.0391 *
```

```
##
```

## Donner Party - Prediction

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	1.8185	0.9994	1.82	0.0688
Age	-0.0665	0.0322	-2.06	0.0391

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

## Donner Party - Prediction

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	1.8185	0.9994	1.82	0.0688
Age	-0.0665	0.0322	-2.06	0.0391

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a newborn (Age=0):

## Donner Party - Prediction

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	1.8185	0.9994	1.82	0.0688
Age	-0.0665	0.0322	-2.06	0.0391

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a newborn (Age=0):

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 0$$

$$\frac{p}{1-p} = \exp(1.8185) = 6.16$$

$$p = 6.16/7.16 = 0.86$$

## Example - Donner Party - Prediction (cont.)

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a 25 year old:

## Example - Donner Party - Prediction (cont.)

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a 25 year old:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 25$$

$$\frac{p}{1-p} = \exp(0.156) = 1.17$$

$$p = 1.17/2.17 = 0.539$$

## Example - Donner Party - Prediction (cont.)

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a 25 year old:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 25$$

$$\frac{p}{1-p} = \exp(0.156) = 1.17$$

$$p = 1.17/2.17 = 0.539$$

Odds / Probability of survival for a 50 year old:

## Example - Donner Party - Prediction (cont.)

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a 25 year old:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 25$$

$$\frac{p}{1-p} = \exp(0.156) = 1.17$$

$$p = 1.17/2.17 = 0.539$$

Odds / Probability of survival for a 50 year old:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 50$$


$$\frac{p}{1-p} = \exp(-1.5065) = 0.222$$

$$p = 0.222/1.222 = 0.181$$




# Donner Predictions: Graphically

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$



# Donner Predictions: Graphically

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$



# Donner Party - Age and Gender

```
summary(glm(Status ~ Age + Sex, data=donner, family=binomial))

## Call:
## glm(formula = Status ~ Age + Sex, family = binomial, data = donner)
##
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept)  1.63312    1.11018   1.471   0.1413
## Age         -0.07820    0.03728  -2.097   0.0359 *
## SexFemale    1.59729    0.75547   2.114   0.0345 *
## ---
##
## (Dispersion parameter for binomial family taken to be 1)
```

**Gender slope:** When the other predictors are held constant this is the log odds ratio between the given level (Female) and the reference level (Male).

## Donner Party - Gender Models

Just like MLR we can plug in gender to arrive at two status vs age models for men and women respectively.

General model:

$$\log\left(\frac{p_1}{1-p_1}\right) = 1.63312 + -0.07820 \times \text{Age} + 1.59729 \times \text{Sex}$$

Male model:

$$\begin{aligned}\log\left(\frac{p_1}{1-p_1}\right) &= 1.63312 + -0.07820 \times \text{Age} + 1.59729 \times 0 \\ &= 1.63312 + -0.07820 \times \text{Age}\end{aligned}$$

Female model:

$$\begin{aligned}\log\left(\frac{p_1}{1-p_1}\right) &= 1.63312 + -0.07820 \times \text{Age} + 1.59729 \times 1 \\ &= 3.23041 + -0.07820 \times \text{Age}\end{aligned}$$

# Donner Party: Gender Models Graphically

