

# Announcements

- Homework
  - Homework 7 and Lab due on Monday
- Test a week from today!
  - Over Chapters 2,3,4
  - I'm trying to get study materials posted today or tomorrow
  - I'm also trying hard to have grading done by that point
- Final Project Info sent out today!
  - I'm getting potential datasets uploaded over the weekend.
  - You are always free to find your own as well. Just make sure they have enough variables to give your options for multiple regression.
- Polling: `rembold-class.ddns.net`

## ANOVA in R

Run the ANOVA test on a linear fit to the data:

```
anova(lm(aldrin$aldrin ~ aldrin$depth))
```

Response: aldrin\$aldrin

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
aldrin\$depth	2	16.961	8.4803	6.1338	0.006367 **
Residuals	27	37.329	1.3826		

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1







## Checking Conditions: (1) independence

Does this condition appear to be satisfied?

In this study the we have no reason to believe that the aldrin concentration won't be independent of each other.


## Checking Conditions: (2) approximately normal

Does this condition appear to be satisfied?



## Checking Conditions: (3) constant variance

Does this condition appear to be satisfied?



## Which means differ?

- Earlier we concluded that at least one pair of means differ. The natural question that follows is “which ones?”

## Which means differ?

- Earlier we concluded that at least one pair of means differ. The natural question that follows is “which ones?”
- We can do two-sample  $t$  tests for differences in each possible pair of groups.

## Which means differ?

- Earlier we concluded that at least one pair of means differ. The natural question that follows is “which ones?”
- We can do two-sample  $t$  tests for differences in each possible pair of groups.

Can you see any pitfalls with this approach?



## Which means differ?

- Earlier we concluded that at least one pair of means differ. The natural question that follows is “which ones?”
- We can do two-sample  $t$  tests for differences in each possible pair of groups.

Can you see any pitfalls with this approach?

- When we run too many tests, the Type 1 Error rate increases.
- This issue is resolved by using a modified significance level.

## Multiple comparisons

- The scenario of testing many pairs of groups is called *multiple comparisons*.

## Multiple comparisons

- The scenario of testing many pairs of groups is called **multiple comparisons**.
- The **Bonferroni correction** suggests that a more **stringent** significance level is more appropriate for these tests:

$$\alpha^{\star} = \alpha/K$$

where  $K$  is the number of comparisons being considered.

## Multiple comparisons

- The scenario of testing many pairs of groups is called **multiple comparisons**.
- The **Bonferroni correction** suggests that a more **stringent** significance level is more appropriate for these tests:

$$\alpha^{\star} = \alpha / K$$

where  $K$  is the number of comparisons being considered.

- If there are  $k$  groups, then usually all possible pairs are compared and  $K = \frac{k(k-1)}{2}$ .

## Understanding Check!

In the aldrin data set depth has 3 levels: bottom, mid-depth, and surface. If  $\alpha = 0.05$ , what should be the modified significance level for two sample  $t$  tests for determining which pairs of groups have significantly different means?

- A)  $\alpha^* = 0.05$
- B)  $\alpha^* = 0.05/2 = 0.025$
- C)  $\alpha^* = 0.05/3 = 0.0167$
- D)  $\alpha^* = 0.05/6 = 0.0083$

## Understanding Check!

In the aldrin data set depth has 3 levels: bottom, mid-depth, and surface. If  $\alpha = 0.05$ , what should be the modified significance level for two sample  $t$  tests for determining which pairs of groups have significantly different means?

- A)  $\alpha^* = 0.05$
- B)  $\alpha^* = 0.05/2 = 0.025$
- C)  $\alpha^* = 0.05/3 = 0.0167$
- D)  $\alpha^* = 0.05/6 = 0.0083$

## Which means differ?

If the ANOVA assumption of equal variability across groups is satisfied, we can use the data from all groups to estimate variability:

- Estimate any within-group standard deviation with  $\sqrt{MSE}$ , which is  $s_{pooled}$
- Use the error degrees of freedom,  $n - k$ , for  $t$ -distributions

### Difference in two means: after ANOVA

$$SE = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \approx \sqrt{\frac{MSE}{n_1} + \frac{MSE}{n_2}}$$

## Bottom and Mid Check

Is there a difference between the average aldrin concentration at the bottom and at mid depth?

	n	mean	sd
bottom	10	6.04	1.58
middepth	10	5.05	1.10
surface	10	4.2	0.66
overall	30	5.1	1.37

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
depth	2	16.96	8.48	6.13	0.0063
Residuals	27	37.33	1.38		
Total	29	54.29			



## Bottom and Mid Check

Is there a difference between the average aldrin concentration at the bottom and at mid depth?

	n	mean	sd
bottom	10	6.04	1.58
middepth	10	5.05	1.10
surface	10	4.2	0.66
overall	30	5.1	1.37

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
depth	2	16.96	8.48	6.13	0.0063
Residuals	27	37.33	1.38		
Total	29	54.29			

$$T_{df_E} = \frac{(\bar{X}_{bottom} - \bar{X}_{middepth})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{middepth}}}}$$

$$T_{27} = \frac{(6.04 - 5.05)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{0.99}{0.53} = 1.87$$

## Bottom and Mid Check

Is there a difference between the average aldrin concentration at the bottom and at mid depth?

	n	mean	sd
bottom	10	6.04	1.58
middepth	10	5.05	1.10
surface	10	4.2	0.66
overall	30	5.1	1.37

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
depth	2	16.96	8.48	6.13	0.0063
Residuals	27	37.33	1.38		
Total	29	54.29			

$$T_{df_E} = \frac{(\bar{X}_{bottom} - \bar{X}_{middepth})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{middepth}}}}$$

$$T_{27} = \frac{(6.04 - 5.05)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{0.99}{0.53} = 1.87$$

$$0.05 < p - value < 0.10 \quad (\text{two-sided})$$

$$\alpha^* = 0.05/3 = 0.0167$$

## Bottom and Mid Check

Is there a difference between the average aldrin concentration at the bottom and at mid depth?

	n	mean	sd
bottom	10	6.04	1.58
middepth	10	5.05	1.10
surface	10	4.2	0.66
overall	30	5.1	1.37

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
depth	2	16.96	8.48	6.13	0.0063
Residuals	27	37.33	1.38		
Total	29	54.29			

$$T_{df_E} = \frac{(\bar{X}_{bottom} - \bar{X}_{middepth})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{middepth}}}}$$

$$0.05 < p\text{-value} < 0.10 \quad (\text{two-sided})$$

$$T_{27} = \frac{(6.04 - 5.05)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{0.99}{0.53} = 1.87$$

$$\alpha^* = 0.05/3 = 0.0167$$

Fail to reject  $H_0$ , the data do not provide convincing evidence of a difference between the average aldrin concentrations at bottom and mid depth.

## Bottom and Surface

Is there a difference between the average aldrin concentration at the bottom and at surface?

## Bottom and Surface

Is there a difference between the average aldrin concentration at the bottom and at surface?

$$T_{df_E} = \frac{(\bar{x}_{bottom} - \bar{x}_{surface})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{surface}}}}$$

## Bottom and Surface

Is there a difference between the average aldrin concentration at the bottom and at surface?

$$T_{df_E} = \frac{(\bar{x}_{bottom} - \bar{x}_{surface})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{surface}}}}$$
$$T_{27} = \frac{(6.04 - 4.02)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{2.02}{0.53} = 3.81$$

## Bottom and Surface

Is there a difference between the average aldrin concentration at the bottom and at surface?

$$T_{df_E} = \frac{(\bar{x}_{bottom} - \bar{x}_{surface})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{surface}}}}$$

$$T_{27} = \frac{(6.04 - 4.02)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{2.02}{0.53} = 3.81$$

$$p - value < 0.01 \quad (\text{two-sided})$$

## Bottom and Surface

Is there a difference between the average aldrin concentration at the bottom and at surface?

$$T_{df_E} = \frac{(\bar{x}_{bottom} - \bar{x}_{surface})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{surface}}}}$$

$$T_{27} = \frac{(6.04 - 4.02)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{2.02}{0.53} = 3.81$$

$$p - value < 0.01 \quad (\text{two-sided})$$

$$\alpha^* = 0.05/3 = 0.0167$$



## Bottom and Surface

Is there a difference between the average aldrin concentration at the bottom and at surface?

$$T_{df_E} = \frac{(\bar{x}_{bottom} - \bar{x}_{surface})}{\sqrt{\frac{MSE}{n_{bottom}} + \frac{MSE}{n_{surface}}}}$$

$$T_{27} = \frac{(6.04 - 4.02)}{\sqrt{\frac{1.38}{10} + \frac{1.38}{10}}} = \frac{2.02}{0.53} = 3.81$$

$$p - value < 0.01 \quad (\text{two-sided})$$

$$\alpha^* = 0.05/3 = 0.0167$$


Reject  $H_0$ , the data provide convincing evidence of a difference between the average aldrin concentrations at bottom and surface.

## **Back to Linear Regression!**

---

## Nature or nurture?

In 1966 Cyril Burt published a paper called “The genetic determination of differences in intelligence: A study of monozygotic twins reared apart?” The data consist of IQ scores for [an assumed random sample of] 27 identical twins, one raised by foster parents, the other by the biological parents.



## Testing for the slope

Assuming that these 27 twins comprise a representative sample of all twins separated at birth, we would like to test if these data provide convincing evidence that the IQ of the biological twin is a significant predictor of IQ of the foster twin. What are the appropriate hypotheses?

- (a)  $H_0 : b_0 = 0$ ;  $H_A : b_0 \neq 0$
- (b)  $H_0 : \beta_0 = 0$ ;  $H_A : \beta_0 \neq 0$
- (c)  $H_0 : b_1 = 0$ ;  $H_A : b_1 \neq 0$
- (d)  $H_0 : \beta_1 = 0$ ;  $H_A : \beta_1 \neq 0$

*Remember:* We use  $\beta_0$  and  $\beta_1$  for the population intercept and slope and  $b_0$  and  $b_1$  for our samples intercept and slope.

## Testing for the slope

Assuming that these 27 twins comprise a representative sample of all twins separated at birth, we would like to test if these data provide convincing evidence that the IQ of the biological twin is a significant predictor of IQ of the foster twin. What are the appropriate hypotheses?

- (a)  $H_0 : b_0 = 0$ ;  $H_A : b_0 \neq 0$
- (b)  $H_0 : \beta_0 = 0$ ;  $H_A : \beta_0 \neq 0$
- (c)  $H_0 : b_1 = 0$ ;  $H_A : b_1 \neq 0$
- (d)  $H_0 : \beta_1 = 0$ ;  $H_A : \beta_1 \neq 0$

*Remember:* We use  $\beta_0$  and  $\beta_1$  for the population intercept and slope and  $b_0$  and  $b_1$  for our samples intercept and slope.

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

- We always use a  $t$ -test in inference for regression.

*Remember:* Test statistic,  $T = \frac{\text{point estimate} - \text{null value}}{SE}$

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

- We always use a  $t$ -test in inference for regression.

*Remember:* Test statistic,  $T = \frac{\text{point estimate} - \text{null value}}{SE}$

- Point estimate =  $b_1$  is the observed slope.



## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

- We always use a  $t$ -test in inference for regression.

*Remember:* Test statistic,  $T = \frac{\text{point estimate} - \text{null value}}{SE}$

- Point estimate =  $b_1$  is the observed slope.
- $SE_{b_1}$  is the standard error associated with the slope.

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

- We always use a  $t$ -test in inference for regression.

*Remember:* Test statistic,  $T = \frac{\text{point estimate} - \text{null value}}{SE}$

- Point estimate =  $b_1$  is the observed slope.
- $SE_{b_1}$  is the standard error associated with the slope.
- Degrees of freedom associated with the slope is  $df = n - 2$ , where  $n$  is the sample size.

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
biolQ	0.9014	0.0963	9.36	0.0000

- We always use a  $t$ -test in inference for regression.

*Remember:* Test statistic,  $T = \frac{\text{point estimate} - \text{null value}}{SE}$

- Point estimate =  $b_1$  is the observed slope.
- $SE_{b_1}$  is the standard error associated with the slope.
- Degrees of freedom associated with the slope is  $df = n - 2$ , where  $n$  is the sample size.

*Remember:* We lose 1 degree of freedom for each parameter we estimate, and in simple linear regression we estimate 2 parameters,  $\beta_0$  and  $\beta_1$ .

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

$$T = \frac{0.9014 - 0}{0.0963} = 9.36$$

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
biolQ	0.9014	0.0963	9.36	0.0000

$$T = \frac{0.9014 - 0}{0.0963} = 9.36$$
$$df = 27 - 2 = 25$$

## Testing for the slope (cont.)

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
biolQ	0.9014	0.0963	9.36	0.0000

$$T = \frac{0.9014 - 0}{0.0963} = 9.36$$

$$df = 27 - 2 = 25$$

$$p\text{-value} = P(|T| > 9.36) < 0.01$$

## Confidence interval for the slope

Remember that a confidence interval is calculated as *point estimate*  $\pm$  *ME* and the degrees of freedom associated with the slope in a simple linear regression is  $n - 2$ . Which of the below is the correct 95% confidence interval for the slope parameter? Note that the model is based on observations from 27 twins.

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

- (a)  $9.2076 \pm 1.65 \times 9.2999$
- (b)  $0.9014 \pm 2.06 \times 0.0963$
- (c)  $0.9014 \pm 1.96 \times 0.0963$
- (d)  $9.2076 \pm 1.96 \times 0.0963$



## Confidence interval for the slope

Remember that a confidence interval is calculated as *point estimate*  $\pm$  *ME* and the degrees of freedom associated with the slope in a simple linear regression is  $n - 2$ . Which of the below is the correct 95% confidence interval for the slope parameter? Note that the model is based on observations from 27 twins.

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.2076	9.2999	0.99	0.3316
bioIQ	0.9014	0.0963	9.36	0.0000

- (a)  $9.2076 \pm 1.65 \times 9.2999$
- (b)  $0.9014 \pm 2.06 \times 0.0963$
- (c)  $0.9014 \pm 1.96 \times 0.0963$
- (d)  $9.2076 \pm 1.96 \times 0.0963$

## Recap

- Inference for the slope for a single-predictor linear regression model:

## Recap

- Inference for the slope for a single-predictor linear regression model:
  - Hypothesis test:

$$T = \frac{b_1 - \text{null value}}{SE_{b_1}} \quad df = n - 2$$

## Recap

- Inference for the slope for a single-predictor linear regression model:
  - Hypothesis test:

$$T = \frac{b_1 - \text{null value}}{SE_{b_1}} \quad df = n - 2$$

- Confidence interval:

$$b_1 \pm t_{df=n-2}^* SE_{b_1}$$

## Recap

- Inference for the slope for a single-predictor linear regression model:
  - Hypothesis test:

$$T = \frac{b_1 - \text{null value}}{SE_{b_1}} \quad df = n - 2$$

- Confidence interval:

$$b_1 \pm t_{df=n-2}^* SE_{b_1}$$

- The null value is often 0 since we are usually checking for **any** relationship between the explanatory and the response variable.

## Recap

- Inference for the slope for a single-predictor linear regression model:
  - Hypothesis test:

$$T = \frac{b_1 - \text{null value}}{SE_{b_1}} \quad df = n - 2$$

- Confidence interval:

$$b_1 \pm t_{df=n-2}^* SE_{b_1}$$

- The null value is often 0 since we are usually checking for **any** relationship between the explanatory and the response variable.
  - The regression output gives  $b_1$ ,  $SE_{b_1}$ , and **two-tailed** p-value for the  $t$ -test for the slope where the null value is 0.

## Recap

- Inference for the slope for a single-predictor linear regression model:
  - Hypothesis test:

$$T = \frac{b_1 - \text{null value}}{SE_{b_1}} \quad df = n - 2$$

- Confidence interval:

$$b_1 \pm t_{df=n-2}^* SE_{b_1}$$

- The null value is often 0 since we are usually checking for **any** relationship between the explanatory and the response variable.
- The regression output gives  $b_1$ ,  $SE_{b_1}$ , and **two-tailed** p-value for the  $t$ -test for the slope where the null value is 0.
- We rarely do inference on the intercept, so we'll be focusing on the estimates and inference for the slope.

## Caution

- Always be aware of the type of data you're working with: random sample, non-random sample, or population.



## Caution

- Always be aware of the type of data you're working with: random sample, non-random sample, or population.
- Statistical inference, and the resulting p-values, are meaningless when you already have population data.

## Caution

- Always be aware of the type of data you're working with: random sample, non-random sample, or population.
- Statistical inference, and the resulting p-values, are meaningless when you already have population data.
- If you have a sample that is non-random (biased), inference on the results will be unreliable.

## Caution

- Always be aware of the type of data you're working with: random sample, non-random sample, or population.
- Statistical inference, and the resulting p-values, are meaningless when you already have population data.
- If you have a sample that is non-random (biased), inference on the results will be unreliable.
- The ultimate goal is to have independent observations.