

COVID-19 Model Containing Contact Tracing and Quarantine

1 Model Equations

$$S_{j+1} = S_j - S_j \alpha c \left(\frac{I_P + b_c I_C + b_a I_A}{N} \right) - (r \delta_E E_{j-4}) q (1 - \alpha) c (b_c (S_j + S_{j-1}) + (S_{j-2} + S_{j-3})) \frac{1}{N} \quad (1)$$

$$E_{j+1} = E_j - S_j \alpha c \left(\frac{I_P + b_c I_C + b_a I_A}{N} \right) - (r \delta_E E_{j-4}) q \alpha c (b_c (S_j + S_{j-1}) + (S_{j-2} + S_{j-3})) \frac{1}{N} - \delta_E E_j \quad (2)$$

$$I_{P_{j+1}} = I_{P_j} + r \delta_E E_j - \delta_{I_P} I_{P_j} \quad (3)$$

$$I_{C_{j+1}} = I_{C_j} + \delta_{I_P} I_{P_j} + \delta_Q Q_j - \delta_{I_C} I_{C_j} \quad (4)$$

$$I_{A_{j+1}} = I_{A_j} + (1 - r) \delta_E E_j - \delta_{I_A} I_{A_j} \quad (5)$$

$$Q_{j+1} = Q_j + (r \delta_E E_{j-4}) c q \alpha r (b_c (S_j + S_{j-1}) + (S_{j-2} + S_{j-3})) \frac{1}{N} - \delta_Q Q_j \quad (6)$$

$$Q_{A_{j+1}} = Q_{A_j} + (r \delta_E E_{j-4}) c q \alpha (1 - r) (b_c (S_j + S_{j-1}) + (S_{j-2} + S_{j-3})) \frac{1}{N} - \delta_{Q_A} Q_{A_j} \quad (7)$$

$$S_{Q_{j+1}} = S_{Q_j} + (r \delta_E E_{j-4}) q (1 - \alpha) c (b_c (S_j + S_{j-1}) + (S_{j-2} + S_{j-3})) \frac{1}{N} - \delta_{S_Q} S_{Q_j} \quad (8)$$

$$R_{j+1} = R_j + \delta_{I_C} I_{C_j} + \delta_{I_A} I_{A_j} + \delta_{Q_A} Q_{A_j} \quad (9)$$

2 Parameters

- c : Contact rate between an infectious individual and susceptible individuals
- α : Probability of transmission given a contact
- τ : Days for covid-19 test results
- r : Probability that an exposed individual becomes presymptomatic and move to I_P
- q : Contact tracing effectiveness rate
- b_A : Decrease in asymptomatic infectivity
- b_C : Decrease in clinical infectivity
- δ_E : Mean time spent in Exposed
- δ_{I_P} : Mean time spent in Infected Pre-Clinical
- δ_{I_C} : Mean time spent in Infected Clinical

- δ_{I_A} : Mean time spent in Infected asymptomatic
- δ_Q : Mean time spent in Q (Quarantine)
- δ_{Q_A} : Mean time spent in Q_A
- δ_{S_Q} : Mean time spent in S_Q