

(Rajshahi, Chapainawabganj, Naogaon and Natore)

Region	Turmeric																																																				Duration: 270–300 days							
Months	April					May					June					July					August					September					October					November					December					January				February										
Std. Week / Normal	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8												
Rainfall (mm)	36	42	51	78	72	66	71	75	72	52	57	64	53	57	54	59	52	44	35	28	28	20	12	8	6	5	6	8	10	12	16	18	14	10	8	5	4	3	4	5	8	12	18	20	22	24	18	18												
Max. Temp. (°C)	34.5	34.2	34.1	33.6	33.2	32.8	32.6	32.4	32.3	32.1	32	32.1	32.4	32.8	33	33.2	32	31.5	31	30.5	30.1	29.8	29.6	29.4	29.5	29.8	30	30.4	29.5	29.7	30.4	30.8	31.2	31.8	32.2	32.5	32.8	33	33.2	33.3	33.5	33.6	33.8	34	33.5	33	33	33.2												
Min. Temp. (°C)	22.5	24	25	26	26.5	26.5	26.5	26	25.8	25.6	25	25.5	25.4	25.3	25.1	24.8	24.5	22.1	21.5	21.1	20.5	20.2	20	19.8	19	18.5	18	18.5	19	19.5	20	20.5	21	21.5	22	22.5	23	23.5	24	24.5	25	25.5	26	26.5	27	27.5	28	28.5	29											
Mean Temp. (°C)	28.5	29.1	29.6	29.8	29.9	29.6	29.6	29.2	29	28.9	28.8	28.8	28.9	28.9	28.9	28.9	27.1	26.5	26.1	25.5	25.1	24.9	24.7	24.2	24	23.9	24.2	24.7	24.5	24.9	25.4	25.9	26.4	26.9	27.4	27.8	28.1	28.5	28.9	29.1	29.5	29.8	30.1	30.5	30.5	30.5	30.8	31.1												
RHmax (%)	95	95	96	96	96	97	97	96	96	96	96	96	96	96	96	95	94	93	92	91	90	89	88	87	86	86	87	88	86	87	88	89	90	91	92	93	94	95	95	95	95	95	95	95	95	95	95	95												
RHmin (%)	52	58	65	72	74	75	74	73	72	71	70	69	68	67	66	64	58	55	52	49	46	43	40	38	36	35	36	38	36	36	38	40	43	47	52	56	60	64	67	70	72	75	77	78	80	82	84	86												
RHmean (%)	73.5	76.5	80.5	84	85	86	85.5	84.5	84	83.5	83	82.5	82	81.5	81	79.5	76	74	72	70	68	66	64	62.5	61	60.5	61.5	63	61	61.5	63	64.5	66.5	69	72	74.5	77	79.5	81	82.5	83.5	85	86	86.5	87.5	88.5	89.5	90.5												
SShr (hrs)	48	45	40	35	31	30	30	30	32	34	33	34	36	35	37	38	40	42	44	47	49	51	53	55	56	57	55	53	57	56	55	53	50	47	44	40	37	33	30	28	28	30	32	34	35	36	38	40												
WD (deg)	135	132	130	128	127	130	134	138	140	138	136	133	131	129	128	126	125	124	122	120	118	115	112	110	108	107	108	112	118	122	126	130	134	138	140	142	145	148	150	152	154	156	158	160	162	164	166	168												
WS (km/hr)	7	6.8	6.5				6	5.8	5.6	5.5	5.4	5.2	5.1	5	4.8	4.7	4.6	4.4	4.2	4	3.8	3.6	3.5	3.5	3.6	3.8	4	4.2	4.5	4.8	5.2	5.6	6	6.4	6.8	7.2	7.5	7.8	8	8.2	8.5	8.7	8.9	9	9.1	9	8.8	8.6	8.4											
Crop growth stages																																																												
	Land preparation & planting (April–May)					Germination & establishment (May–June)					Vegetative growth (June–August)					Rhizome initiation (August–September)					Rhizome enlargement (October–January)					Maturity & harvesting (January–February)																																		
Congenial Weather Conditions for Pests & Diseases																																																												
Rhizome rot High soil moisture, temperature 25–28°C, high relative humidity (>80%), prolonged cloudy/rainy weather																																																												
Leaf spot Temperature 25–32°C, high relative humidity, cloudy weather, frequent rain																																																												
Leaf blotch Temperature 25–32°C, high relative humidity, cloudy weather																																																												
Shoot/stem borer Minimum temperature >20°C, maximum 29–34°C, higher morning RH (>84%), dry weather																																																												
Leaf roller Maximum temperature 32–33°C, relative humidity 92–95%																																																												
Other insects (e.g. scale, mealy bug) High temperature (>32°C), high relative humidity (80–90%), intermittent rain																																																												
Weather Warning																																																												
Heavy rain >50 mm/day: drainage/field inspection; >100 mm/day: high risk of waterlogging and rhizome disease																																																												
Wet spell >25 mm for 3 days or >50 mm for 4 days: drainage and disease surveillance																																																												
High wind >40–50 km/hr: risk of lodging/leaf damage; secure vulnerable fields																																																												
Low temperature Minimum <10°C: unusual cold stress risk, especially for young crop																																																												
High temperature Maximum >35°C: moisture-stress risk; prioritize irrigation/mulching where needed																																																												
Variety production information from BARI																																																												
BARI Holud-1 (Dimla) 270–280 days; planting late March–early April; harvest late January–early March; 28–32 t/ha																																																												
BARI Holud-2 (Sinduri) 270–280 days; planting late March–early April; harvest late January–early March; 25–28 t/ha																																																												
BARI Holud-3 270–290 days; planting late March–early April; harvest late January–early March; 25–30 t/ha																																																												
BARI Holud-4 270–290 days; planting mid-April–early May; harvest mid/late February; 28–30 t/ha																																																												
BARI Holud-5 270–300 days; planting mid-April–early May; harvest mid/late February; 18–20 t/ha																																																												